

# Respiratory Care: Artificial Airways, Mechanical Ventilation, and Chest Tubes

## HOW TO RECEIVE CREDIT

- Read the enclosed course.
- Complete the questions at the end of the course.
- Return your completed Evaluation to NetCE by mail or fax, or complete online at [www.NetCE.com](http://www.NetCE.com). (If you are a Florida nurse, please return the included Answer Sheet/Evaluation.) Your postmark or facsimile date will be used as your completion date.
- Receive your Certificate(s) of Completion by mail, fax, or email.

## Faculty

Jane C. Norman, RN, MSN, CNE, PhD, received her undergraduate education at the University of Tennessee, Knoxville campus. There she completed a double major in Sociology and English. She completed an Associate of Science in Nursing at the University of Tennessee, Nashville campus and began her nursing career at Vanderbilt University Medical Center. Jane received her Masters in Medical-Surgical Nursing from Vanderbilt University. In 1978, she took her first faculty position and served as program director for an associate degree program. In 1982, she received her PhD in Higher Education Administration from Peabody College of Vanderbilt University. In 1988, Dr. Norman took a position at Tennessee State University. There she has achieved tenure and full professor status. She is a member of Sigma Theta Tau National Nursing Honors Society. In 2005, she began her current position as Director of the Masters of Science in Nursing Program.

## Faculty Disclosure

Contributing faculty, Jane C. Norman, RN, MSN, CNE, PhD, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

## Division Planner

Mary Franks, MSN, APRN, FNP-C

## Senior Director of Development and Academic Affairs

Sarah Campbell

## Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

## Audience

This course is designed for nurses working in critical care and general and specialty medical-surgical units in which patients require assistance maintaining a patent airway and respiration.

## Accreditations & Approvals



JOINTLY ACCREDITED PROVIDER<sup>™</sup>  
INTERPROFESSIONAL CONTINUING EDUCATION

In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

## Designations of Credit

NetCE designates this continuing education activity for 3 ANCC contact hours.

NetCE designates this continuing education activity for 3.6 hours for Alabama nurses.

AACN Synergy CERP Category A.

### Individual State Nursing Approvals

In addition to states that accept ANCC, NetCE is approved as a provider of continuing education in nursing by: Alabama, Provider #ABNP0353 (valid through 07/29/2025); Arkansas, Provider #50-2405; California, BRN Provider #CEP9784; California, LVN Provider #V10662; California, PT Provider #V10842; District of Columbia, Provider #50-2405; Florida, Provider #50-2405; Georgia, Provider #50-2405; Kentucky, Provider #7-0054 (valid through 12/31/2025); South Carolina, Provider #50-2405; West Virginia, RN and APRN Provider #50-2405.

### Special Approvals

This activity is designed to comply with the requirements of California Assembly Bill 241, Implicit Bias.

### About the Sponsor

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

### Disclosure Statement

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

### Course Objective

The purpose of this course is to reinforce nurses' knowledge and skills related to the care of patients with artificial airways and/or chest tubes in order to improve outcomes and patient quality of life.

### Learning Objectives

Upon completion of this course, you should be able to:

1. Outline the types and potential complications of artificial airways.
2. Discuss nursing management and patient education for patients with mechanical ventilation.
3. Describe available chest tubes and nursing care of patients who require them.



Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the course material for better application to your daily practice.

---

## INTRODUCTION

---

Artificial airways are used for a variety of patients who require assistance in maintaining a patent airway. The indications for placement of an artificial airway are to relieve obstruction, facilitate suctioning of the lower airway, allow for mechanical ventilation, and/or prevent aspiration. Artificial airways include endotracheal, tracheotomy, and laryngectomy tubes [1].

Mechanical ventilation may be necessary for any patients for whom the process of ventilation has been significantly altered. This alteration may be caused by any condition that prevents the patient from maintaining normal oxygen and carbon dioxide (CO<sub>2</sub>) levels in the blood [2; 3]. Mechanical ventilation may also be indicated for patients with an inability to protect their airway.

In conjunction with artificial airways, some patients with respiratory conditions will require chest tubes. Chest tubes re-establish normal intrapleural pressure in the event of trauma, pleural collection or effusion, or postsurgical drainage.

---

## TYPES OF ARTIFICIAL AIRWAYS

---

### ENDOTRACHEAL TUBES

Endotracheal intubation (by nasal or oral routes) is one of the most widely used methods to institute and maintain an open airway. Placement of an endotracheal tube requires the use of a laryngoscope to visualize the area and facilitate passage of the tube into the trachea. Prior to intubation, whether oral or nasal, the airway should be assessed following the mnemonic LEMON to ensure safe intubation [10]:

- Look for any external signs of trauma or other potential obstructing items, such as a large tongue, neck mass, or dentures.
- Evaluate the 3-3-2 rule. Anything less than three fingers between the incisors, three fingers between the hyoid bone and the mental protuberance, or two fingers between the hyoid bone and the thyroid cartilage (Adam's apple) may be a sign of a difficult airway.
- Mallampati class of 3 or greater predicts a difficult airway.
- Obstruction or obesity can restrict vocal cord visualization.
- Neck mobility or lack thereof contributes to difficult airways and may result in difficulty passing the tube into the correct space.

Endotracheal tubes are single-use and constructed of polyvinylchloride (PVC) [11]. They range in size from 2.5 mm to 8 mm in diameter. The tubes may also be uncuffed or cuffed, the latter of which includes an inflatable balloon to help prevent gastric contents from entering the trachea and to ensure a specific positive pressure is maintained. Uncuffed tubes are generally an option for pediatric patients; most adult patients will receive a cuffed tube. A tube that is too large can damage the airway structure and mucosal lining; a tube that is too small will not allow adequate ventilation. The tube should be inserted gently to avoid structural damage. Usually, adult men will require size 8 and adult women size 7, but this can vary depending on the size of the individual and the patient's specific anatomy. The sizing for pediatric patients is (age/4) +4 for uncuffed tubes; if a cuffed tube is used, size down one-half size [12; 13]. The oral route is most often employed when emergency intubation is required because of the relative ease of insertion.

Generally, with orotracheal intubations, depending on the situation, medications to block neuromuscular actions and provide adequate sedation are used. This also allows for the prevention of vomiting and aspiration of gastric contents [14; 15]. In addition, the airway should be cleared of any debris (e.g., dentures), if applicable.

The proper procedure for orotracheal intubation consists first of preoxygenating the patient (if possible) and administering medications, if indicated, often simultaneously with preoxygenation. The patient is then placed in the “sniffing” position, with the patient in the supine position, head elevated, and neck extended. Intubation of the trachea is then achieved with direct visualization using a laryngoscope. The balloon of the tube is inflated after it is properly placed 3-4 cm past the vocal cords, then the tube is properly secured.

Nasotracheal intubations are also used in a variety of settings. It involves a nasal trumpet along with the endotracheal tube. The nasal trumpet is well lubricated and then inserted through the nare into the nasopharynx. The trumpet dilates the nasal cavity prior to inserting the endotracheal tube. After the tube is passed successfully through the nasopharynx, a laryngoscope is used to visualize the vocal cords for successful passage into the trachea. The cuff on the tube is then inflated to the proper measurement and the tube secured [15; 16; 17]. Preoxygenation of the patient is ideal but not always available in emergent situations [17]. Proper placement is verified by end-tidal carbon dioxide measurements, capnography, chest x-ray, ultrasound, and clinical assessment [14; 15; 17].

Proper sizing of nasal trumpets is important to avoid trauma or poor airway management. The size of the nasal trumpet is chosen particularly based on length rather than diameter, as the correct length ensures the tip sits just superior to the epiglottis. A measurement is taken from the lateral aspect of the nares to the tragus of the ear on the same side to determine the correct length [18]. Nasal trumpets are typically made of soft rubber or plastic.

While nasotracheal intubation does have advantages, such as less sedation required, there are many disadvantages and potential complications as well, including nasal damage, sinusitis, sepsis, and local abscess. Nasopharyngeal airways also pose increased difficulty with pulmonary toilet and have greater airway resistance due to the use of narrower tubes

[19; 20]. Thus, the use of long-term nasopharyngeal airways is limited. Nasotracheal intubation may be indicated in the following situations [21]:

- Impending airway compromise
- For conscious patients in order to avoid loss of airway protection reflexes and less gagging
- Intraoral and oropharyngeal surgery
- Complex intraoral procedures involving mandibular reconstruction
- Rigid laryngoscopy
- Dental surgery
- Maxillofacial or orthognathic surgery

Absolute contraindications for nasotracheal intubation include [21]:

- Suspected epiglottitis
- Midface instability
- Previous history of skull base fractures
- Any known bleeding disorder that could predispose the patient to severe epistaxis
- Choanal atresia
- Anterior skull base fractures, which may result in passage of the tube intracranially

Relative contraindications include:

- Obstruction of the nasal airway (e.g., large nasal polyps, foreign bodies)
- Recent nasal surgery
- History of frequent episodes of epistaxis

Anesthesia uses nasotracheal intubations frequently, dependent on the situation or type of procedure being performed. As the endotracheal tube is introduced into the trachea through the nasal cavity, the tube is easier to stabilize through the nasal pathways due to its small diameter compared with the oral cavity. Nasotracheal intubation is the preferred method for anesthesiologists and surgeons who perform surgeries in the head and neck region, especially oropharyngeal, dental, and maxillofacial surgeries, as this method helps to improve the field of view and access for surgery [22].

## Nursing Care

Immediately after intubation, the lung fields should be auscultated and an x-ray ordered to determine if the tube is placed properly. Because the right bronchus is straighter than the left, the tube may be improperly inserted into the right mainstream bronchus, permitting aeration of the right lung only. If the end of the endotracheal tube rests on the carina (i.e., the area where the trachea bifurcates into the left bronchus and right bronchus), partial or complete obstruction results. Inadvertent slipping of the tube into the esophagus will result in gastric dilatation [1; 4].

Endotracheal tubes can also provide a means of suctioning secretions from the lower respiratory tract. Sterile technique is used, and the tube should be stabilized to prevent inadvertent removal (extubation) or change in placement. When the patient is intubated nasally, tape is applied at the insertion site. Stabilizing the oral endotracheal tube involves placement of an oral airway to prevent the patient from biting down and obstructing the tube. It is important to avoid applying excessive tape at the mouth, which could make inspection of the oral cavity difficult [1; 4]. Plastic devices may also be used rather than tape to secure the tube. Meticulous oral and skin hygiene are important for these patients. The position of the orotracheal tube should be changed daily and the tape replaced at the same time. Care should be taken to avoid skin breakdown from irritation by the tape. The patient's skin should be assessed daily when tape is changed; a rash may indicate an allergic reaction. The use of nonallergenic tape or plastic devices may then be indicated, particularly for patients who are likely to have skin breakdown and/or poor wound healing (e.g., those with diabetes).

Patients with endotracheal tubes should be assessed daily for complications involving the nose, mouth, pharynx, sinuses, and ears. Nasotracheal tubes can obstruct the Eustachian tube, causing otitis media. Damage to the nasal mucosa may occur from pressure or irritation [1; 4]. Other possible complications include damage to the vocal cords, laryngeal edema, and laryngeal ulcers, caused by traumatic insertion, improper stabilization, and/or unneces-

sary manipulation of the tube. Laryngeal ulcers occur more frequently with oral intubation than with nasal intubation. This is because it is difficult to anchor the endotracheal tube in the mouth, and an unanchored tube exerts pressure on the posterior rim of the glottis [1].

The length of time an endotracheal tube is left in place varies with agency policy and the physician. If prolonged intubation (longer than 10 to 14 days) is anticipated, a tracheotomy is usually performed, because longer endotracheal tube placements are associated with greater risks of complications [1].

## Extubation

Patients should be adequately ventilated, oxygenated, and suctioned before extubation is attempted. The nurse deflates the cuff and again suctions the patient to remove secretions that may have accumulated. The tube is removed, and oxygen is administered. An individual qualified to reintubate the patient should be present, and an intubating tray should be ready at the bedside. Complications that may follow tube removal include laryngeal edema and laryngospasm. The patient should be observed for signs of complications, including labored breathing, use of accessory muscles, and stridor. Laryngospasm and laryngeal edema can result in upper airway obstruction and may require the administration of parenteral corticosteroids and reintubation.

## TRACHEOTOMY TUBES

Other methods of establishing an artificial airway include tracheotomy and cricothyroidotomy. The tracheotomy was one of the earliest surgical procedures recorded, dating back to 3600 B.C.E. in Ancient Egypt. Today, tracheotomies can be temporary or permanent. The Eastern Association of Surgical Trauma (EAST) recommends early tracheostomy (within three to seven days after intubation) for patients who have severe closed-head injuries or for patients who require prolonged ventilatory support. In non-trauma-type patients with failed ventilator weaning for a variety of different reasons, tracheostomy at post-intubation between days five and seven has been recommended by various professional organizations [23].



Tracheotomy tubes are placed during these procedures. These short, curved tubes have a flange that assists in stabilization of the device on the neck. A variety of tubes are available, including metal or disposable silicone types in several sizes. Most tubes have an inner cannula to facilitate cleansing and are packaged with an obturator, which is used to minimize trauma when the tube is inserted [5; 6].

The terms “tracheotomy tube,” “tracheostomy tube,” and “laryngectomy tube” are often used interchangeably. Actually, a laryngectomy tube is shorter and consists of an obturator, inner cannula, and outer cannula. These tubes are usually placed following a total laryngectomy to provide a route for removal of secretions and to maintain a patent airway [6].

### **Tracheotomy and Laryngectomy Tube Care**

Tracheotomy care is performed to minimize bacterial contamination and to decrease the possibility of obstruction by secretions. Patients with tracheotomy tubes require suctioning when they experience difficulty breathing, noisy/increased respirations, or cyanosis. Routine care consists of cleaning of the inner cannula and the area around the stoma. The frequency of cleaning varies depending on the amount of secretions present—it may be necessary as frequently as every 30 minutes or only once per shift. Some experts advocate cleaning the inner cannula as often as necessary, while others believe that proper humidification and suctioning negate the need for frequent cleaning [5]. It is important to remember that more frequent cleaning may increase the risk of infection.

When caring for patients with a tracheotomy or laryngectomy tube, it is wise to have another sterile tube at the bedside in case of accidental decannulation or obstruction of the tube. A clearly labeled obturator should also be in the room to be used in cases of emergency reintubation. Some prefer to keep sterile forceps at the bedside as well, because decannulation in the period immediately after tube insertion may result in closure of the stoma. Scissors

should also be at the bedside to cut the tracheotomy tube ties if they become too tight or if the tube is partially dislodged by coughing [6].

Complications can arise with tracheotomies. Early complications include wound infections, frequently related to leaking secretions from the stoma. An abscess can form at the site of the stoma as well. Potentially major issues in the early postoperative period include the development of pneumonia and obstruction of the tube with blood or mucus. Deep infections can be associated with tracheotomy cannula holders being too tight, causing pressure injuries to the site. Inadvertent dislodgement of the tracheostomy tube can be the result of improper sizing or lack of appropriate securement [23]. Possible late complications include necrosis due to overinflation of the tracheostomy tube cuff and stenosis related to improper tracheostomy tube sizing. It is important to remember that the tracheostomy cuff pressure should be no more than 20 cm H<sub>2</sub>O [23].

### **Extubation**

Prior to removal of an endotracheal tube, the patient should be ventilated, oxygenated, and suctioned to remove tracheal and pharyngeal secretions. The cuff is deflated, and the patient is again suctioned. The tube is then removed. The patient is then placed on supplemental oxygen and carefully observed over the next several hours. The choice of supplemental oxygen by nasal cannula, high-flow oxygen mask, continuous positive airway pressure (CPAP), or bilevel positive airway pressure (BIPAP) is dependent on patient factors. CPAP or BIPAP should be considered for patients at high risk of re-intubation. An extubation failure is classified as a patient who is extubated but re-intubated within 48 hours. This occurs in 10% and 20% of patients undergoing extubation [24].

If the patient has a tracheotomy a dressing is placed over the stoma. Normally, the opening of a tracheostomy closes within five days [5; 6].

Fenestrated tracheotomy tubes are used in some cases prior to extubation of the patient. This tube has an opening, or fenestration, in the outer canula; it may or may not have a cuff. If the fenestrated tube is cuffed, the cuff should be deflated to allow the patient to breathe around, as well as through, the tube and adjust gradually to its removal. It also allows healthcare professionals to assess how the patient will tolerate removal. Covering the tracheotomy tube enables the patient to talk, to breathe normally through the upper airway, and to expectorate secretions [5; 6].

Another way of preparing the patient for permanent removal of a tracheotomy tube is by applying a tracheotomy button. The button extends from the tracheotomy opening to just inside the tracheal wall. If the tracheotomy tube has a cuff, the cuff is deflated before the tracheotomy button is placed to allow the patient to breathe around the tube. To determine readiness for decannulation, a tracheotomy button is most effective with a fenestrated tube. Placement of this button facilitates talking, coughing, and normal breathing through the upper airway. If the patient has difficulty breathing or expelling secretions, the button is removed, and the patient breathes through the tracheotomy tube [6].

---

## POSSIBLE COMPLICATIONS OF ARTIFICIAL AIRWAYS

---

The insertion of an artificial airway can result in a variety of complications. Because inspired air bypasses the nose, air should be humidified to prevent drying the mucosa of the lower respiratory tract. Patients with artificial airways are also more vulnerable to the development of infection as a result of altered ciliary function, trauma to the mucosa caused by suctioning and intubation, and colonization of the airway with bacteria. Organisms commonly isolated from the respiratory tract in these patients include *Staphylococcus aureus* and

*Pseudomonas* spp. [5; 7; 24]. Guidelines for the treatment of ventilator-associated pneumonia are available online at [https://www.idsociety.org/practice-guideline/hap\\_vap](https://www.idsociety.org/practice-guideline/hap_vap) [25].



According to the American Association for Respiratory Care, when the upper airway is bypassed during invasive mechanical ventilation, humidification is necessary to prevent hypothermia, disruption of the airway epithelium, bronchospasm, atelectasis, and airway obstruction. When the upper airway is not bypassed, such as in noninvasive mechanical ventilation, active humidification is highly suggested to improve comfort. (<http://rc.rcjournal.com/content/57/5/782>. Last accessed October 16, 2023.)

**Level of Evidence:** Expert Opinion/Consensus Statement

Excessive pressure on the trachea from the endotracheal tube cuff can decrease blood supply to the area and cause tissue necrosis. This may lead to tracheal stenosis or a tracheoesophageal fistula.

Another possible complication is herniation of the cuff over the end of the tube, resulting in partial or complete airway obstruction. Signs include a significant air leak through the stoma, mouth, or nose; the sounding of the high-pressure alarm on the ventilator; and/or obstruction when attempting to suction the patient.

An underinflated cuff may be caused by instillation of an insufficient amount of air or a ruptured cuff. An air leak may be detected around the stoma, nose, or mouth, with the ventilator indicating a decrease in the expired volume of air by sounding the low-pressure alarm [6].

Obstruction of artificial airways may also occur because of accumulation of secretions or a kink in the endotracheal tube. This should be corrected immediately to prevent asphyxiation.

---

## CARE OF ARTIFICIAL AIRWAYS

---

Placement of an artificial airway can be a source of anxiety and apprehension for the patient and family members. One major source of anxiety is an impaired ability to communicate. However, aside from patients who have undergone a total laryngectomy, this problem is temporary. To help allay patients' fear, nurses should provide an alternative means of communication. A white board or a pad of paper and a pencil should be available. The call light should be within reach of the patient at all times and should be answered promptly. The intercom at the nurses' station should be marked to indicate that the individual is unable to speak. Some patients who cannot communicate by writing may use gesturing as an alternative method [8]. Family members should always be included when providing information and reassurance.

Patients with artificial airways also have an impaired ability to remove secretions effectively. The artificial airway prevents the glottis from closing, which impedes the generation of the intrathoracic pressure necessary to dislodge secretions. Thus, cough is less effective. The artificial airway also decreases the effectiveness of the mucociliary mechanism. Secretions that accumulate in the airway may cause the patient to fear suffocation. When the patient is unable to cough and expel secretions, suctioning is necessary. During this procedure, the patient will experience discomfort and a feeling of breathlessness. Nurses should attempt to comfort the patient by providing an explanation of the artificial airway and the suctioning procedure prior to their use, if possible [8].

Obviously, the placement of an artificial airway will result in a change in patients' body image. This not only affects the way patients view themselves but the way they are viewed by family members and significant others. Hospitalization, by its very nature, puts patients in a dependent role, and having an artificial airway compounds the feeling of lost autonomy.

Patients should be encouraged to participate actively in their own care as much as possible. For most patients, an artificial airway is temporary, and they should be informed that the airway will be removed after they are able to ventilate adequately [8].

---

## MECHANICAL VENTILATION

---

### TYPES OF MECHANICAL VENTILATION

Although there are other types of mechanical ventilators, such as negative pressure and CPAP devices, what is commonly referred to as a ventilator is a conventional positive pressure machine. These work by creating higher than atmospheric pressure to push air into the patient's lungs. Positive-pressure mechanical ventilators are pressure-cycled, volume-cycled, or a combination of both. Other, less common types include airway pressure release ventilation (APRV) and bilevel ventilation; discussion of these types is outside of the scope of this course [3; 9; 26].

With pressure-cycled or pressure-limited ventilation, air is delivered to the lungs until a predetermined pressure is reached, then inspiration is terminated. Pressure support cycles are typically used when weaning a patient from mechanical ventilation. The disadvantage of this is that the tidal volume delivered to the patient can vary with each inspiration, depending on his or her airway [3; 9]. The main advantage is the lungs are protected from overinflation due to excessive pressure.

In volume-cycled or volume-limited ventilation, a set amount of air is delivered for each inspiration. The advantage of this form of ventilation is that the tidal volume remains consistent and the pressure necessary to deliver the volume varies, dependent on the patient's lung compliance. Volume-cycled is the most frequently used type of ventilation for adult patients [9].



## **PATTERNS OF MECHANICAL VENTILATION**

There are four common ventilation patterns: assist-control (A/C), intermittent mandatory ventilation (IMV), synchronized intermittent mandatory ventilation (SIMV), and positive end-expiratory pressure (PEEP). In assist-control mode, the inspiratory phase is initiated by the patient. However, modern ventilators have a built-in safety feature to ensure that the machine will deliver the volume at the ordered rate if the patient becomes apneic. In controlled ventilation, the machine has complete control of the patient's rate and depth of ventilation [2; 3].

IMV delivers a preset tidal volume at a specific rate while also providing a continuous flow of air for spontaneous breaths. IMV was originally introduced for the purpose of gradually weaning individuals from ventilators, but the use of this type of ventilation has expanded, and it has become popular for patients who require mechanical ventilation but are able to initiate some inspiratory effort on their own, such as patients with chronic obstructive pulmonary disease. IMV cannot be used for individuals who are apneic (e.g., those who have suffered damage to the brain stem). An advantage of IMV ventilation is that it allows the patient to use respiratory muscles, which prevents atrophy [2; 3; 5; 6].

With SIMV, air is delivered by the ventilator in synchronization with the patient's own ventilation efforts. If the SIMV is set at a rate of 10, the patient is assisted 10 times per minute [6].

PEEP prevents the collapse of the airways and alveoli at the end of expiration, facilitating the diffusion of more oxygen from the alveoli into the pulmonary capillaries. This allows for a reduction in inspired oxygen concentrations, because oxygen can diffuse during both inspiration and expiration. The setting for PEEP is usually in the range of 1–15 cm. However, levels greater than 40 cm have been used when treating patients with severe hypoxemia, as in those with acute respiratory distress syndrome [2; 6].

Physiologic changes occur with the administration of PEEP. PEEP causes an increase in intrathoracic pressure, which decreases the venous return of blood to the heart. The baroreceptors in the thoracic aorta interpret the decreased venous return as hypovolemia and stimulate an increase in production of antidiuretic hormone (ADH). Increased ADH can lead to the development of hypervolemia, which results in decreased cardiac output. Another potential problem encountered is the development of a pneumothorax [2; 6].

## **NURSING CARE OF PATIENTS WHO ARE MECHANICALLY VENTILATED**

Nurses should take steps to ensure the comfort and safety of mechanically ventilated patients. Accidental disconnection of the patient from the ventilator should be prevented. A warning should be placed on all ventilators and on the nursing care plan reminding healthcare personnel to leave the ventilator alarm on at all times, even during suctioning. The sound of the alarm is a minor nuisance compared to the damage that may occur if someone fails to turn the alarm back on (in some cases due to alarm fatigue) [2; 5].

A bag-valve mask (e.g., Ambu bag) should be kept at the patient's bedside to be used in the event of a power failure or ventilator malfunction. If the ventilator is not working properly, do not waste time trying to identify the mechanical difficulty. Instead, use the bag-valve mask to ventilate the patient while another staff member calls the respiratory maintenance technician [5].

Patients who are mechanically ventilated often require monitoring of central venous pressure or pulmonary capillary wedge pressures. The positive pressure exerted by the ventilator will alter these readings. It is important that they always be taken either with the patient connected to or disconnected from the ventilator in order to maintain consistency [2; 5; 26].

As a rule, respiratory therapists are responsible for checking the function of the ventilator at regular intervals. However, nurses should also be familiar with the machine and how it functions [6; 26]. The pressure indicator shows the amount of force required to ventilate the patient's lungs. If the amount of pressure required increases or decreases significantly, assess the situation to determine the cause, if possible. If any alarm sounds on the ventilator, it is crucial to respond immediately. If the reason for the alarm cannot be readily ascertained and rectified, the patient should be disconnected from the ventilator and ventilated manually with a bag-valve mask until the ventilator can be replaced or fixed. Because the alarms can be frightening to patients and any others in the room, reassure them that the patient can be adequately oxygenated [6].

The settings on the ventilator should be checked hourly. This includes making sure the tidal volume, respiratory rate, fraction of inspired oxygen ( $\text{FiO}_2$ ), and other parameters are set as ordered. Also, the ventilator tubing should be checked for water condensation; if any is found, the tubing should be disconnected and the water emptied. This will prevent impairment of ventilation or accidental aspiration of water into the trachea [2; 6; 13; 26].

In addition to checking the ventilator, carefully assess the patient for overt clinical signs of hypoxia, hypocapnia, or hypercapnia. Arterial blood gas values should be obtained at regular intervals. Monitoring these values is particularly important when ventilator settings are changed, as when the  $\text{FiO}_2$ , IMV, or PEEP settings are decreased. Assess the patient every hour to ensure that both lungs are aerated by auscultating lung sounds and observing for symmetrical expansion of the chest. Because hyperventilation can occur, patients should be monitored for signs of respiratory alkalosis [2; 6; 13; 26].

A major responsibility is careful monitoring of the patient's vital signs. Because the application of PEEP can result in a decrease in cardiac output, particular attention should be paid to blood pressure, pulse, and urine output. Any time the setting for PEEP is altered, the patient should be re-evaluated. Increasing the amount of PEEP can further compromise venous return, whereas decreasing could result in cardiac overload and pulmonary edema. Often, patients with PEEP have a central venous pressure (CVP) line, a peripherally inserted central venous catheter, or a pulmonary artery catheter, which allows for more accurate assessment of physiologic parameters. Because of the increased positive pressure exerted with PEEP, the patient should be observed for signs and symptoms of pneumothorax [2; 6; 13; 26].

Nursing care of the patient who is mechanically ventilated also includes position change and skin assessment every hour, putting joints through active or passive range of motion exercises every eight hours, and frequent oral care. Patients who are comatose will also require eye care, including use of artificial tears to keep the eyes moist, removal of crusts on lashes and lids with sterile saline or sterile water-soluble lubricants, and/or protection from corneal drying, irritation, or trauma by taping the eyelids closed [2; 8; 13; 26].

Patients will occasionally "fight" or "buck" the ventilator, also referred to as "being out of phase with the ventilator." In these cases, the patient attempts to actively exhale while the ventilator is delivering the inspired volume. This conflict should be corrected because it results in decreased volume delivery and excessively high airway pressures. A bag-valve mask may be used to increase volume delivery gradually and decrease the respiratory rate. Altered inspiratory flow rates may be ordered for the ventilator to correct the situation, or medication may be prescribed to sedate the patient and allow the ventilator to function more efficiently [2; 8; 13; 26].

Another possible complication of prolonged mechanical ventilation is the development of stress-related mucosal damage. This is a broad term that encompasses a spectrum of inflammatory damage to the upper gastrointestinal tract, ranging from superficial lesions to significant gastrointestinal bleeding (e.g., a stress ulcer). The mechanism of the formation of stress ulcers is not entirely understood, but it is believed to be related to a combination of hypersecretion of hydrochloric acid and decreased resistance of the gastric mucosal lining. The resultant gastric irritation may lead to bleeding, so nurses should assess stool and nasogastric drainage for fresh and occult blood. Many ventilated patients receive a proton pump inhibitor for stress ulcer prophylaxis [27].

Gastric dilatation may also occur from introduction of air into the esophagus. This may necessitate the insertion of a nasogastric tube [2; 8].

### **PSYCHOSOCIAL CONSIDERATIONS AND PATIENT EDUCATION**

Patients receiving mechanical ventilation require frequent attention and care in an environment with a lot of activity and a high noise level. This may result in sensory overload, so it is important that nursing care be organized to give the patient periods of uninterrupted rest.

Mechanical ventilation can be frightening for patients and their families. Staff members are often so familiar with their routines and equipment that they discount the potential impact on patients and their families. It is important to take time to explain equipment, alarms, and lights. The patient and significant others should be assured that a nurse is always close by and will respond promptly to the patient's calls and alarms. Patients should not be left alone in the room until they appear relatively comfortable with the situation. Be careful not to become so involved with monitoring the equipment and performing the procedures that the patient is forgotten; patients and family members should be given time to express fears and concerns. These patients are acutely ill and may have realistic fears about being dependent on the machine and the possibility of dying [2; 6].

Mechanical ventilation limits patient mobility, and patients are confined to their rooms with limited contact with the larger world, which may lead to disorientation. Make a point to interact with the patient frequently and orient him or her to day, time, and place to minimize confusion and prevent anxiety. Lighting can also be controlled to create a day-night change. It can be helpful to keep a large calendar in the room so the patient can remain knowledgeable about passing time. Family members may also bring pictures or other articles to the hospital to personalize the environment. Patients should be encouraged to become actively involved in their care to the extent they are able in order to maintain a sense of control over their lives [2; 6; 19]. In one study, some patients reported being able to remember things that were said or done while they were sedated and ventilated [28].

When patients require indefinite mechanical ventilation after discharge, they and their family members/caregivers will require instruction. Ancillary health-care providers may also be needed in the home. The nurse can facilitate proper care by coordinating activities among different departments, such as respiratory therapy, social service, public health nurses, and any other healthcare team members involved [2; 5].

### **WEANING THE PATIENT FROM THE VENTILATOR**

Weaning the patient from mechanical ventilation involves reducing physiologic and psychologic dependence on the ventilator. Before weaning, the patient's arterial blood gas values on a liter flow of oxygen should be greater than 40% to 50%.

If the patient is receiving PEEP, it should be no higher than 5 cm; otherwise, it will interfere with the patient's independent inspiratory efforts. Vital capacity should be greater than 15 mL/kg of body weight, as this indicates that the patient is able to move enough air. This measurement can be obtained using a respirometer attached to the end of the patient's endotracheal or tracheotomy tube.

Maximum inspiratory effort should also be assessed to determine whether the patient's chest expansion is sufficient to produce negative alveolar pressure and stimulate a deep inspiration. The greater the patient's ability to inhale, the more negative the inspiratory pressure will be. This is assessed by attaching a pressure gauge to the end of the endotracheal or tracheotomy tube and instructing the patient to take in a deep breath. The patient is considered to be ready for weaning with a pressure of -25 to -30 cm. Respiratory rate and minute volume measurements should be taken. The respiratory rate should be at least 12 breaths per minute, but no greater than 20 breaths per minute [6; 8; 13]. Additional criteria for weaning include a stable chest wall, an acceptable chest x-ray, adequate cardiac output, normal body temperature, adequate nutritional status, and a generally well-rested condition.

Before beginning, the weaning process should be explained to the patient. Two methods used to wean patients are the T-piece and IMV. Regardless of the weaning method employed, the cuff of the endotracheal or tracheotomy tube is usually deflated first. In T-piece weaning, the patient is taken off the ventilator and a T-tube carrying humidified oxygen is attached to the airway. Patients should be encouraged to use the breathing exercises taught to them while they were on the ventilator. Weaning is started during the day and scheduled around medications and activities so the patient is not uncomfortable or interrupted. This also helps to keep the patient on a day/night schedule. The patient should be suctioned prior to disconnection from the ventilator. A semi-Fowler's position will facilitate the weaning process by preventing abdominal contents from pressing on the diaphragm [2; 6].

Weaning is usually started with 15 minutes off the ventilator and gradually increased if the patient's respiratory status remains stable. The patient's vital signs should be checked frequently during this period. Indications that weaning is being poorly tolerated include increased respiratory rate and dyspnea, tachycardia, cardiac dysrhythmias, change in mentation, fatigue, and decrease in pulmonary volumes. Arterial blood gas measurements are obtained for objective evidence of how well the patient is independently ventilating [6].

The other method of weaning is via the IMV method. The number of IMV breaths per minute is gradually decreased (without exceeding the patient's tolerance) until the patient is breathing totally spontaneously. The same nursing implications described for T-piece weaning apply [5; 6].

### **Psychologic Considerations and Counseling**

Patients who are mechanically ventilated for extended periods may develop a psychologic dependence on the ventilator that makes weaning difficult. Psychologic factors that impair the weaning process include fear of sudden death, anger from being asked to give up dependence on the ventilator, secondary depression when the illness is chronic, and interpersonal problems (e.g., a resentful or impatient spouse) [2; 6; 13].

Counseling may be necessary for patients and family members when attempts to wean from the ventilator are resisted. This is particularly true when physiologic parameters clearly indicate that weaning is feasible but difficulties are encountered. Along with counseling, psychotropic medications may be necessary. Antianxiety medications administered to patients who are near panic prior to weaning have been found helpful. If depression is a problem, antidepressants may be prescribed. Regardless of whether medications are used, weaning difficulty can be associated with emotional as well as physiologic factors [2; 8].



---

## CHEST TUBES

---

Inspiration and expiration depend partly on the presence of normal intrapleural pressure. Although this pressure varies with breathing, it remains lower than atmospheric pressure. Disruption of intrapleural pressure by trauma, disease, or surgery will decrease the effectiveness of ventilation. To re-establish normal intrapleural pressure, the insertion of chest tubes with a drainage system may be required. This will also prevent mediastinal shift [8].

A chest tube, also known as the thoracostomy tube, is a flexible tube inserted through the chest wall between the ribs into the pleural space. The chest tube is used to allow for re-expansion of the lung in situations such as pneumothorax, hemothorax, pleural effusions, chylothorax, or empyema. Mediastinal chest tubes are used during open-heart surgery to remove blood from around the heart and avoid pericardial tamponade; these types of chest tubes are not further discussed in this course [29; 30].

### CHEST TUBE INSERTION

The insertion of a chest tube can be painful. If the procedure is not an emergency, analgesics or pain medications should be given 30 minutes prior to tube insertion. It is of utmost importance that the nurse explains the procedure and equipment used. Patients and/or their family should be told that the tube(s) will be placed through the chest wall to drain air, fluid, or both, so the lungs can function normally. The tubes are secured by a dressing and connected to a drainage apparatus.

When chest tubes are inserted at the bedside, the nurse is responsible for obtaining equipment. The equipment may vary among institutions, but generally includes:

- Local anesthetic, usually 1% lidocaine
- Antiseptic, such as povidone-iodine
- Sterile gloves
- Collection receptacle

- Chest tube(s)
- Tape and gauze for an occlusive dressing

Unless contraindicated, the patient is usually placed in a recumbent position for insertion of the chest tube. If the tube is being inserted to remove air (e.g., in the case of pneumothorax), the patient should be supine. Because air rises, the chest tube is inserted anteriorly in the midclavicular line in the second to fifth intercostal space. If the tube is being placed to remove fluid (e.g., for hemothorax), the patient should be placed in a semi-Fowler's position. Because fluid collects in the dependent area, the tube is inserted in the sixth to eighth intercostal space in the midaxillary line. The patient who cannot tolerate a semi-Fowler's position should lie on the unaffected side [8]. For visual landmark placement, the fourth intercostal space is generally at nipple level on men or the level of the inframammary fold on women [8; 29]. A chest x-ray is typically completed post-insertion and at regular intervals for to monitor placement and resolution of the space-occupying lesion [31].

### DRAINAGE SYSTEMS

Chest tubes are connected to approximately 6 feet of tubing that leads to the collection system, which is placed considerably below the patient's chest in order to take advantage of gravity flow and facilitate removal of air and fluid from the intrapleural space. The tubing should be long enough that it allows the patient to move and turn without pulling. The dependent position and a long tube also prevent the backflow of fluid into the intrapleural space.

### CLOSED-CHEST DRAINAGE

Chest tube drainage systems are closed devices. Two types are utilized: wet suction and dry suction systems. However, not all patients who require a chest tube also require suction. A wet suction system controls suction by the level of water in the suction control chamber, normally -20 cm for adults. Less water in the chamber equals less suction.

A dry suction system is similar, but it uses a regulator to adjust the amount of suction instead of a water chamber. The dry suction system also responds to air leaks by delivering consistent and stable suction. To prevent the re-entry of air into the intrapleural space, the distal end of the chest tube should be submerged under water. This seal is a necessary part of any chest drainage unit. During exhalation, the pressure in the lungs forces air out of the pleural space through the tubing and into the water (as shown by bubbling) [8].

There are a number of variations of chest drainage units available. Previously, reusable glass systems were the standard. Now, sterile disposable systems are common. These systems are lightweight, take up a small amount of space, and are not breakable. Disposable systems can have one or multiple chambers, a valve or water seal, and/or water- or dry-suction control.

## NURSING CARE

The major objective of nursing care for patients with chest tubes is to facilitate drainage of fluid and air, fostering lung re-expansion. It is imperative to identify which system is being used and why. After insertion of the chest tube, a dressing is applied using sterile technique. Gauze or a hydrocolloid dressing is wrapped around the insertion site, and wide tape is applied to provide an occlusive dressing. This prevents air from entering the intrapleural space. Although the chest tube is sutured in place, further care should be taken to secure the tube when applying the tape. If drainage is present on the dressing, the physician should be notified [8].

Characteristics of the chest tube drainage should be recorded, including amount, color, consistency, and odor. Initially, the drainage characteristics should be noted every 15 to 20 minutes. Persistent drainage of more than 100 mL per hour should be reported. The collection should be labeled with the time, and drainage volumes should be recorded each nursing shift [8].

An important aspect of care for the patient with chest tubes is assessment of the patient's respiratory status. Patients should be questioned about their level of discomfort and any difficulty in breathing. Lung fields should be auscultated to determine if all areas of the lungs are being aerated. Watch for symmetrical expansion of the chest when the patient breathes, and listen for any adventitious breath sounds. The rate, depth, and quality of respirations should be noted and the patient encouraged to cough and deep breathe every hour to facilitate lung re-expansion and prevent complications [8].

## MAINTAINING THE SEAL

It is essential that the seal in the chest drainage unit be maintained to prevent re-entry of air into the pleural space. If a water seal is used, at least 2–2.5 cm of sterile water or normal saline should be placed in the water-seal bottle. The fluid level should be checked frequently and sterile water or saline added to replace water lost to evaporation. It is important that the drainage unit be kept well below the chest level; if any part of the system is lifted above the chest level, fluid may re-enter the pleural space (although most systems now have a one-way valve that prevents air or fluid from entering the chest). When transporting the patient, use a stretcher with a bottom shelf and place the collation unit on the stretcher shelf, keeping the system below the level of the patient's chest. It is recommended that someone familiar with the water-seal system accompany the patient during transport [8].



The Eastern Association for the Surgery of Trauma concludes that there is no evidence to support the routine use of presumptive antibiotics for post-traumatic tube thoracostomy to decrease the incidence of pneumonia or empyema.

(<https://tsaco.bmj.com/content/4/1/e000356>.  
Last accessed October 16, 2023.)

**Level of Evidence:** Expert Opinion/Consensus Statement

The water seal can be lost by inadvertently disconnecting or dislodging the tubing. Disconnection should be suspected if one notices an absence of fluctuation in the water-seal bottle or chamber. In the event that a tube becomes disconnected, the end of the disconnected tubing should be cleaned with an alcohol swab and reconnected. The nurse should then ask the patient to cough deeply to remove any excessive air that may have accumulated in the intrapleural space [8].

A chest tube should never be clamped, because a tension pneumothorax can develop if a continuous air leak is present [5; 6]. If a chest tube is pulled out or falls out of the intrapleural space, the patient should be asked to exhale forcefully, and the opening should be immediately covered. Ideally, the wound should be covered with gauze or dressing and pressure applied until the tube can be re-inserted. If gauze is not available, the palm of a gloved hand or a sheet can be used. If the patient develops signs and symptoms of a tension pneumothorax, the pressure should be momentarily relieved to allow air to escape [5; 6].

### **MAINTAINING PATENCY OF THE DRAINAGE SYSTEM**

Maintaining the patency of the drainage system is necessary to facilitate expansion of the lung, and frequent, systematic observations should be made to determine that the system is patent. This includes checking for loose connections and reinforcing the dressing as needed. All connections should be secured with tape or bands [8].

The water in the seal rises and falls with changes in intrapleural pressure. As the patient inhales, fluid should be pulled up into the water-seal tube or the chest tube, and the level should fall back as the patient exhales. Absence of fluctuation may be caused by an obstructed tube, a disconnected tube, or re-expansion of the lung [8].

Bubbling is normally present in the water-seal chamber during exhalation. However, continuous bubbling during both inspiration and expiration may signal an air leak. The air leak can be from the patient's lung or from an opening in the tubing or collection receptacle. A physician should be notified of any absence of fluctuation or continuous bubbling [5; 6; 8].

The drainage tubing should be regularly inspected for kinks or dependent loops. Kinking, which can occur along any section of the tubing from the chest tube to the collection device, will obstruct the flow of air or fluid. To prevent kinking at the point of entry into the drainage device, the tubing may be secured to a tongue blade with tape. Dependent (hanging) loops, which allow fluid to accumulate and thus alter the pressure within the system, can be avoided by proper positioning of the tubing. The tube should be coiled flat and should fall in a straight line from the coil to the drainage receptacle. The coiled tubing can be secured to the edge of the bed [5; 6].

When suction is applied, the suction control bottle or chamber should be observed for bubbling. Absence of bubbling suggests that sufficient suction is not being applied to the intrapleural space, whether because of an obstructed suction control tube, a leak in the system, an obstructed air inlet, or simply setting the suction too low. Incomplete expansion on one side only is due to increased intrapleural pressure.

Rarely, milking or stripping of the chest tubes may be necessary to prevent the formation of clots or remove clots from the tubing [6]. Evidence to support this practice is scant, and it should only be done if ordered by a physician. The tube should be secured at the incision site to prevent inadvertent dislodgement.

## DISCONTINUATION OF CHEST TUBES

When it is time to discontinue the chest tube, every effort should be made to remove it upon expiration. If the chest tube is removed on inspiration, a pressure gradient develops inside the chest that produces increased intrathoracic pressure, which predisposes the patient to a persistent/recurrent pneumothorax. Many methods can be used to prevent this from occurring. One is to time discontinuation of the tube in synchrony with the patient's breathing, removing it at expiration. Another is to have the patient hold his or her breath or make a seal and blow on their thumb like blowing up a balloon. An occlusive dressing is preferred post-removal. The provider may opt to place a U-stitch around the incision site and tighten it when the tube is discontinued to occlude the wound [29].

## PATIENT TEACHING REGARDING CHEST TUBES

As noted, the insertion of a chest tube is a painful, invasive procedure. Explanations and reassurance should be provided for the patient and family members. During the insertion, remain with the patient to provide emotional support. It is helpful to have one or two extra personnel on hand, if possible, to assist with the insertion and support the patient.

Patients with chest tubes may restrict their breathing and movement not only to minimize pain, but because they fear they may dislodge the tube. They should be informed that the tube is secured with sutures and tape. Movement such as turning, coughing, and deep breathing should be encouraged, as this will facilitate re-expansion of the lung. The patient should be told that pain medication is available to decrease discomfort during breathing exercises. Opioids, however, should be avoided or administered judiciously because they can depress the respiratory rate [8].

The collection receptacle should also be explained to the patient. If suction is necessary, patients should be aware that bubbling and some noise are expected. Patients and family members should be given the opportunity to verbalize fears and ask questions regarding the procedure and equipment.

Be aware that chest tubes may seem more frightening to visitors than other sorts of tubes, and be prepared to reassure patients that it is the visitor's lack of knowledge, not the patient's situation, that has caused the reaction [8].

---

## CONCLUSION

---

With knowledge of the purpose and function of artificial airways, mechanical ventilation, and chest tubes, nurses can readily provide quality and even lifesaving respiratory care. All healthcare professionals involved in the care of these patients should have a knowledge of the function, potential complications, and care skills necessary to ensure the health and safety of individuals who require respiratory assistance.

### Implicit Bias in Health Care

The role of implicit biases on healthcare outcomes has become a concern, as there is some evidence that implicit biases contribute to health disparities, professionals' attitudes toward and interactions with patients, quality of care, diagnoses, and treatment decisions. This may produce differences in help-seeking, diagnoses, and ultimately treatments and interventions. Implicit biases may also unwittingly produce professional behaviors, attitudes, and interactions that reduce patients' trust and comfort with their provider, leading to earlier termination of visits and/or reduced adherence and follow-up. Disadvantaged groups are marginalized in the healthcare system and vulnerable on multiple levels; health professionals' implicit biases can further exacerbate these existing disadvantages.

Interventions or strategies designed to reduce implicit bias may be categorized as change-based or control-based. Change-based interventions focus on reducing or changing cognitive associations underlying implicit biases. These interventions might include challenging stereotypes. Conversely, control-based interventions involve reducing the effects of the implicit bias on the individual's behaviors. These strategies include increasing awareness of biased thoughts and responses. The two types of interventions are not mutually exclusive and may be used synergistically.



## Works Cited

1. Berg SM, Bittner EA. *Massachusetts General Hospital Review of Critical Care Medicine*. Philadelphia, PA: Lippincott Williams & Wilkins; 2014.
2. Brashers VL. *Clinical Applications of Pathophysiology: An Evidence-Based Approach*. 3rd ed. St. Louis, MO: Mosby; 2006.
3. Bope ET, Kellerman RD. *Conn's Current Therapy 2017*. Philadelphia, PA: Elsevier; 2017.
4. Schwartzstein RM, Parker MJ. *Respiratory Physiology: A Clinical Approach*. Philadelphia, PA: Lippincott Williams & Wilkins; 2006.
5. American Thoracic Society, Infectious Diseases Society of America. Guidelines for the management of adults with hospital-acquired, ventilator-associated and healthcare-associated pneumonia. *Am J Respir Crit Care Med*. 2005;171(4):388-416.
6. Garrett L. *The Coming Plague: Newly Emerging Diseases in a World Out of Balance*. New York, NY: Farrar, Straus, Giroux; 1994.
7. Kalil AC, Metersky ML, Klompas M, et al, Infectious Diseases Society of America, American Thoracic Society. Management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clin Infect Dis*. 2016;63(5):e61-e111.
8. Nelms M, Sucher KP. *Nutrition Therapy and Pathophysiology*. 4th ed. Boston, MA: Cengage Learning; 2019.
9. Saunorus Baird M. *Manual of Critical Care Nursing: Nursing Interventions and Collaborative Management*. 7th ed. St. Louis, MO: Elsevier; 2016.
10. Alvarado AC, Panakos P. Endotracheal tube intubation techniques. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
11. Esianor BI, Campbell BR, Casey JD, et al. Endotracheal tube size in critically ill patients. *JAMA Otolaryngol Head Neck Surg*. 2022;148(9):849-853.
12. Ahmed RA, Boyer TJ. Endotracheal tube. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
13. Mora Carpio AL, Mora JI. Ventilator management. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
14. Avva U, Lata JM, Kiel J. Airway management. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
15. Kabrhel C, Thomsen TW, Setnik GS, Walls RM. Orotracheal intubation. *N Engl J Med*. 2007;356:e15.
16. Shindell M. Nasotracheal Intubation . Available at <https://emedicine.medscape.com/article/1663655>. Last accessed October 26, 2023.
17. Atanelov Z, Aina T, Amin B, et al. Nasopharyngeal airway. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
18. Siegmüller C. Endotracheal tubes. Anesthesia Airway Management (AAM). Available at <https://aam.ucsf.edu/endotracheal-tubes>. Last accessed October 25, 2023.
19. Meitner C, Feuerstein RA, Steele AM. Nursing strategies for the mechanically ventilated patient. *Front Vet Sci*. 2023;10:1145758.
20. Grensemann J, Gilmour S, Tariparast PA, Petzoldt M, Kluge S. Comparison of nasotracheal versus orotracheal intubation for sedation, assisted spontaneous breathing, mobilization, and outcome in critically ill patients: an exploratory retrospective analysis. *Scientific Reports*. 2023;13(1):12616.
21. Folino TB, Mckean G, Parks LJ. Nasotracheal intubation. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
22. Park DH, Lee CA, Jeong CY, Yang HS. Nasotracheal intubation for airway management during anesthesia. *Anesthesia and Pain Medicine*. 2021;16(3):232-247.
23. Raimonde AJ, Westhoven N, Winters R. Tracheostomy. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
24. Coelho L, Moniz P, Guerreiro G, Póvoa P. Airway and respiratory devices in the prevention of ventilator-associated pneumonia. *Medicina (Kaunas)*. 2023;59(2):199.
25. Kalil AC, Metersky ML, Klompas M, et al. Management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clin Infect Dis*. 2016;63(5):e61-e111.
26. Hickey SM, Giwa AO. Mechanical ventilation. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
27. PEPTIC Investigators for the Australian and New Zealand Intensive Care Society Clinical Trials Group, Alberta Health Services Critical Care Strategic Clinical Network, and the Irish Critical Care Trials Group, Young PJ, Bagshaw SM, et al. Effect of stress ulcer prophylaxis with proton pump inhibitors vs histamine-2 receptor blockers on in-hospital mortality among ICU patients receiving invasive mechanical ventilation: The PEPTIC Randomized Clinical Trial. *JAMA*. 2020;323(7):616-626.
28. Karlens MW, Holm A, Kvande ME, et al. Communication with mechanically ventilated patients in intensive care units: a concept analysis. *J Adv Nurs*. 2023;79(2):563-580.
29. Ravi C, McKnight CL. Chest tube. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.
30. Wallen M, Morrison A, Gillies D, O'Riordan E, Bridge C, Stoddart F. Mediastinal chest drain clearance for cardiac surgery. *Cochrane Database Syst Rev*. 2004;2002(4):CD003042.
31. Merkle A, Cindass R. Care of a chest tube. In: *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2023.

**Evidence-Based Practice Recommendations Citations**

Restrepo RD, Walsh BK. Humidification during invasive and noninvasive mechanical ventilation: 2012. *Respir Care*. 2012;57(5):782-788. Available at <http://rc.rcjournal.com/content/57/5/782>. Last accessed October 16, 2023.

Cook A, Hu CK, Ward J, et al. Presumptive antibiotics in tube thoracostomy for traumatic hemopneumothorax: a prospective, Multicenter American Association for the Surgery of Trauma Study. *Trauma Surg Acute Care Open*. 2019;4(1):e000356. Available at <https://tsaco.bmj.com/content/4/1/e000356>. Last accessed October 16, 2023.