

2023 CONTINUING EDUCATION FOR PENNSYLVANIA NURSES



INSIDE THIS EDITION

**Child Abuse Identification
& Reporting**

**Anxiety Disorders
in Older Adults**

Pancreatic Cancer

Meets 1 Hour of Pain Mgmt CE

**Pathophysiology: Muscles,
Joints, & Connective Tissues**

Meets 1 Hour of Pain Mgmt CE

We report your Child Abuse credit within 24 hours.

APRNs, we offer additional Pain Management/
Opioid CE (see Course Availability List, page 119).



In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

31 Hours
(10 Advanced Pharmacology Hrs)

\$52

[NetCE.com/PA23](https://www.netce.com/PA23)



31 Hours
Regular Price \$186

#97542 Child Abuse Identification & Reporting: The Pennsylvania Req. (3 contact hour)	1
#96690 Anxiety Disorders in Older Adults (3 contact hours)	19
#90240 Pancreatic Cancer (10 contact hours)	32
#38950 Pathophysiology: Muscles, Joints, and Connective Tissues (15 contact hours)	71
Customer Information/Evaluation	Located between pgs 88–89
Course Availability List	118–120

Special Offer price of \$52 valid through December 31, 2023

After December 31, 2023, price increases to \$62



LEARN

Read the enclosed course(s). Study questions will appear throughout the courses to create a link between the learning objectives and the supporting text.



CLICK

Go to **NetCE.com/PA23**. Enter your Quick Code and Customer ID (found on the back of your booklet), or complete the enclosed forms located between pages 88–89.



DONE

Receive certificates of completion:

FAST
By Mail

FASTER
Fax/Email

FASTEST!
Online

Print from your NetCE transcript.

Advanced Practice Nurses,

Need more Advanced Pharmacology hours?

Our Advanced Practice special offers are designed just for you.

You can also create your own **Special Offer** with **Design a Deal**.

Choose from our entire library of courses to create your own Special Offer. Your price is based on the number of hours you select.

Starting at only \$52!

(See inside back cover for details.)

Nurses,

Need more hours?

Get One Year of Unlimited Online CE Starting at Only \$85.

NetCE.com/NurseUnlimited

(See inside back cover for details.)

CONTINUING EDUCATION
FOR PENNSYLVANIA NURSES
2023

Published by NetCE,
a TRC Healthcare Company
P.O. Box 997571
Sacramento, CA 95899
Tel: 800-232-4238 (within the U.S.)
916-783-4238 (outside the U.S.)
Fax: 916-783-6067
Email: Info@NetCE.com
Website: www.NetCE.com

NETCE

Sr. Director of Development and Academic Affairs,
Sarah Campbell

Director of NetCE, Julie Goodwin

Chief Information Officer, Kevin Bluck

Director of Graphic Services, Kathryn Harris

Director of Operations, Alma Parra

Division Planners

Margaret Donohue, PhD

Alice Yick Flanagan, PhD, MSW

Margo A. Halm, RN, PhD, ACNS-BC

John V. Jurica, MD, MPH

John M. Leonard, MD

Ronald Runciman, MD

Shannon E. Smith, MHSC, CST, CSFA

Mark J. Szarejko, DDS, FAGD

Featured Contributing Faculty

Alice Yick Flanagan, PhD, MSW

Beyon Miloyan, PhD

Jane C. Norman, RN, MSN, CNE, PhD

Mark Rose, BS, MA, LP

Copyright © 2023 NetCE

Child Abuse Identification and Reporting: The Pennsylvania Requirement

This course is approved by the Pennsylvania Department of Human Services to fulfill the requirement for 2 hours of Child Abuse Recognition and Reporting (Act 31) training for healthcare professionals renewing their license. Provider number CACE000020.

Audience

This course is designed for all Pennsylvania nurses, physicians, physician assistants, social workers, counselors, pharmacists, and allied health professionals required to complete child abuse education.

Course Objective

The purpose of this course is to enable healthcare professionals in all practice settings to define child abuse and identify the children who are affected by violence. This course describes how a victim can be accurately diagnosed and identifies the community resources available in the state of Pennsylvania for child abuse victims.

Learning Objectives

Upon completion of this course, you should be able to:

1. Summarize the historical context of child abuse.
2. Discuss the emergence of the child welfare system in Pennsylvania.
3. Define child abuse and neglect and identify the different forms of child abuse and neglect.
4. Discuss the scope of child abuse and neglect in the United States and specifically in Pennsylvania.
5. Review the mandatory reporting process and mandated reporters in the state of Pennsylvania, including possible barriers to reporting suspected cases of child abuse.

Faculty

Alice Yick Flanagan, PhD, MSW, received her Master's in Social Work from Columbia University, School of Social Work. She has clinical experience in mental health in correctional settings, psychiatric hospitals, and community health centers. In 1997, she received her PhD from UCLA, School of Public Policy and Social Research. Dr. Yick Flanagan completed a year-long post-doctoral fellowship at Hunter College, School of Social Work in 1999. In that year she taught the course Research Methods and Violence Against Women to Masters degree students, as well as conducting qualitative research studies on death and dying in Chinese American families. (A complete biography can be found at NetCE.com/faculty.)

Faculty Disclosure

Contributing faculty, Alice Yick Flanagan, PhD, MSW, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Division Planner

Jane C. Norman, RN, MSN, CNE, PhD

Senior Director of Development and Academic Affairs

Sarah Campbell

Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Accreditations & Approvals



JOINTLY ACCREDITED PROVIDER™
INTERPROFESSIONAL CONTINUING EDUCATION

In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American

Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Designations of Credit

NetCE designates this continuing education activity for 3 ANCC contact hours.



IPCE CREDIT™

This activity was planned by and for the healthcare team, and learners will receive 3 Interprofessional Continuing Education (IPCE) credits for learning and change.

AACN Synergy CERP Category B.

Individual State Nursing Approvals

In addition to states that accept ANCC, NetCE is approved as a provider of continuing education in nursing by: Alabama, Provider #ABNP0353 (valid through 07/29/2025); Arkansas, Provider #50-2405; California, BRN Provider #CEP9784; California, LVN Provider #V10662; California, PT Provider #V10842; District of Columbia, Provider #50-2405; Florida, Provider #50-2405; Georgia, Provider #50-2405; Kentucky, Provider #7-0054 (valid through 12/31/2023); South Carolina, Provider #50-2405; West Virginia, RN and APRN Provider #50-2405.

Special Approvals

This course is approved by the Pennsylvania Department of Human Services to fulfill the requirement for 3 hours of Child Abuse Recognition and Reporting (Act 31) training for healthcare professionals applying for licensure. Provider number CACE000020.

This course is approved by the Pennsylvania Department of Human Services to fulfill the requirement for 2 hours of Child Abuse Recognition and Reporting (Act 31) training for healthcare professionals renewing their license. Provider number CACE000020.

About the Sponsor

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

Disclosure Statement

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

How to Receive Credit

- Read the following course.
- Complete the Self-Assessment Questions at the end of the course. (No test required.)
- Complete the Evaluation.
- Return your Customer Information/Evaluation and payment to NetCE by mail or fax, or complete online at www.NetCE.com/PA23.
- A full Works Cited list is available online at www.NetCE.com.



EVIDENCE-BASED
PRACTICE
RECOMMENDATION

Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the study questions and course material for better application to your daily practice.

HISTORICAL CONTEXT

Today, there is an established system in the United States to respond to reports of child abuse and neglect; however, this has not always been the case. This is not because child abuse, neglect, and maltreatment are new social phenomena. Rather, the terms “child abuse,” “child neglect,” and “child maltreatment” are relatively new, despite the fact that this social problem has existed for thousands of years [1]. Cruelty to children by adults has been documented throughout history and across cultures. In China, infant girls were often neglected during times of famine or sold during times of extreme poverty. There is also historical evidence that cultures have taken steps to stop child abuse and cruelty. For example, 6,000 years ago in Mesopotamia, orphans had their own patron goddesses for help and protection [2].

In many cases, the physical abuse of children has been linked to punishment. Throughout history, physical child abuse was justified because it was believed that severe physical punishment was necessary to discipline, rid the child of evil, or educate [2; 13]. It was not until 1861 that there was a public outcry in the United States against extreme corporal punishment. This reform was instigated by Samuel Halliday, who reported the occurrence of many child beatings by parents in New York City [2].

Sexual abuse of children, particularly incest (defined as sex between family members), is very much a taboo. The first concerted efforts to protect children from sexual abuse occurred in England during the 16th century. During this period, boys were protected from forced sodomy and girls younger than 10 years of age from forcible rape [2]. However, in the 1920s, sexual abuse of children was described solely as an assault committed by “strangers,” and the victim of such abuse was perceived as a “temptress” rather than an innocent child [2].

The first public case of child abuse in the United States that garnered widespread interest took place in 1866 in New York City. Mary Ellen Wilson was an illegitimate child, 10 years of age, who lived with her foster parents [3]. Neighbors were concerned that she was being mistreated; however, her foster parents refused to change their behaviors and said that they could treat the child as they wished [2]. Because there were no agencies established to protect children specifically, Henry Berge, founder of the Society for the Prevention of Cruelty to Animals, intervened on Mary’s behalf [3]. He argued that she was a member of the animal kingdom and deserved protection. The case received much publicity, and as a result, in 1874 the New York Society for the Prevention of Cruelty to Children was formed [3]. Because of this case, every state now has a system in place for reporting child abuse. The Pennsylvania Department of Public Welfare (now known as the Department for Human Services) was established in 1921 and part of its original intent was to care for “dependent, defective, and delinquent children” [7].

As a result of Berge’s advocacy for children’s safety, other non-governmental agencies were formed throughout the United States, and the establishment of the juvenile court was a direct result of the Society for the Prevention of Cruelty to Children [13]. By 1919, all but three states had juvenile courts. However, many of these nongovernmental agencies could not sustain themselves during the Depression [13].

The topic of child abuse and neglect received renewed interest in the 1960s, when a famous study titled “The Battered-Child Syndrome” was published [1; 4]. In the study, researchers argued that the battered-child syndrome consisted of traumatic injuries to the head and long bones, most commonly to children younger than 3 years of age, inflicted by parents [1; 4]. The study was viewed as the seminal work on child abuse, alerting both the general public and the academic community to the problems of child abuse [1; 2]. Soon, all 50 states required physicians to report child abuse [14]. In the early 1970s, Senator Walter Mondale noted that there was no official agency that spent its energies on preventing and treating child maltreatment [13]. Congress passed the Child Abuse Prevention and Treatment Act (CAPTA) of 1974, which targeted federal funds to improve states’ interventions for the identification and reporting of abuse [13]. In 2010, additional prevention and treatment programs were funded through CAPTA, and in 2012, the Administration on Children, Youth, and Families began to focus on protective factors to child abuse and neglect [61].

Today, child abuse and neglect are considered significant social problems with deleterious consequences. As noted, a system has been implemented in all 50 states to ensure the safety of children, with laws defining what constitutes abuse and neglect and who is mandated to report.

CHILD WELFARE IN PENNSYLVANIA

The Children’s Aid Society of Pennsylvania, one of the first organizations to advocate for children and their welfare in the United States, was founded in 1882 [62]. In the following years, the Children’s Aid Society was instrumental in educating the public about the unsanitary and unsafe conditions in almshouses, which were sometimes used for orphaned or abandoned children. Subsequently, legislation was passed in Pennsylvania to ensure that children were not permanently placed in almshouses [62].

In the state of Pennsylvania, Act 91 was passed in 1967 and gave child welfare agencies in all counties the responsibility to investigate child abuse reports made by physicians [18]. Three years later, Act 91 was modified to include school nurses and teachers as mandated reporters [18].

Pennsylvania was also the first state to take a noncriminal view of child abuse [22; 26]. In 1975, the Child Protective Services Law was enacted, which established a child abuse hotline and a statewide central registry in Pennsylvania in order to encourage the reporting of child abuse [18; 26].

The child welfare system in Pennsylvania is supervised by the state but administered by the different local counties [27]. This means that there are a total of 67 county agencies that administer the child welfare and juvenile justice services [27]. Aside from frank abuse, reports of other acts that might affect the well-being of a child are also accepted. The State of Pennsylvania delineates two functions for the local agencies: child protective services (CPS) and general protective services (GPS).

In 2017, Governor Tom Wolf approved Act 68 (also known as the Newborn Protection Act) to increase the number of locations for parents to give up their newborn without criminal liability [63]. In 2018, Act 29 was signed and expanded the definition of child abuse in Pennsylvania to include leaving a child unsupervised with a sexual predator [64].

CHILD PROTECTIVE SERVICES

CPS is in place to address acts that are “non-accidental serious physical or mental injury, sexual abuse, or exploitation, or serious physical neglect caused by acts or omissions of the parent or caretaker” [32]. In other words, these are cases in which there is reasonable cause to suspect child abuse and conduct an investigation.

Case Scenario

A young boy comes into the community health clinic for a physical exam. The boy’s mother hovers and does not seem to want to let her son answer any questions. During the exam, in the process of taking blood, the nurse notices some bruises and lacerations on the boy’s arm. Later, bruises in the shape of a belt are observed on the boy’s back as well. Upon questioning, the boy will only say that he was “bad.”

In this case, the nurse should make a report to ChildLine. This would be classified as a CPS case, and an investigation would be conducted. More information will be presented about reporting in later sections of this course.

GENERAL PROTECTIVE SERVICES

GPS is involved in non-abuse cases or acts that involve “non-serious injury or neglect” [38]. This includes children who experience “inadequate shelter, food, clothing, health care, truancy, inappropriate discipline, lack of supervision, hygiene issues, abandonment, or other problems that threaten a child’s opportunity for healthy growth and development” [38]. One of the following criteria must be met for GPS to be involved [55]:

- Lack of parental control
- Deprivation of the essentials of life
- Illegal placement for adoption or care
- Abandonment by parents or guardians

- Chronic truancy
- Habitual disobedience
- Formal adjudication
- Commitment of a delinquent act at an age younger than 10 years
- Defined as ungovernable
- Born to parents with terminated parental rights

Case Scenario

Ms. J, a neighbor, notices E (5 years of age) and S (6 years of age) running around their front yard at 8 p.m. The front door of the house is wide open, and Ms. J asks if their mother is home. S states that her mother went out with her girlfriend to a party. Ms. J asks if a babysitter is at the house, and S answers “no” again. This is not the first time neighbors have noticed that the kids are left at home alone. The neighbors report that the mother often comes home late, intoxicated.

In this case, a bystander (likely Ms. J or one of the neighbors) could call ChildLine, the local county agency, or even the police, and the case would be addressed by GPS. More information will be presented about reporting in later sections of this course.

DEFINITIONS OF CHILD ABUSE AND NEGLECT

The federal definition of child abuse is evident in CAPTA, published as a product of federal legislation. CAPTA defines a child to be any individual younger than 18 years of age, except in cases of sexual abuse. In cases of sexual abuse, the age specified by the child protection laws varies depending on the state in which the child resides [5]. CAPTA defines child abuse as, “any recent act or failure to act on the part of a parent or caretaker, which results in death, serious physical or emotional harm, sexual abuse, or exploitation, or an act or failure to act that presents an imminent risk of serious harm” [6].

In Pennsylvania, the child abuse law takes a very comprehensive approach to defining of child abuse [26]. According to Pennsylvania law, child abuse refers to intentionally, knowingly, or recklessly doing any of the following [43; 54]:

- Causing bodily injury to a child through any recent act or failure to act
- Fabricating, feigning, or intentionally exaggerating or inducing a medical symptom or disease that results in a potentially harmful medical evaluation or treatment to the child through any recent act
- Causing or substantially contributing to serious mental injury to a child through any act or failure to act or a series of such acts or failures to act
- Causing sexual abuse or exploitation of a child through any act or failure to act

- Creating a reasonable likelihood of bodily injury to a child through any recent act or failure to act
 - Creating a likelihood of sexual abuse or exploitation of a child through any recent act or failure to act
 - Causing serious physical neglect of a child
 - Engaging in any of the following recent acts:
 - Kicking, biting, throwing, burning, stabbing, or cutting a child in a manner that endangers the child
 - Unreasonably restraining or confining a child, based on consideration of the method, location, or duration of the restraint or confinement
 - Forcefully shaking a child younger than 1 year of age
 - Forcefully slapping or otherwise striking a child younger than 1 year of age
 - Interfering with the breathing of a child
 - Causing a child to be present at a location while a violation relating to the operation of methamphetamine laboratory is occurring, provided that the violation is being investigated by law enforcement
 - Leaving a child unsupervised with an individual, other than the child's parent, who the actor knows or reasonably should have known a) is required to register as a Tier II or Tier III sexual offender, where the victim of the sexual offense was younger than 18 years of age when the crime was committed; b) has been determined to be a sexually violent predator; or c) has been determined to be a sexually violent delinquent child
 - Causing the death of the child through any act or failure to act
 - Engaging a child in a severe form of trafficking in persons or sex trafficking, as those terms are defined under section 103 of the Trafficking Victims Protection Act of 2000
- In addition, the Code explicitly excludes specific acts and injuries from the definition of child abuse. Effective December 31, 2014, the following are considered exclusions to the definition of child abuse [44]:
- Environmental factors: No child shall be deemed to be physically or mentally abused based on injuries that result solely from environmental factors, such as inadequate housing, furnishings, income, clothing, and medical care, that are beyond the control of the parent or person responsible for the child's welfare with whom the child resides. This shall not apply to any child-care service, excluding an adoptive parent.
 - Practice of religious beliefs: If, upon investigation, the county agency determines that a child has not been provided needed medical or surgical care because of sincerely held religious beliefs of the child's parents or relative within the third degree of consanguinity and with whom the child resides, which beliefs are consistent with those of a bona fide religion, the child shall not be deemed to be physically or mentally abused. In such cases the following shall apply:
 - The county agency shall closely monitor the child and the child's family and shall seek court-ordered medical intervention when the lack of medical or surgical care threatens the child's life or long-term health.
 - All correspondence with a subject of the report and the records of the department and the county agency shall not reference child abuse and shall acknowledge the religious basis for the child's condition.
 - The family shall be referred for general protective services, if appropriate.
 - This subsection shall not apply if the failure to provide needed medical or surgical care causes the death of the child.
 - This subsection shall not apply to any child-care service as defined in this chapter, excluding an adoptive parent.
 - Use of force for supervision, control, and safety purposes: Subject to the rights of parents, the use of reasonable force on or against a child by the child's own parent or person responsible for the child's welfare shall not be considered child abuse if any of the following conditions apply:
 - The use of reasonable force constitutes incidental, minor, or reasonable physical contact with the child or other actions that are designed to maintain order and control.
 - The use of reasonable force is necessary to quell a disturbance or remove the child from the scene of a disturbance that threatens physical injury to persons or damage to property; to prevent the child from self-inflicted physical harm; for self-defense or the defense of another individual; or to obtain possession of weapons or other dangerous objects or controlled substances or paraphernalia that are on the child or within the control of the child.
 - Rights of parents: Nothing in this chapter shall be construed to restrict the generally recognized existing rights of parents to use reasonable force on or against their children for the purposes of supervision, control, and discipline of their children. Such reasonable force shall not constitute child abuse.

- Participation in events that involve physical contact with child: An individual participating in a practice or competition in an interscholastic sport, physical education, recreational activity, or extracurricular activity that involves physical contact with a child does not, in itself, constitute contact that is subject to the reporting requirements of this chapter.
- Defensive force: Reasonable force for self-defense or the defense of another individual shall not be considered child abuse.
 - Child-on-child contact: Harm or injury to a child that results from the act of another child shall not constitute child abuse unless the child who caused the harm or injury is a perpetrator. Notwithstanding this, the following shall apply: Acts constituting any of the following crimes against a child shall be subject to the reporting requirements: rape, involuntary deviate sexual intercourse, sexual assault, aggravated indecent assault, indecent assault, and indecent exposure.
 - No child shall be deemed to be a perpetrator of child abuse based solely on physical or mental injuries caused to another child in the course of a dispute, fight, or scuffle entered into by mutual consent.
 - A law enforcement official who receives a report of suspected child abuse is not required to make a report to the department if the person allegedly responsible for the child abuse is a nonperpetrator child.

It is important to note that exclusions are utilized by the CPS agency when investigating suspected abuse and should not be considered exclusions from reporting suspected abuse.

For the purposes of this course, a perpetrator is defined as a person who has committed child abuse. According to the Pennsylvania Code, the term includes only [42; 54]:

- A parent of the child
- A spouse or former spouse of the child's parent
- A paramour or former paramour of the child's parent
- A person 14 years of age or older and responsible for the child's welfare, including a person who provides temporary or permanent care, supervision, mental health diagnosis or treatment, or training or control of a child in lieu of parental care, supervision, and control
- An individual 14 years of age or older who resides in the same home as the child
- An individual 18 years of age or older who does not reside in the same home as the child but is related within the third degree of consanguinity or affinity by birth or adoption to the child

- An individual 18 years of age or older who engages a child in severe forms of trafficking in persons or sex trafficking, as those terms are defined under section 103 of the Trafficking Victims Protection Act of 2000

In a significant revision to the definition of perpetrator, school personnel and other childcare providers are considered "individuals responsible for the child's welfare" and may be perpetrators of child abuse; there is no longer a separate definition for student abuse [42]. As such, a perpetrator may be any such person who has direct or regular contact with a child through any program, activity, or services sponsored by a school, for-profit organization, or religious or other not-for-profit organization.

In addition, only the following may be considered a perpetrator for failing to act [42; 54]:

- A parent of the child
- A spouse or former spouse of the child's parent
- A paramour or former paramour of the child's parent
- A person 18 years of age or older and responsible for the child's welfare or who resides in the same home as the child

FORMS OF CHILD ABUSE AND NEGLECT

There are several acts that may be considered abusive, and knowledge of what constitutes abuse is vital for healthcare providers and other mandated reporters. In this section, specific behaviors that fall under the category of abuse and neglect will be reviewed.

Physical Abuse

Physical abuse injuries can range from minor bruises and lacerations to severe neurologic trauma and death. Physical abuse is one of the most easily identifiable forms of abuse and the type most commonly seen by healthcare professionals. Physical injuries that may be indicative of abuse include bruises/welts, burns, fractures, abdominal injuries, lacerations/abrasions, and central nervous system trauma [8; 34].

Bruises and welts are of particular concern, especially those that appear on:

- The face, lips, mouth, ears, eyes, neck, or head
- The trunk, back, buttocks, thighs, or extremities
- Multiple body surfaces

Patterns such as the shape of the article (e.g., a cord, belt buckle, teeth, hand) used to inflict the bruise or welt are common. Cigar or cigarette burns may be present, and they will often appear on the child's soles, palms, back, or buttocks. Patterned burns that resemble shapes of appliances, such as irons, burners, or grills, are of concern as well.

Fractures that result from abuse might be found on the child's skull, ribs, nose, or any facial structure. These may be multiple or spiral fractures at various stages of healing. When examining patients, note bruises on the abdominal wall, any intestinal perforation, ruptured liver or spleen, and blood vessel, kidney, bladder, or pancreatic injury, especially if accounts for the cause do not make sense. Look for signs of abrasions on the child's wrists, ankles, neck, or torso. Lacerations might also appear on the child's lips, ears, eyes, mouth, or genitalia. If violent shaking or trauma occurred, the child might experience a subdural hematoma [8; 34].



According to the American College of Radiology, fractures highly suggestive of physical abuse include rib fractures, classic metaphyseal lesions, those unsuspected or inconsistent with the history or age of the child, multiple fractures involving more than one skeletal area, and fractures of differing ages.

(<https://acsearch.acr.org/docs/69443/Narrative>. Last accessed July 26, 2022.)

Level of Evidence: Expert Opinion/Consensus Statement

Sexual Abuse/Exploitation

According to the Pennsylvania Code, sexual abuse or exploitation is defined as [45]:

- The employment, use, persuasion, inducement, enticement, or coercion of a child to engage in or assist another individual to engage in sexually explicit conduct, which includes, but is not limited to, the following:
 - Looking at the sexual or other intimate parts of a child or another individual for the purpose of arousing or gratifying sexual desire in any individual
 - Participating in sexually explicit conversation either in person, by telephone, by computer, or by a computer-aided device for the purpose of sexual stimulation or gratification of any individual
 - Actual or simulated sexual activity or nudity for the purpose of sexual stimulation or gratification of any individual
 - Actual or simulated sexual activity for the purpose of producing visual depiction, including photographing, videotaping, computer depicting, or filming

- Any of the following offenses committed against a child:
 - Rape
 - Statutory sexual assault
 - Involuntary deviate sexual intercourse
 - Sexual assault
 - Institutional sexual assault
 - Aggravated indecent assault
 - Indecent assault
 - Indecent exposure
 - Incest
 - Prostitution
 - Sexual abuse
 - Unlawful contact with a minor
 - Sexual exploitation

This does not include consensual activities between a child who is 14 years of age or older and another person who is 14 years of age or older and whose age is within four years of the child's age.

Child sexual abuse can be committed by a stranger or an individual known to the child. Sexual abuse may be manifested in many different ways, including [9; 10]:

- Verbal: Obscene phone calls or talking about sexual acts for the purpose of sexually arousing the adult perpetrator
- Voyeurism: Watching a child get dressed or encouraging the child to masturbate while the perpetrator watches
- Child prostitution: Involving the child in sexual acts for monetary profit
- Child pornography: Taking photos of a child in sexually explicit poses or acts
- Exhibitionism: Exposing his/her genitals to the child or forcing the child to observe the adult or other children in sexual acts
- Molestation: Touching, fondling, or kissing the child in a provocative manner; for example, fondling the child's genital area or long, lingering kisses
- Sexual penetration: The penetration of part of the perpetrator's body (e.g., finger, penis, tongue) into the child's body (e.g., mouth, vagina, anus)
- Rape: Usually involves sexual intercourse without the victim's consent and usually involves violence or the threat of violence
- Commercial sex act: Any sex act on account of which anything of value is given to or received by any person

Physical Neglect

Pennsylvania law defines serious physical neglect of a child as repeated, prolonged, or egregious failure to supervise a child in a manner that is appropriate considering the child's developmental age and abilities, and/or the failure to provide a child with adequate essentials of life, including food, shelter, or medical care, when committed by a perpetrator that endangers a child's life or health, threatens a child's well-being, causes bodily injury, or impairs a child's health, development, or functioning. Due to the ambiguity of definitions of child abuse and neglect, CAPTA provides minimum standards that each state must incorporate in its definition of neglect. Examples of child neglect may include [6; 11; 12]:

- Failure to provide adequate food, clothing, shelter, hygiene, supervision, and protection
- Refusal and/or delay in medical attention and care (e.g., failure to provide needed medical attention as recommended by a healthcare professional or failure to seek timely and appropriate medical care for a health problem)
- Abandonment, characterized by desertion of a child without arranging adequate care and supervision. Children who are not claimed within two days or who are left alone with no supervision and without any information about their parents'/caretakers' whereabouts are examples of abandonment.
- Expulsion or blatant refusals of custody on the part of parent/caretaker, such as ordering a child to leave the home without adequate arrangement of care by others
- Inadequate supervision (i.e., child is left unsupervised or inadequately supervised for extended periods of time)

Emotional Abuse

Under Pennsylvania law, emotional abuse involves an act or failure to act by a perpetrator that causes nonaccidental serious mental injury. Serious mental injury is "a psychological condition, as diagnosed by a physician or licensed psychologist, including the refusal of appropriate treatment, that renders a child chronically and severely anxious, agitated, depressed, socially withdrawn, psychotic, or in reasonable fear that his or her life or safety is threatened, or that seriously interferes with a child's ability to accomplish age-appropriate development and social tasks" [45].

The following behaviors could constitute emotional abuse [6; 11; 12]:

- Verbal abuse: Belittling or making pejorative statements in front of the child, which results in a loss or negative impact on the child's self-esteem or self-worth
- Inadequate nurturance/affection: Inattention to the child's needs for affection and emotional support

- Witnessing domestic violence: Chronic spousal abuse in homes where the child witnesses the violence
- Substance and/or alcohol abuse: The parent/caretaker is aware of the child's substance misuse problem but chooses not to intervene or allows the behavior to continue
- Refusal or delay of psychological care: Failure or delay in obtaining services for the child's emotional, mental, or behavioral impairments
- Permitted chronic truancy: The child averages at least five days per month of school absence and the parent/guardian does not intervene
- Failure to enroll: Failure to enroll or register a child of mandatory school age or causing the child to remain at home for nonlegitimate reasons
- Failure to access special education services: Refusal or failure to obtain recommended services or treatment for remedial or special education for a child's diagnosed learning disorder

Trafficking and Exploitation

It can be difficult to identify and intervene to stop human trafficking and exploitation, because it is hidden and even people who interact with victims may not recognize that it is happening. However, in many cases, women and children are considered the typical victims of human trafficking. Trafficking and exploitation are real risks to child safety and well-being and are reportable as forms of abuse.

There are several different types of child or minor human trafficking, but the term is generally defined as the recruitment, transportation, provision, or obtaining of a child for labor or services through the use of force, fraud, or coercion. Severe forms of human trafficking include sex and labor trafficking, including debt bondage and slavery.

Labor Trafficking

Labor trafficking is defined as labor obtained by the use of threat of serious harm, physical restraint, or abuse of the legal process. Severe labor trafficking includes the recruitment, harboring, transportation, provision, or obtaining of a person for labor or services, through the use of force, fraud, or coercion, for the purpose of subjection to involuntary servitude, peonage (i.e., paying off debt through work), debt bondage (i.e., debt slavery, bonded labor or services for a debt or other obligation), or slavery (i.e., a condition compared to that of a slave in respect of exhausting labor or restricted freedom).

Typically, children involved in forced labor are being given little or no pay. In the United States, forced labor is predominantly found in five sectors [57]:

- Prostitution and sex industry (46%)
- Domestic servitude (27%)

CHILD ABUSE VICTIMIZATION IN THE UNITED STATES ACCORDING TO RACE/ETHNICITY, 2020	
Race/Ethnicity	Child Abuse Rate per 1,000 Children
Native American/Alaska Native	15.5
African American	13.2
Multi-race	10.3
Pacific Islander	9.0
Hispanic	7.8
White	7.4
Asian American	1.6
<i>Source: [15]</i>	

Table 1

- Agriculture (10%)
- Sweatshops and factories (5%)
- Restaurant and hotel work (4%)

Among child victims, forced domestic servitude is a serious concern, particularly related to the provision of domestic services for 10 to 16 hours per day on activities such as child care, cooking, cleaning, and yard work/gardening.

Sex Trafficking

The Victims of Trafficking and Violence Protection Act defines sex trafficking as, “the recruitment, harboring, transportation, provision, or obtaining of a person for the purpose of a commercial sex act” [58]. A commercial sex act is, “any sex act on account of which anything of value is given to or received by any person” [58]. In other words, it involves the illegal transport of humans to be exploited in a sexual manner for financial gains [59]. Victims of sex trafficking could be forced into prostitution, stripping, pornography, escort services, and other sexual services [60]. Under federal law, sex trafficking (such as prostitution, pornography, or exotic dancing) does not require there be force, fraud, or coercion if the victim is younger than 18 years of age.

The term “domestic minor sex trafficking” has become a popular term used to connote the buying, selling, and/or trading of children for sexual services within the country, not internationally [60]. In the United States, the children most vulnerable to domestic minor trafficking are [60]:

- Youth in the foster care system
- Youth who identify as LGBTQIA+
- Youth who are homeless or runaway
- Youth with disabilities
- Youth with mental health or substance abuse disorders
- Youth with a history of sexual abuse

- Youth with a history of being involved in the welfare system
- Youth who identify as native or aboriginal
- Youth with family dysfunction

EPIDEMIOLOGY OF CHILD ABUSE AND NEGLECT

NATIONAL PREVALENCE

In 2020, there were 3.9 million referrals to child protective agencies in the United States [15]. More than 2.1 million (or 54%) were assessed to be appropriate for a response, and 27.6% of reports were made by health, social service, and/or mental health professionals [15]. Girls tend to be victims at a slightly higher rate (8.9 per 1,000 population) compared with boys (7.9 per 1,000 population) [15]. The most common perpetrators were parents; 90.6% of victims are maltreated by one or both parents [15]. Specifically, mothers are more often perpetrators compared with fathers (58.3% of victims were abused by a mother vs. 44.3% of victims were abused by a father) [15].

As of 2020, 8.4 of every 1,000 children in the United States were victims of abuse and/or neglect [15]. This is the unique rate, meaning each child is counted only once regardless the number of times a report may have been filed for abuse/neglect. The fatality rate for 2020 was 2.38 deaths per 100,000 children [15].

Research has shown that racial and ethnic minority children (particularly African American, Native American/Alaska Native, and multi-racial children) tend to have higher rates of reported child maltreatment compared with their White counterparts (*Table 1*) [15]. However, the lowest reported rate is among Asian American children [15].

PENNSYLVANIA STATE PREVALENCE

According to the Annual Child Protective Services Report, a yearly statistical report that documents child abuse cases in Pennsylvania, the child abuse hotline registered a total of 32,919 reports of suspected abuse or neglect in 2020 [27]. Approximately 14% of these cases were substantiated, which translates to 4,593 cases of child abuse in 2020 [27]. This is a decrease of 9,333 reports (22%) compared with 2019, a decline attributed to the COVID-19 pandemic, and the reduced contact between children and mandated reporters during that time [27]. Of the substantiated child abuse cases, there were 73 fatalities, 22 more than in 2019 [27]. More than half (51.4%) of perpetrators of child abuse in 2020 were the parent of the child victim [27].

RECOGNIZING WARNING SIGNS

It is crucial that practitioners become familiar with the indications of child abuse and neglect. These factors do not necessarily conclusively indicate the presence of abuse or neglect; rather, they are clues that require further interpretation and clinical investigation. Some parental risk indicators include [8; 10; 12; 16]:

- Recounting of events that do not conform either with the physical findings or the child's physical and/or developmental capabilities
- Inappropriate delay in bringing the child to a health facility
- Unwillingness to provide information or the information provided is vague
- History of family violence in the home
- Parental misuse of substances and/or alcohol
- Minimal knowledge or concern about the child's development and care
- Environmental stressors, such as poverty, single parenthood, unemployment, or chronic illness in the family
- Unwanted pregnancy
- Early adolescent parent
- Expression that the parent(s) wanted a baby in order to feel loved
- Unrealistic expectations of the child
- Use of excessive physical punishment
- Healthcare service "shopping"
- History of parent "losing control" or "hitting too hard"

Child risk indicators include [8; 10; 12; 16]:

- Multiple school absences
- Learning or developmental disabilities
- History of multiple, unexplained illnesses, hospitalizations, or accidents

- Poor general appearance (e.g., fearful, poor hygiene, malnourished appearance, inappropriate clothing for weather conditions)
- Stress-related symptoms, such as headaches or stomachaches
- Frozen watchfulness
- Mental illness or symptoms, such as psychosis, depression, anxiety, eating disorders, or panic attacks
- Regression to wetting and soiling
- Sexually explicit play
- Excessive or out-of-the-ordinary clinging behavior
- Difficulties with concentration
- Disruptions in sleep patterns and/or nightmares

In addition, warning signs specifically associated with victims of child trafficking and/or exploitation include (but are not limited to):

- A youth that has been verified to be younger than 18 years of age and is in any way involved in the commercial sex industry or has a record of prior arrest for prostitution or related charges
- An explicitly sexual online profile
- Excessive frequenting of Internet chat rooms or classified sites
- Depicting elements of sexual exploitation in drawing, poetry, or other modes of creative expression
- Frequent or multiple sexually transmitted infections or pregnancies
- Lying about or not being aware of their true age
- Having no knowledge of personal data (e.g., age, name, date of birth)
- Having no identification
- Wearing sexually provocative clothing
- Wearing new clothes of any style, getting hair and/or nails done with no financial means
- Being secretive about whereabouts
- Having late nights or unusual hours
- Having a tattoo that s/he is reluctant to explain
- Being in a controlling or dominating relationship
- Not having control of own finances
- Exhibiting hypervigilance or paranoid behaviors
- Expressing interest in or being in relationships with adults or much older men or women

Some of the types of behaviors and symptoms discussed in the definitions of physical, sexual, and emotional abuse/neglect are also warning signs. For example, any of the injuries that may result from physical abuse, such as a child presenting with bruises in the shape of electric cords or belt buckles, should be considered risk factors for abuse.

CONSEQUENCES OF CHILD ABUSE

The consequences of child abuse and neglect vary from child to child, and these differences continue as victims grow older. Several factors will mediate the outcomes, including the [17]:

- Severity, intensity, frequency, duration, and nature of the abuse and/or neglect
- Age or developmental stage of the child when the abuse occurred
- Relationship between the victim and the perpetrator
- Support from family members and friends
- Level of acknowledgment of the abuse by the perpetrator
- Quality of family functioning

In examining some of the effects of physical abuse, it is helpful to frame the consequences along a lifespan perspective [3]. During infancy, physical abuse can cause neurologic impairments. Most cases of infant head trauma are the result of child abuse [19]. Neurologic damage may also affect future cognitive, behavioral, and developmental outcomes. Some studies have noted that, in early childhood, physically abused children show less secure attachments to their caretakers compared to their non-abused counterparts [20].

By middle to late childhood, the consequences are more notable. Studies have shown significant intellectual and linguistic deficits in physically abused children [3]. Other environmental conditions, such as poverty, may also compound this effect. In addition, a number of affective and behavioral problems have been reported among child abuse victims, including anxiety, depression, low self-esteem, excessive aggressive behaviors, conduct disorders, delinquency, hyperactivity, and social detachment [3; 8; 10; 12].

Surprisingly, there has been little research on the effects of childhood physical abuse on adolescents [3]. However, differences have been noted in parents who abuse their children during adolescence rather than preadolescence. It appears that lower socioeconomic status plays a lesser role in adolescent abuse as compared with abuse during preadolescence [21]. In addition, parents who abuse their children during adolescence are less likely to have been abused as children themselves compared with those parents who abused their children during preadolescence [21]. It is believed that the psychosocial effects of physical abuse manifest similarly in late childhood and adolescence.

Research findings regarding the effects of childhood physical abuse on adult survivors indicate an increased risk for major psychiatric disorders, including depression, post-traumatic stress disorder, and substance abuse [36]. Some adult survivors function well socially and in terms of mental and physical health, even developing increased resilience as a result of their

experiences, while others exhibit depression, anxiety, post-traumatic stress, substance abuse, criminal behavior, violent behavior, and poor interpersonal relationships [3; 17; 46]. A meta-analysis found that adult survivors of child abuse were more likely to experience depression than non-abused counterparts, with the rates varying according to the type of abuse sustained (1.5-fold increase for physical child abuse, 2.11-fold increase for neglect, and 3-fold increase for emotional abuse) [24]. Similar results were found in a longitudinal study that compared a child welfare cohort to a group with no child welfare involvement. The child welfare group was twice as likely to experience moderate-to-severe depression and generalized anxiety compared with the control group [25]. There is some evidence that vulnerability to long-term effects of maltreatment in childhood may be at least partially genetically mediated [50].

Although not all adult survivors of sexual abuse experience long-term psychological consequences, it is estimated that 20% to 50% of all adult survivors have identifiable adverse mental health outcomes [23]. Possible psychological outcomes include [10]:

- Affective symptoms: Numbing, post-traumatic stress disorder, anxiety, depression, obsessions and compulsions, somatization
- Interpersonal problems: Difficulties trusting others, social isolation, feelings of inadequacy, sexual difficulties (e.g., difficulties experiencing arousal and orgasm), avoidance of sex
- Distorted self-perceptions: Poor self-esteem, self-loathing, self-criticism, guilt, shame
- Behavioral problems: Risk of suicide, substance abuse, self-mutilation, violence
- Increased risk-taking behaviors: Abuse of substances, cigarette smoking, sexual risk-taking

Adult male survivors of child sexual abuse are three times as likely to perpetrate domestic violence as non-victims. In addition, female survivors of child sexual abuse are more vulnerable to bulimia, being a victim of domestic violence, and alcohol use disorder [28].

In more recent years, research has focused on the impact of adverse childhood experiences (ACEs) in general. ACEs are defined as potentially traumatic experiences that affect an individual during childhood (before 18 years of age) and increase the risk for future health and mental health problems (including increased engagement in risky behaviors) as adults [47]. Abuse and neglect during childhood are clear ACEs, but other examples include witnessing family or community violence; experiencing a family member attempting or completing suicide; parental divorce; parental or guardian substance abuse; and parental incarceration [47]. Adults who experienced ACEs are at increased risk for chronic illness, impaired health, violence, arrest, and substance use disorder [28; 52].

REPORTING SUSPECTED CHILD ABUSE

Pennsylvania has a delineated process in place to facilitate the reporting of suspected child abuse. In addition, in 2014, Governor Corbett signed four new bills intended to streamline and clarify the child abuse reporting process in Pennsylvania. These bills were spurred by the Sandusky child sexual abuse case.

PERMISSIVE REPORTERS

There are two general categories of child abuse reporters: mandated reporters and permissive reporters. Permissive reporters are individuals who report an incident of suspected child abuse. These persons are not required to act or intervene in cases of suspected abuse. Put plainly, permissive reporters can report abuse while mandated reporters must report. However, it is important to note that any person is encouraged to report suspected child abuse or cause a report of suspected child abuse to be made to the department, county agency, or law enforcement, if that person has reasonable cause to suspect that a child is a victim of child abuse.

MANDATED REPORTERS

In Pennsylvania, a mandated reporter is required to make a report of suspected child abuse when he or she has reasonable cause to suspect that a child is a victim of child abuse if [48]:

- The mandated reporter comes into contact with the child in the course of employment, occupation, and practice of a profession or through a regularly scheduled program, activity, or service.
- The mandated reporter is directly responsible for the care, supervision, guidance, or training of the child, or is affiliated with an agency, institution, organization, school, regularly established church or religious organization, or other entity that is directly responsible for the care, supervision, guidance, or training of the child, regardless of the setting of the disclosure of abuse (within or outside of the reporter's professional role).
- A person makes a specific disclosure to the mandated reporter that an identifiable child is the victim of child abuse either within or outside of the reporter's professional role.
- An individual 14 years of age or older makes a specific disclosure to the mandated reporter (either within or outside of the reporter's professional role) that the individual has committed child abuse.

The mandated reporter is not required to interrogate the victim or identify the person responsible for the child abuse in order to make a report of suspected child abuse.

By law, individuals who come into contact with children on a frequent and consistent basis due to their work are legally required to report any suspected child abuse [39]. Mandated reporters in the state of Pennsylvania include, but are not limited to, [39]:

- Physicians (including osteopaths)
- Medical examiners
- Coroners
- Funeral directors
- Dentists
- Optometrists
- Chiropractors
- Podiatrists
- Interns
- Registered nurses
- Licensed practical nurses
- Hospital personnel engaged in the admission, examination, care, or treatment of persons
- Christian Science practitioners
- Members of the clergy
- School administrators
- School teachers
- School nurses
- Social services workers
- Day-care center workers or any other child-care or foster-care workers
- Mental health professionals
- Peace officers or law enforcement officials

Senate Bill 21 and House Bill 436 were two of the bills signed into law and enacted in 2014. These bills elucidate that mandated reporters are “to include anyone who comes in contact with a child, or is directly responsible for the care, supervision, guidance, or training of a child” [51]. Under this expanded definition, additional individuals who are also classified as mandatory reporters include [39]:

- A person licensed or certified to practice in any health-related field under the jurisdiction of the Department of State
- A school employee
- A foster parent
- An individual, paid or unpaid, who, on the basis of the individual's role as an integral part of a regularly scheduled program, activity, or service, accepts responsibility for a child
- An employee of a social services agency
- An employee of a public library

- Those who are supervised by mandated reporters
- An independent contractor with direct contact with children
- An attorney affiliated with an agency, institution, or organization that is responsible for the care, supervision, guidance, or control of children

It has long been debated whether attorneys should be included as mandated reporters. With this new definition, there is a seeming compromise, limiting the mandate to attorneys who are affiliated with an organization that is responsible for the care or supervision of children [37].

Privileged communication between any mandated reporter and his or her patient or client does not apply in cases of child abuse, and failure to report this information is considered a violation of the law [39]. There are exceptions: confidential communication made to an ordained member of the clergy (within the scope of 42 Pennsylvania CS §§ 5943), and confidential communications made to an attorney so long as they are within the scope of 42 Pennsylvania CS §§ 5916 (relating to confidential communications to attorney) and 5928 (relating to confidential communications to attorney), the attorney work product doctrine, or the rules of professional conduct for attorneys [39].

The Pennsylvania Code states that whenever a person is a mandated reporter in his or her capacity as a member of the staff of a medical or other public or private institution, school, facility, or agency, that person shall report immediately and immediately thereafter notify the person in charge of the institution, school, facility, or agency (or the designated agent) [48]. Upon notification, the person in charge or the designated agent is responsible for facilitating the cooperation of the institution, school, facility, or agency with the investigation of the report.

Not surprisingly, more than three-quarters (80%) of suspected child abuse reports are made by mandated reporters [27]. More specifically, the majority of child abuse reports come from mandated reporters in public/private social services agencies.

THE PROCESS OF REPORTING CHILD ABUSE IN PENNSYLVANIA

In Pennsylvania, mandated reports of potential child abuse (CPS or GPS cases) are made either in writing (through the online portal) or orally to ChildLine. The ChildLine is available seven days per week, 24 hours per day at 800-932-0313 or 412-473-2000. In 2020, ChildLine answered 163,215 calls, including suspected child abuse cases, referrals for GPS, and inquiries for general information to services [27]. Electronic submission of suspected child abuse reports may be made in lieu of calling ChildLine.

All mandated reporters who report via telephone shall also make a written report, which may be submitted electronically, within 48 hours [51]. The written reports are made through the Child Welfare Information Solution (CWIS) Portal, available online at <https://www.compass.state.pa.us/cwis>. The written report will include all of the following information, if known [55]:

- The names and addresses of the child, the child's parents, and any other person responsible for the child's welfare
- Where the suspected abuse occurred
- The age and sex of each subject of the report
- The nature and extent of the suspected child abuse, including any evidence of prior abuse to the child or any sibling of the child
- The name and relationship of each individual responsible for causing the suspected abuse and any evidence of prior abuse by each individual
- Family composition
- The source of the report
- The name, telephone number, and e-mail address of the person making the report
- The actions taken by the person making the report, including collection of evidence, protective custody, or admission to hospital
- Any other information required by federal law or regulation
- Any other information that the department requires by regulation

According to Pennsylvania law, a person or official required to report cases of suspected child abuse may take or request photographs of the child who is subject to a report and, if clinically indicated, request a radiologic examination and other medical tests on the child [56]. If completed, medical summaries or reports of the photographs, x-rays, and relevant medical tests should be sent along with the written report or within 48 hours after a report is made electronically.

Mandated reporters must identify themselves when reporting [54]. However, their names are usually not released; only the Secretary of the Department of Human Services has this authority. If a mandated reporter so chooses, he/she can sign a consent form that gives consent to have his/her name released [54].

A specialist at ChildLine will interview the caller to determine what the next step should be. This includes assessing if the report will be forwarded to a county agency for investigation as CPS or GPS; if a report should be forward directly to law enforcement officials; or if the caller will be referred to local services [53].

For both GPS and CPS cases, the appropriate county agency is contacted immediately [35]. The county agency is then responsible for its investigation, completing both a “risk assessment” and a “safety assessment.” In CPS cases, the agency sees and evaluates the child within 24 hours of receiving the report. The primary goal of the evaluations are to assess the nature and extent of the abuse reported; to evaluate the level of risk or harm if the child were to stay in the current living situation; and to determine action(s) needed to ensure the child’s safety [53].

A GPS referral will be assessed for any further needs, and appropriate referrals for services may be made for the child and family. If it is a CPS case, further investigation will be conducted. During the investigation, the agency may take photographs of the child and his/her injuries for the files. All investigations must be completed within 30 days from the date the report is taken at ChildLine [27]. Mandated reporters have a right to know of the findings of the investigation and the services provided to the child and may follow the case [33].

PROTECTIONS FOR REPORTERS

Reporters are afforded protections after reporting a suspected incidence of child abuse. Any person or institution who, in good faith, makes a report of child abuse, cooperates with a child abuse investigation, or testifies in a child abuse proceeding is considered immune from civil and criminal liability [44]. Mandated reporters who make a report in good faith and then later face discrimination in their workplace can take legal action [44]. For the most part, the reporter’s identity is kept confidential. If a case is referred to law enforcement, then the name of the reporter must be given upon request; however, reporters are treated as confidential informants [49].

PENALTIES FOR FAILURE TO REPORT

According to Pennsylvania statutes, a person or official required to report a case of suspected child abuse or to make a referral to the appropriate authorities who willfully fails to do so commits a misdemeanor of the third degree for the first violation and a misdemeanor of the second degree for a second or subsequent violation [44; 54]. An offense is a felony of the third degree if all three of the following are true:

- The person or official willfully fails to report.
- The child abuse constitutes a felony of the first degree or higher.
- The person or official has direct knowledge of the nature of the abuse.

A person who commits a second or subsequent offense commits a felony of the third degree, except if the child abuse constitutes a felony of the first degree or higher, in which case the penalty for the second or subsequent offenses is a felony of the second degree. In addition, if a person’s willful failure continues while the person knows or has reasonable cause to believe the child is actively being subjected to child abuse, the person commits a misdemeanor of the first degree; if the child abuse constitutes a felony of the first degree or higher, the person commits a felony of the third degree [44; 54].

BARRIERS TO REPORTING

Studies have shown that many professionals who are mandated to report child abuse and neglect are concerned and/or anxious about reporting. Identified barriers to reporting include [29; 30; 31; 40]:

- Professionals may not feel skilled in their knowledge base about child abuse and neglect. In addition, they lack the confidence to identify sexual and emotional abuse.
- Professionals may be frustrated with how little they can do about poverty, unemployment, drug use, and the intergenerational nature of abuse.
- Although professionals understand their legal obligation, they may still feel that they are violating patient confidentiality.
- Many professionals are skeptical about the effectiveness of reporting child abuse cases given the bureaucracy of the child welfare system.
- Practitioners may be concerned that they do not have adequate or sufficient evidence of child abuse.
- Practitioners may have a belief that government entities do not have the right to get involved in matters within the family.
- There may be some confusion and emotional distress in the reporting process.
- Practitioners may fear that reporting will negatively impact the therapeutic relationship.
- Some professionals have concerns that there might be negative repercussions against the child by the perpetrator.
- Some simply underestimate the seriousness and risk of the situation and may make excuses for the parents.

When interviewing children whose first language is not English, it is highly recommended that they be interviewed through the use of an interpreter. It can cause additional stress for children who struggle to find the right words in English, which can result in more feelings of fear, disempowerment, and voicelessness [41].

CASE SCENARIOS

In the following case scenarios, consider if the case should be reported as possible child abuse in accordance with Pennsylvania law.

A young girl, 2 years of age, is brought to the emergency department by her mother and stepfather for a scalp laceration. The girl is very quiet and appears listless and out of sorts. Her mother reports that she was injured when she fell onto a rock outside, but that the injury occurred when the girl was being watched by the stepfather. The girl undergoes assessment for traumatic brain injury, including assessment of function using the modified Glasgow Coma Score. The toddler is found to have mild impairment (a score of 13), and the follow-up test two hours later indicates normal functioning. The nurse notices that the toddler appears to be afraid of the stepfather, leaning away and crying when he is near her. The stepfather also appears to be easily frustrated with the child, saying that he does not know why she cries so much.

A boy, 13 years of age, is undergoing a routine physical exam with his family physician. The physician asks the boy if he is excited to start school in the next few weeks and how his baseball team is doing. The boy becomes quiet and states that he is nervous about an upcoming trip with his baseball team but does not give additional information. When asked directly, the boy says that he is uncomfortable with the new assistant coach, who watches pornography with them during out-of-town tournaments and supplies them with pornographic magazines. However, the boy states that he doesn't think it's a big deal and that "all of the other kids seem to really like it."

CONCLUSION

Child abuse and neglect are considered significant social problems with deleterious consequences. As noted, a system has been implemented in all 50 states to ensure the safety of children, with laws defining what constitutes abuse and neglect and who is mandated to report. Healthcare professionals, regardless of their discipline or field, are in a unique position to assist in the identification, education, and prevention of child abuse and neglect.

It is the duty of all mandated reporters in the state of Pennsylvania to know their responsibilities and the laws that govern the reporting process. All reporters should adhere to the established laws and rules that govern child abuse reporting, taking into account the expanded definition of perpetrator, the updated processes in place for reporting cases of suspected child abuse, and the delineated roles of mandated reporters. Doing so will help ensure the safety of millions of children in Pennsylvania.

RESOURCES

ChildLine: Pennsylvania Child Abuse Hotline
1-800-932-0313
<https://www.dhs.pa.gov/keepkidssafe>

Child Welfare Information Gateway
330 C Street SW
Washington, DC 20201
1-800-394-3366
To report abuse: 1-800-422-4453
<https://www.childwelfare.gov>

Child Welfare League of America
727 15th Street NW, 12th Floor
Washington, DC 20005
202-688-4200
<https://www.cwla.org>

National Council on Child Abuse and Family Violence
P.O. Box 5222
Arlington, VA 22205
202-429-6695
<https://www.preventfamilyviolence.org>

Pennsylvania Chapter of Children's Advocacy Centers and Multidisciplinary Teams
P.O. Box 3323
Erie, PA 16508
814-431-8151
<https://penncac.org>

Pennsylvania Child Welfare Information Solution
877-343-0494
<https://www.compass.state.pa.us/cwis>

Pennsylvania Department of Human Services
P.O. Box 2675
Harrisburg, PA 17105
1-800-692-7462
<https://www.dhs.pa.gov>

University of Pittsburgh, Pennsylvania Child Welfare Resource Center
403 East Winding Hill Road
Mechanicsburg, PA 17055
717-795-9048
<http://www.pacwrc.pitt.edu>

Customer Information/Evaluation insert located between pages 88–89.

SELF-ASSESSMENT QUESTIONS

#97542 CHILD ABUSE IDENTIFICATION AND REPORTING: THE PENNSYLVANIA REQUIREMENT

After reviewing the course, complete the following Self-Assessment Questions. Receive immediate feedback by reviewing the Study Guide provided on page 18. Please record the number of questions you correctly answered on the Evaluation. The Evaluation must be completed in order to receive credit for this course.

This 3 contact hour activity must be completed by July 31, 2025.

1. The first child abuse case in the United States that garnered widespread interest involved Mary Ellen Wilson, a foster child in New York City. This case took place in
 - A) 1790.
 - B) 1866.
 - C) 1921.
 - D) 1965.
2. How is the child welfare system in Pennsylvania characterized?
 - A) It is monitored by CAPTA.
 - B) The child welfare system is founded on the criminal justice model.
 - C) It is supervised by the state and administered by the various local county agencies.
 - D) It is supervised by each respective local county agency and administered by the federal government.
3. Child abuse is defined at the federal level by the
 - A) Child Protective Services.
 - B) Office of Child and Family Welfare.
 - C) Child Abuse Prevention and Treatment Act.
 - D) National Council on Child Abuse and Family Violence.
4. Which of the following injuries is NOT considered a possible indicator of physical abuse?
 - A) Patterned burns
 - B) Bruises on multiple body areas
 - C) Abrasions to the knees and elbows
 - D) Multiple or spiral fractures at various stages of healing
5. Child sexual abuse is categorized as exhibitionism if the act involves
 - A) obscene phone calls.
 - B) forcing a child to observe sexual acts.
 - C) watching a child get dressed or undressed.
 - D) touching, fondling, or kissing the child in a provocative manner.
6. How many substantiated cases of child abuse occurred in Pennsylvania in 2020?
 - A) 280
 - B) 1,967
 - C) 4,593
 - D) 26,944
7. Patient A, a child 10 years of age, arrives at the emergency department with a burn. Upon intake, a registered nurse notices that the burn on the child's thigh resembles the face of an iron. In addition, the child has bruising on her upper arm. The nurse suspects abuse and therefore calls the toll-free number for mandated reporters to report the case. Which of the following steps must the nurse take following the call?
 - A) This nurse must report the suspected abuse in writing within 48 hours.
 - B) The nurse must contact a physician for a complete evaluation of the child, including assessment for sexual abuse.
 - C) The nurse should call a legal aid society to ask for a lawyer to represent her in the event he/she is held liable if the case is not substantiated.
 - D) This nurse should inform the mother that he/she will be contacting the appropriate agencies regarding suspected child abuse and maltreatment.

8. When making a written report of suspected child abuse, the mandated reporter may be asked for
- A) *family composition.*
 - B) *photographs of the injuries, if available.*
 - C) *the location where the suspected abuse occurred.*
 - D) *All of the above*
9. The identity of the individual who reported a child abuse incident is NOT kept confidential if
- A) *the report is substantiated.*
 - B) *the incident is reported to law enforcement officials.*
 - C) *the intake specialist determines that the incident falls under general protective services.*
 - D) *the individual who reported the incident is determined to not have made the call in good faith.*
10. A failure to report suspected child abuse by a mandated reporter is considered a felony of the third degree if
- A) *the person willfully fails to report.*
 - B) *the child abuse constitutes a felony of the first degree or higher.*
 - C) *the person has direct knowledge of the nature of the abuse.*
 - D) *All of the above*

STUDY GUIDE

COURSE #97542 CHILD ABUSE IDENTIFICATION & REPORTING: THE PENNSYLVANIA REQUIREMENT

This Study Guide has been included to provide immediate feedback about your achievement for the course learning objectives. The correct answer for the Self-Assessment Questions and their locations within the text are indicated below.

- 1. The first child abuse case in the United States that garnered widespread interest involved Mary Ellen Wilson, a foster child in New York City. This case took place in**
B) 1866. (Historical Context)
- 2. How is the child welfare system in Pennsylvania characterized?**
C) It is supervised by the state and administered by the various local county agencies. (Child Welfare in Pennsylvania)
- 3. Child abuse is defined at the federal level by the**
C) Child Abuse Prevention and Treatment Act. (Definitions of Child Abuse and Neglect)
- 4. Which of the following injuries is NOT considered a possible indicator of physical abuse?**
C) Abrasions to the knees and elbows (Definitions of Child Abuse and Neglect; Forms of Child Abuse and Neglect; Physical Abuse)
- 5. Child sexual abuse is categorized as exhibitionism if the act involves**
B) forcing a child to observe sexual acts. (Definitions of Child Abuse and Neglect; Forms of Child Abuse and Neglect; Sexual Abuse/Exploitation)
- 6. How many substantiated cases of child abuse occurred in Pennsylvania in 2020?**
C) 4,593 (Epidemiology of Child Abuse and Neglect; Pennsylvania State Prevalence)
- 7. Patient A, a child 10 years of age, arrives at the emergency department with a burn. Upon intake, a registered nurse notices that the burn on the child's thigh resembles the face of an iron. In addition, the child has bruising on her upper arm. The nurse suspects abuse and therefore calls the toll-free number for mandated reporters to report the case. Which of the following steps must the nurse take following the call?**
A) This nurse must report the suspected abuse in writing within 48 hours. (Reporting Suspected Child Abuse; The Process of Reporting Child Abuse in Pennsylvania)
- 8. When making a written report of suspected child abuse, the mandated reporter may be asked for**
D) All of the above (Reporting Suspected Child Abuse; The Process of Reporting Child Abuse in Pennsylvania)
- 9. The identity of the individual who reported a child abuse incident is NOT kept confidential if**
B) the incident is reported to law enforcement officials. (Reporting Suspected Child Abuse; Protections for Reporters)
- 10. A failure to report suspected child abuse by a mandated reporter is considered a felony of the third degree if**
D) All of the above (Reporting Suspected Child Abuse; Penalties for Failure to Report)

Anxiety Disorders in Older Adults

Includes 1 Advanced Pharmacology Hour

Audience

This course is designed for the benefit of a broad range of allied health professionals, including but not limited to, nurses, physicians, medical assistants, and nursing home administrators.

Course Objective

Older adults are the fastest growing demographic in the world, and anxiety disorders are the most common mental disorder in this age group. The purpose of this course is to provide clinicians with the knowledge and skills necessary in order to improve the assessment and treatment of anxiety disorders in older adults.

Learning Objectives

Upon completion of this course, you should be able to:

1. Describe the history and neuroanatomy of anxiety and anxiety disorder.
2. Discuss the assessment and classification of anxiety disorders in older adults.
3. Analyze the epidemiology of anxiety disorders in elderly patients.
4. Describe the clinical implications of late-life anxiety disorders and their treatment.

Faculty

Beyon Miloyan, PhD, received his PhD in Psychology from the University of Queensland in 2015 for his thesis on late-life anxiety disorders. He completed his postdoctoral training in the Epidemiology and Biostatistics of Aging program at the Johns Hopkins University before taking a tenure-track position in the School of Psychology and Health Sciences at Federation University, Australia. Dr. Miloyan has published 30 peer-reviewed journal articles and book chapters and has been teaching since 2012. He has supervised 10 student theses at doctoral, Master's, and undergraduate levels and served as an ad hoc peer reviewer for various journals in the fields of psychology, psychiatry, and public health.

Faculty Disclosure

Contributing faculty, Beyon Miloyan, PhD, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Division Planner

Jane C. Norman, RN, MSN, CNE, PhD

Senior Director of Development and Academic Affairs
Sarah Campbell

Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Accreditations & Approvals



JOINTLY ACCREDITED PROVIDER™
INTERPROFESSIONAL CONTINUING EDUCATION

In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCM), the Accreditation Council for Pharmacy Education (ACPE), and the American

Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Designations of Credit

NetCE designates this continuing education activity for 3 ANCC contact hours.



IPCE CREDIT™

This activity was planned by and for the healthcare team, and learners will receive 3 Interprofessional Continuing Education (IPCE) credits for learning and change.

NetCE designates this continuing education activity for 1 pharmacotherapeutic/pharmacology contact hour.

AACN Synergy CERP Category A.

Individual State Nursing Approvals

In addition to states that accept ANCC, NetCE is approved as a provider of continuing education in nursing by: Alabama, Provider #ABNP0353 (valid through 07/29/2025); Arkansas, Provider #50-2405; California, BRN Provider #CEP9784; California, LVN Provider #V10662; California, PT Provider #V10842; District of Columbia, Provider #50-2405; Florida, Provider #50-2405; Georgia, Provider #50-2405; Kentucky, Provider #7-0054 (valid through 12/31/2023); South Carolina, Provider #50-2405; West Virginia, RN and APRN Provider #50-2405.

About the Sponsor

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

Disclosure Statement

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

How to Receive Credit

- Read the following course.
- Complete the Self-Assessment Questions at the end of the course. (No test required.)
- Complete the Evaluation.
- Return your Customer Information/Evaluation and payment to NetCE by mail or fax, or complete online at www.NetCE.com/PA23.
- A full Works Cited list is available online at www.NetCE.com.



Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the study questions and course material for better application to your daily practice.

INTRODUCTION

The first known clinical case description of an anxiety disorder appeared in the medical corpus of the Ancient Greek physician Hippocrates. The description tells of Nicanor, a man who developed an extreme fear of a “flute girl” whom he encountered one night at a drinking party and who haunted him every night for many years. Five hundred years after this case description, the Ancient Roman Stoic philosophers Seneca the Younger and Cicero addressed the topic of anxiety at length, recognizing both its benefits and harms, depending on the severity and circumstances of the anxiety [1]. These texts reveal a sophisticated understanding of fear and anxiety among

these ancient authors, even by modern medical standards. It was not until the 19th century that Charles Darwin noted essential similarities in the expression of fear and anxiety in mammals, reinforcing Seneca’s notion that fear and anxiety are ultimately adaptive traits [2]. In its normal state, anxiety facilitates the management of potential future hazards [3; 4; 5]. In its extreme state, the individual regards it as excessive or distressing or it can cause impairment in the individual’s daily life, thus constituting a disorder [6; 7].

The analogy of a smoke detector demonstrates the adaptive and maladaptive aspects of anxiety [8; 9]. Just as the function of a smoke detector is to signal potential fires so that one can take action to prevent harm, the function of anxiety is to signal any potential hazards so that preventive actions can be taken. In this analogy, an anxiety disorder is an extreme that renders the individual more sensitive to threat signals [10]. Although those with higher anxiety experience more false alarms (signals for a threat that does not occur), this is advantageous to the extent that it reduces the risk of a fatal miss. In other words, the costs associated with false alarms and misses are not equal: over-reacting to non-threats is generally less costly than failing to detect one danger. Nonetheless, living in a chronic state of high anxiety can take a long-term toll on an individual’s health and quality of life, and in these cases, intervention is warranted.

Age-related changes in anxiety can occur over the course of one’s life, and understanding these changes is key to facilitating clinical detection and treatment, particularly among older adults, who are the largest and fastest-growing age demographic in the United States. This course begins by addressing the neuroanatomy of anxiety, followed by its classification and a review of commonly used methods of assessment. The course goes on to cover the epidemiology of anxiety disorders in older adults, including its prevalence, incidence, course, risk factors, and consequences. Finally, treatment considerations are addressed.

NEUROANATOMY

In 1949, the Nobel Prize in Medicine was awarded to António Egas Moniz for his discovery of “a simple operation, always safe, [and] which may prove to be an effective surgical treatment in certain cases of mental disorder” [11]. Specifically, Moniz discovered the prefrontal leukotomy as a treatment for mental disorders, including anxiety disorders [12]. Since then, studies have found that damage to the ventromedial prefrontal cortex produces resistance against anxiety and depression [13; 14; 15; 16]. Despite the effective reduction of anxiety in these patients, it took many decades until research began to address the harms imposed by damage to the prefrontal cortex. For example, in addition to reducing anxiety, damage to the ventromedial prefrontal cortex also impairs self-regulation and decision-making and can induce sociopathic behaviors [17; 18; 19; 20; 21]. Similar patterns of anxiety reduction were also observed

in one patient with focal bilateral lesions to the amygdalae who showed a similar pattern of impairment in her daily life as those with damage to the prefrontal cortex [22]. Although the prefrontal cortex and amygdala are critical structures in a neural network that is necessary for anxiety, these findings highlight the fact that damage to these structures comes with unintended consequences. These findings also highlight the more general point that, in treating anxiety disorders, it is also important to not abolish otherwise useful traits as it is to reduce the anxiety to a manageable level.

CLASSIFICATION

The *Diagnostic and Statistical Manual of Mental Disorders* (DSM) sub-classifies anxiety disorders into panic disorder, agoraphobia, specific phobia, social anxiety disorder, and generalized anxiety disorder (GAD) [6]. In the following sections, each of these subtypes are described, including the relevant criteria for diagnosis and a description of age-related differences in symptom patterns. The most frequently reported symptom(s) for each disorder in older adults are based on data from the National Epidemiological Survey on Alcohol and Related Conditions (NESARC).

PANIC DISORDER AND AGORAPHOBIA

Panic disorder is characterized by the occurrence of panic attacks. Panic attacks are defined as sudden, unexpected, and brief onsets of terror, accompanied by at least four of the following symptoms: sweating, trembling, chest pain, dizziness, nausea, chills or hot flashes, numbness or tingling, shortness of breath or choking, a feeling of loss of control, desensitization, or a fear of death. In order to be classified as panic disorder, the DSM requires such panic attacks to be accompanied by a period of at least one month in which the individual also fears the possibility of a future panic attack [6].

Two main subtypes of panic disorder have been observed, diverging between individuals with respiratory and non-respiratory symptoms [23; 24; 25]. Determining the subtype may be informative for treatment purposes. Older adults with panic disorder experience fewer symptoms of panic compared with younger adults, and their panic attacks are also reported to be less intense and shorter in duration [26; 27; 28; 29].

Agoraphobia is characterized by a fear or avoidance of situations from which escape is difficult. The diagnosis requires a fear or avoidance of two or more of the following specific situations: public transportation, open spaces, closed spaces, crowds, or being alone in public. Although the presence of these fears is also associated with panic disorder and specific phobia, a distinguishing factor of agoraphobia is defined by the frequency of the aforementioned fears. Individuals with greater and more frequent occurrences of these fears tend to be classified as having agoraphobia [30].

Table 1 displays the most commonly reported panic symptoms among older adults (55 years of age and older) with a diagnosis of panic disorder (with or without agoraphobia). The total percentage of each symptom is displayed. The prevalence of panic disorder in this sample was 1.2% (95% confidence interval [CI]: 1.0-1.5).

SPECIFIC PHOBIA

The central feature of specific phobia is the fear or avoidance of specific objects or situations. These include, but are not limited to, animals (e.g., snakes or insects), the natural environment (e.g., storms, water, or heights), situations (e.g., typically closed or open spaces), and blood, injections, or injury. A diagnosis of specific phobia requires the individual to recognize that the fear or avoidance is unreasonable and to regard it as distressing or interfering with their everyday life. The most common fears reported by adults involve animals, heights, and flying [32; 33; 34]. However, older adults frequently report situational fears [35]. Individuals who report having at least one specific fear are likely to report having other fears [33; 36].

Table 2 displays the most commonly reported specific fears among older adults (55 years of age and older) with a diagnosis of specific phobia. The prevalence of specific phobia in this sample was 5.5% (95% CI: 5.0-6.0).

SOCIAL ANXIETY DISORDER

The core feature of social anxiety disorder is the fear or avoidance of social situations. The fear or avoidance concerns the possibility of negative judgment by others (e.g., resulting in embarrassment or humiliation). Social anxiety may pertain to particular types of social settings or situations, such as small or large group settings, or the anxiety may generalize to a variety of social situations. Older adults endorse fewer social anxiety symptoms relative to younger adults [37]. The most common social fears among older adults include public speaking or being confronted or criticized by others, while discomfort with and avoidance of social situations and experiencing anxiety when thinking about social situations appears equally common to both younger and older adults [37; 38].

Table 3 displays the most commonly reported social fears among older adults (55 years of age and older) with a diagnosis of social anxiety disorder. The prevalence of social anxiety disorder in this sample was 2.0% (95% CI: 1.7-2.3).

GENERALIZED ANXIETY DISORDER

The central feature of GAD is intrusive worry, defined as repetitive thinking about potentially harmful future events. Worries generally pertain to everyday concerns and involve attempts to minimize the likelihood or consequences of disadvantageous outcomes. Although some degree of worry is recognized as helpful, when the individual reports experiences of excessive and uncontrollable worry for a period of six months or more, this constitutes a diagnosis of GAD if, and only if, the worry is also regarded by the individual as causing distress or impairment [39]. Age-related reductions in worry frequency have

PREVALENCE OF PANIC SYMPTOMS AMONG OLDER ADULTS WITH PANIC DISORDER			
Symptom	Age Groups		
	55 to 64 Years	65 Years and Older	55 Years and Older
Shortness of breath	82%	85%	83%
Heart racing/pounding	91%	82%	88%
Trembling/shaking	73%	69%	72%
Perspiring/sweating	75%	65%	72%
Felt as if choking	45%	57%	50%
Dizzy/lightheaded	74%	74%	74%
Things seemed unreal	61%	56%	59%
Tingling/numbness	57%	50%	54%
Flushes/hot flashes/chills	71%	53%	64%
Nauseous/upset stomach	50%	56%	53%
Pain/pressure in chest	63%	56%	60%
Going crazy/losing control	63%	61%	62%
Felt might die	58%	64%	60%

Source: [31] Table 1

PREVALENCE OF SPECIFIC FEARS AMONG OLDER ADULTS WITH SPECIFIC PHOBIA				
Object of Fear/ Avoidance	Age Groups			
	55 to 64 Years	65 to 74 Years	75 Years and Older	55 Years and Older
Animals	57%	56%	62%	58%
Heights	53%	60%	49%	55%
Storms	26%	30%	45%	31%
Being in/on water	31%	41%	45%	37%
Flying	35%	36%	33%	35%
Crowds/lines	13%	12%	23%	15%
Closed spaces	40%	38%	41%	40%
Blood/injections	16%	11%	17%	15%
Public transportation	9%	6%	7%	7%
Going to the dentist	31%	32%	27%	30%
Hospitals	15%	12%	17%	15%

Source: [31] Table 2

been observed in older adult samples from the United States, United Kingdom, Canada, and Australia [40; 41; 42; 43; 44; 45; 46]. There are also age-related differences in the subjects of individuals' worries. For example, for younger adults, common worries concern work, finances, and personal relationships. For older adults, these concerns give way to worries about personal health and the health and welfare of loved ones [41; 44; 46]. These "world issue" worries typically focus outwardly on problems that could be faced by future generations, which may be of particular relevance during this developmental life stage [43]. In fact, late-life developmental transitions have

been associated with other context-specific worries, such as concerns of becoming a burden after transitioning out of a primary caregiver role and into retirement [47; 48]. Caregiving, too, can be a significant source of worry, anxiety, and distress in later life [49; 50; 51]. Older adults who report financial worries tend to be concerned about receiving care and about their own capacity to make decisions [52]. However, despite the observation that older adults with GAD tend to endorse a greater variety of worries than matched non-GAD controls, there are fewer differences in the experience of worry between older adults with and without GAD than there are between

**PREVALENCE OF SOCIAL ANXIETY SYMPTOMS AMONG
OLDER ADULTS WITH SOCIAL ANXIETY DISORDER**

Symptom	Age Groups		
	55 to 64 Years	65 Years and Older	55 Years and Older
Social situations made you nervous	81%	85%	83%
Social situations made you upset/anxious	93%	93%	93%
Endured social situations that frightened you	88%	85%	87%
Avoided social situations out of strong fear	82%	72%	77%
More frightened in social situations than most people	80%	76%	78%
Thought fear of social situations stronger than it should be	92%	89%	90%
Had a panic attack in social situations	22%	8%	15%
Frightened of social situations out of fear of panic attack	16%	8%	12%
Remained in social situation despite fear of panic attack	20%	11%	15%

Source: [31] Table 3

**PREVALENCE OF GENERALIZED ANXIETY DISORDER SYMPTOMS
AMONG OLDER ADULTS WITH GENERALIZED ANXIETY DISORDER**

Symptom	Age Groups		
	55 to 64 Years	65 Years and Older	55 Years and Older
Worry a lot about things you usually didn't worry about?	83%	72%	78%
Ever think your worrying was excessive?	56%	47%	52%
Often got tired easily	83%	81%	82%
Often had tense, sore, aching muscles	76%	67%	71%
Became so restless you paced, fidgeted, or could not sit still	62%	55%	58%
Often felt keyed up or on edge	82%	78%	80%
Often had trouble concentrating	83%	83%	83%
Often felt irritable	80%	62%	71%
Often had trouble falling/staying asleep	77%	72%	74%
Often forgot what you were talking about/mind went blank	75%	66%	70%
Often felt heart racing, skipping, or pounding in chest	59%	45%	52%
Often perspired/sweated	50%	35%	43%
Often had cold and clammy hands	44%	27%	36%
Often had dry mouth	58%	49%	54%
Often felt dizzy/lightheaded/like might faint	48%	53%	50%
Often felt nauseous	54%	34%	45%
Often urinated frequently	54%	47%	51%
Often had trouble swallowing/felt like lump in throat	37%	34%	35%
Often had pain/pressure in chest	40%	31%	36%
Often trembled/shook	34%	39%	36%
Often had trouble catching breath/felt like smothering	43%	33%	39%

Source: [31] Table 4

younger adults with and without GAD [53; 54]. In essence, the expression of worry may vary significantly as a function of the developmental stage of the individual, with older adults endorsing worries commensurate with their changing life circumstances [55].

Table 4 displays the most commonly reported generalized anxiety disorder symptoms among older adults (55 years of age and older) with a diagnosis of GAD. The prevalence of GAD in this sample was 2.0% (95% CI: 1.7–2.3).

ASSESSMENT

STRUCTURED AND SEMI-STRUCTURED INTERVIEWS

The standard procedure for anxiety disorder assessment is the structured diagnostic interview, which is administered by a trained professional. The structured interview consists of pre-determined questions that assess for relevant symptoms based on diagnostic criteria. For example, an interview for GAD would start by asking the individual questions about the presence of worry symptoms over the past six months. If the interviewee answers this question affirmatively, the interviewer would then ask the individual about the presence of secondary symptoms associated with the worry (e.g., sleep, irritability). If the individual responds affirmatively to the minimum number of secondary symptoms required for a diagnosis of GAD, the individual would then be queried about the presence of distress or impairment due to the worry. The key advantage of the structured interview is its standardized administration, procedure, and scoring, which minimize bias and error in assessment. Two commonly used structured interviews for the assessment of mental disorders are the Diagnostic Interview Schedule (DIS) and the Composite International Diagnostic Interview (CIDI) [56; 57]. In addition, the Anxiety Disorders Interview Schedule (ADIS) is a structured diagnostic interview that was developed specifically for anxiety disorder assessment [58]. These interviews are regularly updated along with diagnostic criteria, as for example with new editions of the DSM. Structured interviews rely essentially on self-report; in addition to being administered by clinicians, they may also be conducted by trained lay persons and/or computer-assisted technology (as in epidemiologic surveys).

The examination modality of assessment contrasts with the interview technique in that the person conducting the assessment, typically a trained clinician, decides about the presence or absence of a symptom instead of relying on the report of the individual. The Structured Clinical Interview for the DSM (SCID) and the Schedules for Clinical Assessment in Neuropsychiatry (SCAN) are examples of semi-structured interviews/examinations that allow the clinician to take

a more flexible approach to the interview while retaining some degree of structure [59; 60]. Structured interviews and semi-structured examinations are not always practical to use because they are time-consuming to administer. Nonetheless, they are essential for validating briefer, easier to administer, and more widely used questionnaires and screening tools for use in particular contexts.

As a result of the evolving racial and immigration demographics in the United States, interaction with patients for whom English is not a native language is inevitable. Because diagnosing anxiety disorders is reliant on good communication, it is each practitioner's responsibility to ensure that interviews and assessments are conducted in such a way that allows for patient understanding. When there is an obvious disconnect in the communication process between the practitioner and patient due to the patient's lack of proficiency in the English language, an interpreter is required.

Mental health professionals should consider undertaking a language needs analysis for the service population and consider how to best meet identified needs. If possible, 10 to 15 minutes should be reserved in advance of sessions to brief the interpreter about the purpose of the meeting and to enable them to explain any cultural issues that may have bearing on the session.

RATING SCALES

Generalized Anxiety Disorder

The Generalized Anxiety Disorder 7-item (GAD-7) scale is a brief, self-administered screening instrument for use in medical settings. The scale assesses for symptoms occurring over the previous two weeks of the respondent's life [61; 62]. Each item is rated on a four-point scale (0 to 3) yielding a maximum score of 21. A score of 10 or greater indicates a probable diagnosis of GAD based on validation against the psychiatrist-administered SCID [62]. There is also a shorter, two-item version called the GAD-2. The GAD-7 (along with scoring instructions) can be accessed online at https://adaa.org/sites/default/files/GAD-7_Anxiety-updated_0.pdf. The GAD-2 consists of only the first two items of the GAD-7, with scores of 3 or greater indicating clinically significant anxiety symptoms [63].

Panic Disorder

The Panic Disorder Severity Scale (PDSS) is a brief, self-administered screening instrument [64]. There are seven items, each rated on a five-point scale (0 to 4) yielding a maximum score of 28. A score of eight or greater indicates a probable diagnosis of panic disorder based on validation against the ADIS or the psychiatrist-administered SCID [65]. The PDSS and scoring instructions can be accessed online at <http://www.goodmedicine.org.uk/files/panic,%20assessment%20pdss.pdf>.

Social Anxiety Disorder

The Social Phobia Inventory (SPIN) is a self-administered screening instrument [66]. There are 17 items, each rated on a five-point scale (0 to 4), yielding a total score of 68. A score of 19 or greater indicates a probable diagnosis of social anxiety disorder. There also a shorter, three-item version called the Mini-SPIN [67; 68]. The SPIN and its scoring instructions can be accessed online at http://www.goodmedicine.org.uk/files/social%20anxiety,%20assessment%20spin.full_.tahoma_0.pdf.

Specific Phobia/Agoraphobia

There are currently no validated rating scales for the assessment of specific phobia or agoraphobia. However, screening for specific phobia is simpler than for other anxiety subtypes. The clinician starts by assessing whether the individual has a fear or avoidance of any specific stimulus or situation. If the individual answers affirmatively, then the clinician assesses whether the fear/avoidance is regarded as excessive or unreasonable and whether it interferes with the individual's life. Both criteria must be met for a patient to screen positive.

General (Transdiagnostic) Anxiety Screening

The Overall Anxiety Severity and Impairment Scale (OASIS) is a brief, transdiagnostic screening tool designed to assess for the severity of anxiety in the past week of the individual's life [69]. There are five items, each rated on a five-point scale (0 to 4), yielding a total possible score of 20. A raw score of 8 or greater indicates the presence of anxiety disorder based on validation against anxiety disorder diagnosis using the psychiatrist-administered SCID [70]. Raw scores of 10 and 12 indicate the presence of marked and severe anxiety, respectively, based on validation against the clinician-rated Clinical Global Impression-Severity (CGI-S) scale in a sample of individuals with any anxiety disorder ascertained using the Mini International Neuropsychiatric Interview (MINI) [71].

Scales for Older Adult/Geriatric Use

The Geriatric Anxiety Inventory (GAI) is a 20-item questionnaire designed specifically for older adults (65 years of age and older) [72; 73]. It is a self-report or clinician-administered measure, with each item rated on a binary response scale (agree/disagree), for a total score of 20. Scores of 10 or greater indicate a probable diagnosis of anxiety disorder. The GAI has been translated to more than 20 languages. The GAI has also been validated in clinical and non-clinical samples and in those with cognitive impairment and Parkinson disease [74; 75; 76]. There is also a five-item version called the Geriatric Anxiety Inventory-Short Form (GAI-SF) [77]. Both versions can be accessed online at <http://gai.net.au>.

The Geriatric Anxiety Scale (GAS) is a self-report questionnaire designed for use among adults 65 years of age and older [78]. The GAS contains 30 items, of which only 25 are used to derive a total score. The remaining five questions are used to help the

clinician identify areas of concern for the respondent. Each item is rated on a four-point scale (from 0 to 3), yielding a total score of 100. The GAS consists of three sub-scales assessing somatic (9 items, total possible score: 36), cognitive (8 items, total possible score: 32), and affective (8 items, total possible score: 32) symptoms. There is also a shorter, 10-item version called the GAS-10 [79]. The standard GAS can be accessed at https://gerocentral.org/wp-content/uploads/2013/03/Geriatric-Anxiety-Scale-v2.0_FINAL.pdf, and the GAS-10 can be accessed at <https://gerocentral.org/wp-content/uploads/2013/03/GAS-10-item-version-2015-1-15.pdf>.

Assessment Implications

Compared with younger adults, older adults report fewer and less concrete anxiety symptoms across anxiety subtypes [40; 41; 42; 43; 45]. In addition to this, age-related neurocognitive impairment makes self-reporting a more difficult method of assessment [55]. For example, those with memory impairment can experience stressors that evoke negative effects without leaving memory traces [22; 80]. Although informant report can be a way of effectively gathering information about observable (e.g., physical) symptoms, it is ineffective for identifying unobservable (i.e., subjective) symptoms [81].

EPIDEMIOLOGY

This section addresses the epidemiology of anxiety disorders in older adults, which focuses on the occurrence, determinants, and course and consequences of anxiety disorders in the population. The focus is on the U.S. population, using data from nationally representative surveys. The section begins by addressing the occurrence of anxiety disorders by describing their prevalence and incidence. It then addresses the course and consequences of anxiety disorder, which includes their chronicity, persistence, and comorbidity. Finally, the determinants of anxiety disorders are explored by focusing on risk factors.

PREVALENCE

Prevalence is an estimate of the percent of individuals in the population who meet diagnostic criteria for anxiety disorder, either overall or by subtype. While lifetime prevalence estimates are concerned with the presence of anxiety disorders within the lifetime of individuals, these estimates are typically unreliable because they require respondents to recall prior episodes of anxiety and associated symptoms [82]. Estimating lifetime prevalence in older adults is particularly unreliable, due to general age-related memory deficits [83]. In contrast to lifetime prevalence, period prevalence estimates focus on the presence of anxiety disorder within a given timeframe, typically 12 months. The data in this section are one-year prevalence estimates of anxiety disorder (i.e., whether anxiety disorders were present or absent in the past year of respondents' lives) in nationally representative samples of the U.S. population.

ONE-YEAR PREVALENCE OF ANXIETY DISORDER AMONG ADULTS 55 YEARS OF AGE AND OLDER IN TWO NATIONAL SAMPLES										
Population	Specific Phobia		Social Anxiety Disorder		Generalized Anxiety Disorder		Panic Disorder		Any Anxiety Disorder	
	NESARC	CPES ^a	NESARC	CPES	NESARC	CPES	NESARC	CPES	NESARC	CPES ^b
Total	5%	6%	2%	3%	1%	3%	1%	2%	9%	6%
Age (years)										
55-64	6%	8%	3%	5%	2%	4%	2%	2%	11%	9%
65-74	5%	5%	2%	3%	1%	2%	1%	1%	8%	4%
75+	3%	4%	1%	1%	1%	15%	1%	2%	6%	4%
Sex										
Male	4%	4%	2%	2%	1%	2%	1%	1%	6%	5%
Female	7%	7%	2%	4%	2%	3%	2%	2%	11%	7%
Education										
Less than high school	6%	10%	3%	4%	2%	3%	2%	2%	9%	7%
Completed high school	6%	5%	2%	3%	1%	2%	1%	1%	9%	5%
Some college	6%	6%	2%	3%	2%	4%	1%	2%	9%	9%
Bachelor's degree	4%	4%	1%	2%	1%	2%	1%	1%	7%	5%
Marital status										
Married or cohabiting	5%	5%	2%	2%	1%	2%	1%	1%	8%	4%
Widowed, divorced or separated	6%	8%	2%	5%	2%	4%	2%	2%	10%	9%
Never married	5%	7%	2%	6%	2%	2%	1%	2%	9%	7%

^aSpecific phobia was assessed in a sub-sample of 9,282 respondents from the NCS-R.
^bSpecific phobia was not included in the overall anxiety disorder estimate for the CPES.

Source: [31; 84] Table 5

Anxiety disorders are the most prevalent mental disorders in older adults [54; 84]. The most prevalent subtypes are, in descending order, specific phobia, GAD, social anxiety disorder, and panic disorder. **Table 5** displays the one-year prevalence of anxiety disorders, both overall and by subtype, in the NESARC and the Collaborative Psychiatric Epidemiology Surveys (CPES) of the United States. The prevalence of anxiety disorders is higher among women relative to men, and the prevalence of all anxiety subtypes decreases among persons 75 years of age or older. Previous studies have also reported ethnic differences in prevalence, such that Native and White Americans have the highest prevalence, and Hispanic and Asian Americans have the lowest prevalence of anxiety disorders [85]. Black Americans have a higher or lower prevalence of anxiety disorders depending on subtype; specific phobias and GAD are more prevalent, comparable to Native and White Americans, whereas panic disorder and social anxiety disorder are less prevalent, closer to levels observed in Hispanic and Asian Americans. The prevalence of anxiety disorders does not vary substantially by educational attainment or marital status.

INCIDENCE

The incidence of a disease is defined as the rate at which new cases occur. In contrast to prevalence estimates, which are based on single diagnostic assessments, incidence estimates require at least two diagnostic assessments. The reason for

this is that anyone meeting criteria for an anxiety disorder at any time is counted as a case for the purpose of prevalence estimation, whereas only those individuals who did not have anxiety disorder at time one and who went on to be diagnosed with anxiety disorder at time two are counted as cases for the purpose of incidence estimation, showing that the individuals represent new occurrences of the disorder. The individuals at time one who do not meet criteria for an anxiety disorder are the “risk set” and form the denominator of the incidence ratio, and the individuals at time two or later who meet criteria for an anxiety disorder form the numerator over the period in which the diagnostic assessments were made.

Just as the prevalence of anxiety disorder is higher in older adults than other mental disorders, so too is the incidence, or the rate of newly diagnosed cases [86]. The subtypes with the highest incidence, in descending order, are specific phobia, social anxiety disorder, panic disorder, agoraphobia, and GAD [87; 88]. Data from the Epidemiologic Catchment Area (ECA) study and National Comorbidity Survey (NCS) in the United States, and the Netherlands Mental Health Survey and Incidence Study (NEMESIS) indicate that women have a higher incidence than men [88; 89]. Although anxiety disorder often peaks in young adulthood, there is a smaller but important second peak that occurs in older adulthood [88; 89].

COURSE

The chronicity of a disease refers to its persistence. Persistence is defined here as the percentage of respondents who meet diagnostic criteria for an anxiety disorder at baseline and who then meet criteria again at follow-up. Data from the NESARC indicate that approximately 30% of older adults (55 years of age and older) have persistent cases of anxiety disorder, or chronicity, assessed over a three-year follow-up period. The most persistent subtypes were specific phobia (25%) and GAD (20%), followed by social anxiety disorder (16%) and panic disorder (10%) [31].

There is high co-occurrence between anxiety and other mental disorders, particularly major depression [90; 92; 93]. Panic disorder and GAD have a particularly high comorbidity with mood and substance use disorders in adults [94]. Specific phobia has a strong association with social anxiety disorder and depression [95]. Finally, there are strong associations between social anxiety disorder, GAD, and bipolar disorder [91]. These patterns generally persist among older adults [54]. Anxiety subtypes also have high degrees of overlap [91; 94; 95].



When assessing an adult with possible social anxiety disorder, the National Collaborating Centre for Mental Health recommends that clinicians be aware of comorbid disorders, including avoidant personality disorder, alcohol and substance misuse, mood disorders, other anxiety disorders, psychosis, and autism.

(<https://www.nice.org.uk/guidance/cg159/resources/social-anxiety-disorder-recognition-assessment-and-treatment-pdf-35109639699397>. Last accessed February 17, 2022.)

Level of Evidence: Expert Opinion/Consensus Statement

Data from the NCS suggest that anxiety disorders are also associated with various physical conditions [96]. While panic attacks are associated with vascular conditions, specific phobias are linked with respiratory conditions, and social anxiety disorder with metabolic conditions. Among older adults, there are high rates of anxiety disorders in individuals who have chronic obstructive pulmonary disease (COPD) and/or cardiovascular diseases [35; 97]. The Baltimore ECA study reports an association between blood-injection phobia and vascular complications among individuals with diabetes, which suggests the possibility that fear of blood and injections may interfere with medical treatment [98]. Blood-injection phobia is also associated with respiratory conditions, similar to the data on overall phobias in the NCS sample [99]. The prevalence of blood-injection phobia ranges from 4% to 8% in older adults [36; 100].

CONSEQUENCES

As discussed, anxiety is diagnosed as a disorder only when it is deemed by the individual to be a cause of distress and/or to interfere with daily life. In the NEMESIS study, those with anxiety disorder at baseline had more suicidal ideation and suicide attempts at three-year follow-up, after adjustment for demographic characteristics and past history of mental disorders [101]. Similar associations were observed in cross-sectional studies of the NESARC and NCS-R samples of adults residing in the United States [102].

Importantly, the findings of a 2016 systematic review and meta-analysis of prospective, longitudinal studies suggest that a diagnosis of any anxiety disorder at baseline is not associated with increased risk of all-cause mortality at follow-up [5]. In fact, in a population study of Norwegians, high anxiety symptoms were associated with lower mortality among individuals with depression [103]. In a population study of a 1946 UK birth cohort, individuals who demonstrated lower levels of trait anxiety in adolescence were associated with higher risk of accident mortality at follow-up [104]. Thus, low anxiety (but not high anxiety) is associated with increased mortality risk, and some degree of anxiety is beneficial for survival. Some anxiety likely encourages individuals to engage in preventive health behaviors. For example, women who worry about the possibility of breast cancer are more likely to seek routine screenings, people who are more worry-prone are more likely to vaccinate than those who worry less, and smokers with higher worries about their health have been found to be more likely to quit [105; 106; 107].

RISK FACTORS, RISK ASSESSMENT, AND PREVENTION

The two strongest risk factors for anxiety disorders among older adults are female sex and younger age [84; 108; 109]. However, other risk factors have also been identified. Cigarette smoking is shown to be a major risk factor of anxiety disorder onset, while smoking cessation is associated with reduced anxiety, suggesting that smoking interventions would have a significant effect on anxiety disorder onset [110; 111]. Another important risk factor of anxiety disorder onset in longitudinal studies is the occurrence of adverse life events, such as the ending of a relationship or the injury, illness, or death of a loved one [112; 113; 114].

The presence of adverse events at baseline is associated with an increased risk of overall anxiety disorder onset in older adults. In addition to female sex, history of any mood disorder, and cigarette smoking at baseline, lower levels of educational achievement are associated with higher risk of anxiety disorder onset at follow-up. Although previous studies have reported that excessive anxiety may be a result of licit or illicit substance use or abuse, this has not been replicated in more recent analyses [115; 116; 117]. The association between anxiety disorder and increased substance abuse (including prescription medication) observed in prior studies has been interpreted as evidence of self-medication for emotional distress [118; 119; 120]. A

2019 study assessed the longitudinal association of baseline social anxiety disorder and incident alcohol use disorder at 3- and 10-year follow-up periods in two national samples and did not find evidence of an association between social anxiety and self-medication with alcohol [121].

The prevalence of anxiety disorder is substantially higher in medical versus community settings, and there is a particularly high prevalence of anxiety disorder in individuals with Parkinson disease and among caregivers of older adults [51; 61; 122; 123; 124; 125]. Studies have demonstrated that, in part, the psychological distress (e.g., anxiety and depression) experienced by caregivers is linked to their patients' overall cognitive well-being, patient functional ability, and the reported caregiver burden [126; 127; 128; 129].

TREATMENT

This section will review the available evidence base for the treatment of anxiety disorders. First, preference is given to systematic reviews and network meta-analyses in the general population. However, individual studies are also used in discussion of specific phobia due to a lack of more rigorous research.

PANIC DISORDER

A 2016 network meta-analysis of 54 intervention studies assessed the effectiveness of eight methods of psychological interventions for treating panic disorder with or without agoraphobia [130]. These interventions included [130]:

- Psychoeducation
- Supportive psychotherapy
- Physiological therapies
- Behavior therapy
- Cognitive therapy
- Cognitive behavioral therapy (CBT)
- Third-wave CBT
- Psychodynamic therapies

Researchers found that not one of these treatments was supported as being more efficacious than the others, although any psychological treatment was generally mildly efficacious in comparison with a wait-list control condition [130]. In a subsequent study, the same investigators assessed whether particular components of CBT were associated with better responses to treatment. They reported that face-to-face administration (as compared to self-help) and graded interoceptive exposure to the physiological aspects of the panic response are the most effective features of CBT for treating panic disorder, although it is important to note that the principle of totality applies: the whole of a treatment is more than the sum of its parts [131]. Individual studies addressing treatments for late-life panic disorder have found that both psychological and pharmacological interventions tend to be less efficacious for older adults compared with younger adults [132].

SOCIAL ANXIETY DISORDER

A systematic review and network meta-analysis compared the effectiveness of seven classes of psychological interventions, five classes of pharmacological interventions, and three control groups [133]. Interventions included:

- Promotion of exercise
- Exposure and social skills
- Group CBT
- Individual CBT
- Other psychological therapy (including interpersonal psychotherapy, mindfulness training, and supportive therapy)
- Psychodynamic psychotherapy
- Self-help with or without support

Individual CBT was found to be effective for acute treatment compared with waitlist control groups. Pharmacologic interventions included anticonvulsants, benzodiazepines, monoamine oxidase inhibitors (MAOIs), noradrenergic and serotonergic antidepressants, selective serotonin reuptake inhibitors (SSRIs), and selective norepinephrine reuptake inhibitors (SNRIs). SSRIs and SNRIs were found to be the most effective class of pharmacological treatment compared with placebo control groups [133].

In this study, the promotion-of-exercise intervention was not found to be effective; however, this was not actually an exercise intervention. A 2020 systematic review and network meta-analysis assessing the efficacy of aerobic, resistance, and mind-body training regimens for treating depression reported that actual exercise interventions elicit high levels of treatment compliance and can be effective in reducing depressive symptoms [134]. Thus, similar treatments may also prove to be efficacious for treating anxiety disorders or subtypes.

GENERALIZED ANXIETY DISORDER

The results of two systematic reviews and meta-analyses suggest that psychological therapy has short-term efficacy for treating GAD [135]. A systematic review and network meta-analysis of 27 randomized, double-blind, placebo-controlled studies compared the relative effectiveness of nine pharmacologic treatments of GAD [136]. Although none of the treatments stood out as being clearly more successful than the others, it was concluded that fluoxetine may be preferred for response and remission and sertraline for treatment tolerance. Sertraline is also the most cost-effective pharmacologic treatment of GAD [137]. A separate systematic review and meta-analysis of 27 clinical trials assessed the effectiveness of psychological and pharmacologic treatments for late-life GAD [41]. In this study, benzodiazepines were found to be mildly efficacious relative to placebo, and psychotherapy was found to be mildly efficacious relative to waitlist control groups. A 2016 meta-analysis also reported that CBT is effective for treating GAD in older adults [138].



According to the National Collaborating Centre for Mental Health, the recommended high-intensity psychological intervention for persons with generalized anxiety disorder is cognitive-behavioral therapy (CBT) or applied relaxation.

(<https://www.nice.org.uk/guidance/cg113>. Last accessed February 17, 2022.)

Level of Evidence: Expert Opinion/Consensus Statement

therapy alone. A second form of pharmacologic augmentation for exposure therapy, introduced more than 20 years ago, is the antibiotic D-cycloserine, which is thought to facilitate fear extinction due to its role as an N-methyl D-aspartate (NMDA) receptor agonist [149; 150]. D-cycloserine has also been used to augment exposure therapy for social anxiety disorder, with studies suggesting that this antibiotic can produce a marginal benefit for treating specific phobias and social anxiety disorder when combined with exposure therapy [151]. However, while these studies mention that the antibiotic is of a low dosage, they do not mention that this marginal benefit needs to be traded off against the risk of accelerating antibiotic resistance, which is a pressing global public health challenge. Computational studies suggest that increasing administration of low doses of antibiotics (as these studies suggest doing in conjunction with exposure therapy) accelerates resistance [152; 153].

Although the short-term efficacy of exposure therapy for specific phobias is moderately high, it is important to note that specific phobias are prone to high rates of relapse [139; 140; 154; 155; 156; 157]. Accordingly, studies have sought to eliminate conditioned responses to phobic stimuli over multiple contexts to make for a more successful extinction [158; 159; 160; 161; 162].

TREATMENT IMPLICATIONS

Given that anxiety itself is an adaptive trait, anxiety disorders are better seen as poorly regulated defenses than as defects. As decades of lesion studies indicate, a lack of anxiety may also create non-trivial problems for individuals' lives. Low levels of anxiety are associated with higher mortality risk, and those who report greater worries about particular health problems are likely to seek medical care and take preventative or corrective action [39; 103; 104; 105; 106; 107]. If some degree of anxiety is advantageous, then insufficient and excessive anxiety can both be considered maladaptive.

CONCLUSION

Anxiety facilitates the management of potential future hazards. Even though anxiety is effective at reducing danger, excessive anxiety is often a cause of significant distress and impairment, and anxiety disorders are the most prevalent mental disorders among older adults. Female sex and smoking are the strongest risk factors for late-life anxiety disorders, although adverse life events are also an important factor. About one-third of all cases have considerable chronicity, and therefore prevention is important. Interventions should focus on reducing anxiety to a sufficient, but not excessive, degree.

SPECIFIC PHOBIAS

Exposure therapy is the treatment of choice for specific phobias [139; 140]. This includes in vivo (real-life) and virtual reality exposure to phobic stimuli or situations. Virtual reality exposure therapy was introduced in the 1990s, and although it may have some treatment benefit, it has not been found to have strong efficacy [141]. A one-session exposure therapy treatment for specific phobias was pioneered more than 30 years ago with a suggested duration of two hours and was subsequently used to treat various specific phobia subtypes [142; 143; 144; 145]. More recent studies suggest that one session does not always turn out to be adequate and that multiple sessions are generally more efficacious [140; 146]. However, there may be some cases where the single-session approach is viable.



The National Collaborating Centre for Mental Health recommends against routinely offering computerized CBT to treat specific phobias in adults.

(<https://www.nice.org.uk/guidance/cg159/resources/social-anxiety-disorder-recognition-assessment-and-treatment-pdf-35109639699397>. Last accessed February 17, 2022.)

Level of Evidence: Expert Opinion/Consensus Statement

Pharmacotherapy is not a common treatment for specific phobias. However, studies have sought to supplement exposure therapy using pharmacologic approaches. One such intervention administers cortisol to augment exposure therapy due to its role in interfering with memory for fearful scenarios [147; 148]. Although this treatment shows some efficacy, it does not seem to be particularly advantageous relative to exposure

SELF-ASSESSMENT QUESTIONS
#96690 ANXIETY DISORDERS IN OLDER ADULTS

After reviewing the course, complete the following Self-Assessment Questions. Receive immediate feedback by reviewing the Study Guide provided on page 31. Please record the number of questions you correctly answered on the Evaluation. The Evaluation must be completed in order to receive credit for this course.

This 3 contact hour activity must be completed by February 28, 2025.

1. António Egas Moniz was awarded the 1949 Nobel Prize for treating mental disorders with
 - A) leukotomy.
 - B) psychotherapy.
 - C) holistic medicine.
 - D) pharmacotherapy.
2. Two neural structures that are necessary for anxiety responses are the
 - A) temporal pole and amygdala.
 - B) temporal pole and hippocampus.
 - C) ventromedial prefrontal cortex and amygdala.
 - D) ventromedial prefrontal cortex and hippocampus.
3. The standard for anxiety disorder assessment is the
 - A) ICD.
 - B) DSM.
 - C) structured interview.
 - D) semi-structured interview.
4. The Overall Anxiety Severity and Impairment Scale (OASIS) consists of
 - A) two items.
 - B) five items.
 - C) 12 items.
 - D) 25 items.
5. The most prevalent anxiety disorder subtype in older adults is
 - A) panic disorder.
 - B) specific phobia.
 - C) social anxiety disorder.
 - D) generalized anxiety disorder.
6. Which of the following groups of older adults (55 years of age or older) has the highest prevalence of any anxiety disorder according to data from the NESARC?
 - A) Women
 - B) Persons with a Bachelor's degree
 - C) Adults 75 years of age and older
 - D) Those who are married or cohabitating
7. The most persistent anxiety disorder subtype in older adults is
 - A) panic disorder.
 - B) specific phobia.
 - C) social anxiety disorder.
 - D) generalized anxiety disorder.
8. The most significant modifiable risk factor for anxiety disorder is
 - A) age.
 - B) education.
 - C) adverse events.
 - D) cigarette smoking.
9. Which class(es) of pharmacotherapy is the most effective in the treatment of social anxiety disorder?
 - A) MAOIs
 - B) SSRIs and SNRIs
 - C) Benzodiazepines
 - D) Tricyclic antidepressants
10. The treatment of choice for specific phobias is
 - A) cortisol.
 - B) lithium.
 - C) D-cycloserine.
 - D) exposure therapy.

STUDY GUIDE

COURSE #96690 ANXIETY DISORDERS IN OLDER ADULTS

This Study Guide has been included to provide immediate feedback about your achievement for the course learning objectives. The correct answer for the Self-Assessment Questions and their locations within the text are indicated below.

1. **Antônio Egas Moniz was awarded the 1949 Nobel Prize for treating mental disorders with**
A) *leucotomy. (Neuroanatomy)*
2. **Two neural structures that are necessary for anxiety responses are the**
C) *ventromedial prefrontal cortex and amygdala. (Neuroanatomy)*
3. **The standard for anxiety disorder assessment is the**
C) *structured interview. (Assessment; Structured and Semi-Structured Interviews)*
4. **The Overall Anxiety Severity and Impairment Scale (OASIS) consists of**
B) *five items. (Assessment; Rating Scales; General (Transdiagnostic) Anxiety Screening)*
5. **The most prevalent anxiety disorder subtype in older adults is**
B) *specific phobia. (Epidemiology; Prevalence)*
6. **Which of the following groups of older adults (55 years of age or older) has the highest prevalence of any anxiety disorder according to data from the NESARC?**
A) *Women (Epidemiology; Table 5)*
7. **The most persistent anxiety disorder subtype in older adults is**
B) *specific phobia. (Epidemiology; Course)*
8. **The most significant modifiable risk factor for anxiety disorder is**
D) *cigarette smoking. (Epidemiology; Risk Factors, Risk Assessment, and Prevention)*
9. **Which class(es) of pharmacotherapy is the most effective in the treatment of social anxiety disorder?**
B) *SSRIs and SNRIs (Treatment; Social Anxiety Disorder)*
10. **The treatment of choice for specific phobias is**
D) *exposure therapy. (Treatment; Specific Phobias)*

Pancreatic Cancer

Includes 9 Advanced Pharmacology Hours

This course meets one hour of Pain Management education.

Audience

This course is designed for nurses, physicians, and other members of the interprofessional healthcare team involved in the care of patients with pancreatic cancer.

Course Objective

The purpose of this course is to provide healthcare professionals with the knowledge and skills necessary to recognize and appropriately manage pancreatic cancer in their patients.

Learning Objectives

Upon completion of this course, you should be able to:

1. Outline the epidemiology of and risk factors for pancreatic cancer.
2. Describe the pathophysiology of pancreatic cancers.
3. Discuss recommendations for screening for pancreatic cancer in various patient populations.
4. Describe key aspects of the clinical evaluation of patients with suspected pancreatic cancer.
5. Select the appropriate tools for diagnosis and staging of pancreatic cancer.
6. Apply models of assessing the functional performance status of patients with diagnosed pancreatic cancer.
7. Discuss the role of resection in pancreatic cancer treatment, including most appropriate approaches.
8. Compare and contrast chemotherapy regimens used in the treatment of pancreatic cancer.
9. Describe the use of radiation therapy as a component of pancreatic cancer treatment according to evidence-based guidelines.
10. Evaluate available interventions to manage symptoms and provide palliative care to patients with pancreatic cancer.

Faculty

Mark Rose, BS, MA, LP, is a licensed psychologist in the State of Minnesota with a private consulting practice and a medical research analyst with a biomedical communications firm. Earlier healthcare technology assessment work led to medical device and pharmaceutical sector experience in new product development involving cancer ablative devices and pain therapeutics. Along with substantial experience in addiction research, Mr. Rose has contributed to the authorship

of numerous papers on CNS, oncology, and other medical disorders. He is the lead author of papers published in peer-reviewed addiction, psychiatry, and pain medicine journals and has written books on prescription opioids and alcoholism published by the Hazelden Foundation. He also serves as an Expert Advisor and Expert Witness to law firms that represent disability claimants or criminal defendants on cases related to chronic pain, psychiatric/substance use disorders, and acute pharmacologic/toxicologic effects. Mr. Rose is on the Board of Directors of the Minneapolis-based International Institute of Anti-Aging Medicine and is a member of several professional organizations.

Faculty Disclosure

Contributing faculty, Mark Rose, BS, MA, LP, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Division Planner

Jane C. Norman, RN, MSN, CNE, PhD

Senior Director of Development and Academic Affairs

Sarah Campbell

Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Accreditations & Approvals



JOINTLY ACCREDITED PROVIDER™
INTERPROFESSIONAL CONTINUING EDUCATION

In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Designations of Credit

NetCE designates this continuing education activity for 10 ANCC contact hours.



IPCE CREDIT™

This activity was planned by and for the healthcare team, and learners will receive 10 Interprofessional Continuing Education (IPCE) credits for learning and change.

NetCE designates this continuing education activity for 9 pharmacotherapeutic/pharmacology contact hours.

AACN Synergy CERP Category A.

Individual State Nursing Approvals

In addition to states that accept ANCC, NetCE is approved as a provider of continuing education in nursing by: Alabama, Provider #ABNP0353 (valid through 07/29/2025); Arkansas, Provider #50-2405; California, BRN Provider #CEP9784; California, LVN Provider #V10662; California, PT Provider #V10842; District of Columbia, Provider #50-2405; Florida, Provider #50-2405; Georgia, Provider #50-2405; Kentucky, Provider #7-0054 (valid through 12/31/2023); South Carolina, Provider #50-2405; West Virginia, RN and APRN Provider #50-2405.

About the Sponsor

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

Disclosure Statement

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

How to Receive Credit

- Read the following course.
- Complete the Self-Assessment Questions at the end of the course. (**No test required.**)
- Complete the Evaluation.
- Return your Customer Information/Evaluation and payment to NetCE by mail or fax, or complete online at www.NetCE.com/PA23.
- A full Works Cited list is available online at www.NetCE.com.



Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the study questions and course material for better application to your daily practice.

INTRODUCTION

Pancreatic ductal adenocarcinoma (PDAC) has the worst prognosis of any common cancer. The five-year overall survival rate is approximately 10% and has improved only marginally in five decades [1]. There are four fundamental challenges that underlie the high mortality of PDAC: pancreatic anatomy, aggressive biology, systemic effects, and treatment resistance.

The retroperitoneal position of the pancreas is situated deep within the upper abdomen, behind the stomach, and between the aorta and its major upper abdominal branches. Shielded from detection, the tumors often grow around and encase these vessels, making the cancer inoperable in nearly 85% of patients [2]. With this aggressive cancer, more than 50% of patients have distant metastases at diagnosis, and micrometastases are already present in most patients undergoing resection for apparently localized tumors [2; 3; 4].

At diagnosis, up to 80% of patients with PDAC present with cachexia, a wasting syndrome and physiologic effect of PDAC. Cachexia dramatically weakens patients, limiting their ability to withstand aggressive treatment. The poor treatment tolerance of patients with cachexia is evidenced by decreased survival after resection or chemotherapy [2].

The complex tumor microenvironment and heterogeneity of gene mutations make PDAC one of the most drug-resistant cancers. Most treatment options are ineffective, with rapid progression and low complete responses to the most effective chemotherapy and radiotherapy [1; 4].

Surgical resection of the pancreas with microscopically free margins (R0 resection) followed by chemotherapy remains the only realistic option for remission, but this is potentially achievable in only a fraction of patients [4; 5]. Nonetheless, incremental gains have been increasingly frequent over the past decade, and more substantive gains are anticipated, pending clinical trial results. This course will describe the current standard of care for patients with pancreatic cancer and present information that may help increase earlier detection of this malignancy and improve the symptom burden and quality of life in these patients, regardless of disease stage.

Clinical practice guidelines for patients with pancreatic cancer have been published by the American Society of Clinical Oncology (ASCO), the NCCN (National Comprehensive Cancer Network), the American Society for Radiation Oncology (ASTRO), the European Society for Medical Oncology (ESMO), the National Institute for Health and Care Excellence (NICE), and others [6; 7; 8; 9; 10; 11; 12; 13; 14; 15]. The recommendations are largely concordant on what constitutes multidisciplinary standards of care in the management of pancreatic cancer [2; 16].

PANCREATIC CANCER STAGE AT DIAGNOSIS AND ASSOCIATED SURVIVAL

Stage	Progression at Diagnosis	Five-Year Survival
Localized	11%	41.6%
Regional	30%	14.4%
Distant	52%	3.0%
Unknown	7%	6.5%

Source: [19] Table 1

Most pancreatic cancers arise in the exocrine pancreas (95%). Tumors of the endocrine pancreas (<5%) are distinct from exocrine pancreas cancers and will not be discussed in this course [4].

PDACs account for more than 95% of exocrine pancreatic cancers. PDAC and pancreatic cancer are commonly used as interchangeable terms in the literature and will be in this course [17].

EPIDEMIOLOGY

During 2021 in the United States, an estimated 60,430 people will be diagnosed with pancreatic cancer, which represents 3.2% of all new cancer cases and the 11th most common new cancer diagnosis. The median age at diagnosis is 70 years [18].

Approximately 1.7% of men and women will be diagnosed with pancreatic cancer at some point during their lifetime, based on 2016–2018 data. In 2018, an estimated 83,777 people were living with PDAC in the United States [19].

With an estimated 48,220 deaths in 2021, pancreatic cancer is the third leading cause of cancer death (after lung and colorectal cancer) in both men and women; it is expected to become the second leading cause of cancer death by 2030 [2; 19; 20]. The median age at death is 72 years [18].

Pancreatic cancer stage at diagnosis strongly influences the length of survival, as shown by data from 2011 to 2017 (*Table 1*) [19]. The five-year survival of PDAC, 10.8%, remains the lowest of all common cancers [19; 21].

During 2013–2017, annual pancreatic cancer incidence and mortality rates (per 100,000 persons) were higher among men (14.9 and 12.7) than women (11.6 and 9.6). These rates were highest for Blacks (15.3 and 13.3), followed by non-Hispanic Whites (13.1 and 10.9) and Hispanics. The rates were lowest for Asian/Pacific Islanders and American Indian/Alaska Natives [2].

Since 2010, both incidence and mortality rates increased by an average of 0.3% per year. Underlying these trends is a combination of an aging population, a longer lifespan, and the high prevalence of obesity and diabetes [11; 18]. In 2015, lost earnings from person-years of life lost from pancreatic cancer were estimated at more than \$6 billion [2].

RACIAL SURVIVAL DISPARITIES

In examining PDAC survival disparities over 2004–2015, the unadjusted median overall survival was slightly longer for White patients than Black patients (6.6 months vs. 6.0 months). Decreased survival for Black patients persisted after controlling for sociodemographic parameters. Conversely, controlling specifically for clinical parameters (e.g., disease stage, treatment) found a modest survival advantage for Black patients [22].

Black patients with PDAC present at younger ages with more advanced disease than White patients, possibly suggesting differences in tumor biology. Black patients receive less treatment stage-for-stage and fewer surgeries for resectable PDAC than White patients; these findings may be only partly associated with socioeconomic differences. In one study, when disease stage and treatment were controlled for, Black patients had no decrease in survival compared to other races [22].

Role of Implicit Bias

Health professionals' implicit biases shape behaviors, communications, and interactions, which then produce differences in diagnoses and ultimately treatments and interventions. Implicit biases are subtle and unconscious and may unwittingly produce professional behaviors, attitudes, and interactions that reduce patients' trust and comfort with their provider.

Racial and socioeconomic differences in surgical intervention rates, treatment at high-volume hospitals/centers, and morbidity and mortality rates have been noted, with the largest disparities between Black (and to a slightly lesser extent Hispanic) and White Americans [23]. Several factors are implicated, but implicit biases and insurance status are identified as potentially modifiable contributors.

NON-GENETIC RISK FACTORS

The most common recognized risk factor for pancreatic cancer is cigarette smoking followed by obesity. Others include pancreatitis, diabetes, and family history of pancreatic cancer (*Table 2*) [13; 24]. Periodontal disease is increasingly linked to pancreatic and other gastric cancers. Chronic pancreatitis substantially elevates the risk of developing pancreatic cancer and represents an opportunity for surveillance and monitoring. Most importantly, new-onset hyperglycemia or diabetes is now recognized as an early symptom of PDAC in an otherwise

COMMON RISK FACTORS FOR THE DEVELOPMENT OF PANCREATIC CANCER	
Factor	Relative Risk
Cigarette smoking	1.7-fold to 2.6-fold
Obesity	1.1-fold to 1.5-fold
Diabetes	1.5-fold to 2-fold
Family history	1.7-fold to 2.3-fold
Chronic pancreatitis	13.3-fold
Source: [2]	Table 2

asymptomatic patient. Many recognized risk factors are modifiable for prevention of pancreatic cancer.

Smoking

Cigarette smokers have at least a two-fold greater risk for pancreatic cancer than nonsmokers. The risk increases with the amount of cigarettes consumed and duration of smoking. In heavy smokers with polymorphism in the carcinogen-metabolizing enzyme gene glutathione S-transferase theta 1 (*GSTT1*), the risk is up to five-fold greater [25; 26].

Excess risk decreases with smoking cessation. The risk of pancreatic cancer among current smokers (relative risk: 2.5) decreased 48% two years after smoking cessation, and within 10 to 15 years after cessation, it approximated that of nonsmokers [26].

In the United States, estimates indicate that 11% to 32% of deaths from PDAC are attributable to tobacco smoking. It is estimated that cessation of smoking could eliminate up to 25% of pancreatic cancer deaths [24; 26].

Alcohol Consumption

Limited evidence suggests alcohol consumption may be associated with risk of developing PDAC, but findings of population-based studies are inconsistent. In pooled cohort data of 1.5 million light, heavy, or never-drinkers, heavy drinkers had a greater relative risk of developing PDAC than never-drinkers (relative risk: 1.29) or light drinkers (relative risk: 1.36). Light drinkers had no difference compared to never-drinkers (relative risk: 0.96) [27].

Smoking and Drinking

Most studies have assumed additivity between average effects of smoking and alcohol and oversimplified their impact on burden of pancreatic cancer. However, the combined effect of smoking and total alcohol intake on risk of PDAC is likely non-additive. It appears that only heavy consumption of liquor (but not wine or beer) increases the risk of PDAC in ever-smokers [27].

Obesity

A number of studies have associated obesity with a higher incidence of pancreatic cancer. Obesity (defined as a body mass index [BMI] >30) during early adulthood was associated with a greater risk of PDAC and younger age of disease onset. Tumorigenesis is enhanced by excess adipose tissue. Obesity is associated with a 20% to 40% higher mortality rate from PDAC, and obesity at an older age is associated with lower overall survival [13; 28].

Although BMI is widely used as a marker for general adiposity, visceral obesity has a stronger correlation to metabolic syndrome, insulin resistance, and certain gastrointestinal (GI) malignancies. The close proximity to visceral organs and drainage via the portal system may explain the strong correlation of inflamed visceral adipose tissue (VAT) in obese subjects with metabolic dysfunction and pancreatic cancer [29].

Diet

There is some evidence that higher consumption of red/processed meat is associated with elevation in pancreatic cancer risk, but other studies have failed to identify dietary risk factors for PDAC [11]. Pancreatic cancer incidence may be lower in persons with higher intake of fresh fruits and vegetables rich in folate and lycopenes (e.g., tomatoes) [30].

A link between vitamin D and risk for pancreatic cancer is inconsistent, but some data suggest low plasma 25-hydroxyvitamin D levels may increase the risk for pancreatic cancer, especially in those with low retinol/vitamin A intake [31]. Coffee and tea consumption are not associated with pancreatic cancer risk, despite early reports to the contrary [24].

Systemic/Nonmodifiable Risks

Numerous studies and meta-analyses have found systemic/nonmodifiable factors that increased the relative risk, hazard ratio, or odds ratio of developing pancreatic cancer. These include individuals with greater height (relative risk: 1.81); individuals with blood groups A, AB, and B (hazard ratio: 1.32, 1.51, and 1.72, respectively); and patients with hepatitis B infection (odds ratio: 1.50) or systemic lupus erythematosus (hazard ratio: 1.43). Biologic explanations for some of these associations are not yet understood, and some data may have potential confounders. Infectious etiologies warrant more investigation [11; 32].

Periodontal Disease

Periodontitis describes a chronic inflammatory response to a disease-associated, multispecies bacterial community in the subgingival region. Periodontal disease is associated with pancreatic cancer, even when controlling for gender, smoking, BMI, diabetes, and alcohol consumption [33]. The inflammatory processes of periodontitis occur locally, but systemic dissemination of inflammatory mediators, subgingival species, and bacterial components contribute to digestive cancers (including PDAC) by activating proinflammatory pathways, inducing gene

expression related to cell proliferation, apoptosis, and immune responses linked to carcinogenesis, cell migration, invasion, and metastasis [34].

Chronic Pancreatitis

A high-risk subgroup for PDAC are patients with chronic pancreatitis, often secondary to chronic alcohol use disorder, smoking, hypertriglyceridemia, diabetes, or renal failure [2]. Patients with chronic pancreatitis show a 26-fold increase in risk of developing PDAC. This risk increases with duration. Among patients with chronic pancreatitis of 20 years' duration, approximately 5% will progress to PDAC.

Concomitant smoking enhances the risk of neoplastic progression [2; 35]. Hereditary pancreatitis further increases the risk of pancreatic cancer by more than 50-fold. In these individuals, the cumulative risk of pancreatic cancer by age 70 years is 40% [24].

Long-Standing Diabetes

Pancreatic cancer has complex relationships with diabetes and obesity that are only recently becoming understood. A population cohort study underscored the complex relationship between metabolic abnormalities and PDAC. Glycemic status, insulin resistance, and hyperinsulinemia were independently associated with an increased risk of pancreatic cancer mortality, even in individuals without diabetes [36].

The association between pancreatic cancer and diabetes was noted as early as 1833, clearly documented by the 1930s, and characterized in a large cohort of patients with pancreatic cancer from Mayo Clinic in 1958 [37]. Several meta-analyses have greatly refined the risk-factor status of diabetes.

Long-standing (i.e., more than five years) diabetes (both type 1 and type 2) is associated with increased risk of developing PDAC [13]. The overall risk for PDAC increases 4- to 7-fold in those with diabetes of a duration less than three years [38]. The relative risk associated with diabetes levels off after five years, with a 1.5-fold greater risk [39]. Increased baseline hemoglobin A1C (HbA1C) levels correlate with subsequent development of PDAC [40].

Long-standing diabetes modestly increases the risk of PDAC, which decreases with diabetes duration [11; 37]. The initial three-year period after diabetes diagnosis is high risk for PDAC, as confirmed by prospective pancreatographic screening [41].

With diabetes medications, insulin use has been associated with increased risk of PDAC, but this finding is attributed to reverse causality [11; 42]. Metformin use in patients with diabetes and PDAC was associated with improved two-year survival (30.1% vs. 15.4%) and median overall survival (15.2 months vs. 11.1 months) in patients without metastases [43]. One metformin study reported negative findings [44].

Long-standing diabetes in patients who develop PDAC is associated with significantly lower overall survival (14.4 months vs. 21.7 months) and significantly higher mortality (harm ratio: 1.52) compared with patients without diabetes who develop PDAC [11; 45].

Postpancreatitis Diabetes Mellitus

Diabetes of the exocrine pancreas (formerly type 3c diabetes) is the second most common type of new-onset diabetes in adults (behind type 2 diabetes) [42]. Acute or chronic pancreatitis is one of the most prevalent risk factors for PDAC and the most frequent cause of diabetes of the exocrine pancreas. Pancreatitis leads to postpancreatitis diabetes mellitus in up to 83% of patients [42]. In a registry study involving 139,843 individuals, the proportion of pancreatic cancer was 3.1% among those with postpancreatitis diabetes mellitus, compared with 2.3% in those with type 2 diabetes followed by pancreatitis, 2.0% in those with pancreatitis alone, and 0.6% in individuals with type 2 diabetes alone [42].

Prediagnostic Metabolic and Soft Tissue Changes

Numerous studies have identified new-onset diabetes, weight loss, and soft tissue changes in patients with PDAC at diagnosis, but their inter-relationship and connection to PDAC remained unaddressed. From 2000 through 2015, temporal changes in the five years preceding PDAC diagnosis of 219 patients diagnosed with PDAC were compared to 657 controls [46]. From 60 to 30 months before PDAC diagnosis, patients did not significantly differ from controls. However, starting at 30 months prediagnosis, PDAC showed three distinct metabolic phases, each marked by onset and significant progressive worsening of one or more metabolic abnormalities [46]:

- Phase 1, hyperglycemia (30 to 18 months before PDAC diagnosis): A significant proportion of patients develop hyperglycemia, without soft tissue changes.
- Phase 2, pre-cachexia (18 to 6 months before PDAC diagnosis): Decreases in serum lipids, weight loss, and the first soft tissue change (subcutaneous abdominal tissue loss) are seen. A profile appears of advanced prediabetes (i.e., fasting blood glucose 120–126 mg/dL or A1c of 6% to 6.5%). In type 2 diabetes, this is associated with weight gain and hyperlipidemia due to insulin resistance. In PDAC, decreases in weight and serum lipids despite rising glucose levels are paradoxical.
- Phase 3, cachexia (less than 6 months before PDAC diagnosis): Onset of muscle loss, visceral adipose tissue loss, and decreasing high-density lipoprotein. Continued decreases in all other serum lipids, subcutaneous abdominal tissue, and weight. Fasting blood glucose continues rising.

Based on evidence of increases in body temperature before PDAC diagnosis, browning and loss of subcutaneous abdominal tissue is estimated to begin 18 months before PDAC. Browning of white abdominal tissue is a mechanism of subcutaneous abdominal tissue loss in cancer; its purpose is to generate heat [46].

Symptoms of cachexia and muscle loss (e.g., anorexia, fatigue, reduced exercise tolerance) appear shortly (less than six months) before PDAC diagnosis. The onset of objective weight loss precedes PDAC diagnosis by one year or more. New-onset diabetes appears a median of six to nine months before PDAC diagnosis [46].

Pancreatic Cancer Cachexia and Diabetes

Cancer cachexia is a paraneoplastic syndrome characterized by pronounced weight loss and muscle wasting triggered by cancer-induced systemic inflammation [47]. Cachexia develops in about 80% of patients with PDAC during the disease course, often before the tumor is clinically apparent. Cachexia negatively impacts treatment response and survival, and one-third of patients with PDAC die from cachexia-associated complications, including impaired immunity and cardiopulmonary dysfunction. No curative treatments exist [47].

Pancreatic cancer-associated diabetes mellitus might be a major contributor to PDAC-induced cachexia. The co-occurrence is frequent, and the relationship between pancreatic cancer-associated diabetes and PDAC-induced cachexia was clarified in a 2020 study [47]. Compared with patients without pancreatic cancer-associated diabetes, those with pancreatic cancer-associated diabetes did not have a higher risk of cachexia, a greater degree of weight loss, or lower skeletal muscle mass. Among patients with cachexia, weight loss and skeletal muscle mass were comparable between patients with and without pancreatic cancer-associated diabetes. Fasting blood glucose levels and PDAC-derived diabetogenic factors did not correlate with weight loss or muscle mass or predict cachexia in patients with pancreatic cancer-associated diabetes. A notable finding was the consistently high prevalence of cachexia and muscle wasting regardless of tumor size and stage in PDAC [47]. These results argue against pancreatic cancer-associated diabetes and hyperglycemia in mediating PDAC-induced cachexia.

Cancer cachexia is characterized by systemic inflammation with resultant skeletal muscle breakdown and increased circulating amino acids to support tumor growth. Pancreatic cancer-associated diabetes is a metabolic strategy by PDAC to fuel tumor growth. PDAC cells have a high demand for glucose (termed “glucose addiction”); hyperglycemia promotes invasion and migration of PDAC cells. PDAC-induced cachexia and pancreatic cancer-associated diabetes are distinct metabolic reprogramming induced by PDAC cells to secure amino acids and glucose for tumor growth [47].

Unexplained weight loss/cachexia is a clue to occult PDAC, but a modality that can identify PDAC-induced cachexia is needed to take advantage of this screening opportunity [47]. Optimizing glycemic control may not alleviate weight loss or muscle wasting, and therapies targeting mediators of pancreatic cancer-associated diabetes may not protect against the development of cachexia [47]. Management of cachexia in patients with PDAC is discussed in detail later in this course.

PATHOPHYSIOLOGY

PDAC is caused by somatic (acquired) and germline (inherited) mutations in specific cancer-associated genes. In PDAC, the accumulation of multiple combinations of gene mutations significantly perturbs major signaling pathways, leading to a malignant phenotype [13; 48; 49; 50].

Like most solid tumors, PDACs are driven by mutations that disrupt intra- and extracellular networks that normally restrain abnormal growth, proliferation, survival, and invasion [51]. Four major genetic drivers are fundamental in nearly all PDACs. These involve mutational activation of the oncogene *KRAS*, and mutational inactivation of the tumor suppressor genes *CDKN2A*, *TP53*, and *SMAD4* [3; 50; 52; 53]. Inactivation of genome maintenance genes that repair DNA damage is a third broad type of mutation in PDAC.

PRIMARY MUTATIONAL DRIVERS IN PDAC

KRAS encodes a GTPase molecule that acts as a transducer for growth factor receptors on the cell surface. *KRAS* mutations dysregulate intrinsic GTPase activity, stimulating downstream pathways that drive uncontrolled cellular proliferation, angiogenesis, suppression of apoptosis, and evasion of immune response [54].

CDKN2A encodes the proteins p16 and p14ARF, which are both cell-cycle regulators. With loss of *CDKN2A* gene function, inactivation of p16 results in unchecked cell cycle progression and enhanced tumor cell proliferation [3; 49]. *TP53* encodes the protein p53, called the “guardian of the genome,” which plays a central role in DNA repair, cell cycle arrest, and induction of apoptosis in response to DNA damage or cellular stress [55].

Inactivation of p53 (loss of function mutation) allows DNA damage to go unchecked with failed apoptosis and unregulated G1/S cell cycle transition. Mutant p53 can also gain pro-oncogenic activities (gain-of-function mutation), promoting cell proliferation, survival, angiogenesis, and metastases [54].

SMAD4 encodes the protein Smad4, a downstream effector of transforming growth factor-beta (TGF- β) signaling pathway. *SMAD4* inactivation and loss of Smad4 promotes cancer progression by removing the early growth inhibitory effect of the TGF- β pathway and is associated with higher rates of distant metastasis and poorer prognosis [54].

MUTATIONAL SEQUENCE OF PDAC DEVELOPMENT

Through pathways and somatic mutations that differ modestly in each lesion, PDAC develops from precancerous precursor lesions: pancreatic intraepithelial neoplasia (PanIN), intraductal papillary mucinous neoplasm (IPMN), and mucinous cystic neoplasms (MCNs). The most common are PanINs (approximately 90%), and the least common are MCNs. However, all precursor lesions have key similarities [4; 48; 50]:

- Early oncogene mutations initiate tumorigenesis.
- Later loss of tumor suppressor genes drive tumor progression, high-grade dysplasia, and invasive cancer.
- Increasing grades of dysplasia are associated with accumulation of somatic mutations in key driver genes.

Pancreatic Intraepithelial Neoplasia (PanIN)

PDAC develops in PanINs through a specific process [56]. First, mutational *KRAS* activation initiates pancreatic carcinogenesis. With tumor suppressor inactivation, cancer progresses. *CDKN2A* or *SMAD4* are implicated in locally destructive disease; *TP53* is involved in metastatic seeding; and concurrent *SMAD4* and *TP53* are often present in locally or metastatic dominant disease. IPMNs and MCNs often share the driver gene mutations and sequence of PanINs, but also show specific patterns.

Intraductal Papillary Mucinous Neoplasms (IPMN)

More than 90% of IPMNs are marked by activating mutations in the oncogene *GNAS* and/or inactivating mutations in the tumor suppressor gene *RNF43* [48; 53; 54]. *GNAS* mutation causes constitutive activation of adenylyl cyclase, with downstream effects driving proliferation. *RNF43* encodes E3 ubiquitin-protein ligase, which functions as a tumor suppressor in the Wnt-signaling pathway. After the initiating oncogene mutation, the progression of IPMN resembles PanIN.

Mucinous Cystic Neoplasms (MCN)

RNF43 mutation is also a prevalent event in MCNs (50%). As in PanINs, genetic changes accumulate with higher grade of dysplasia and invasiveness [48; 53; 54].

NATURAL HISTORY OF PDAC ONCOGENESIS

The PanIN Progression Model has been critical in shaping the perspective of how PDAC develops and progresses over the past two decades. PDAC arises through a specific sequence of genetic alterations over a gradual progression from early PanIN to late-stage metastatic disease [57; 58; 59].

The timeframe of PanIN progression has also been established. Based on computational modeling using autopsy cases, the estimated average time interval from initiation in normal cells to invasive ability (11.7 years), metastatic dissemination (6.8 years) and death (2.7 years) corresponds to an average of about 21 years from the initiating mutation until a patient's death [17].

Most cases with PDAC are diagnosed toward the end of this lifetime span, suggesting that poor prognosis is a result of late diagnosis in the natural history of PDAC, and that a golden opportunity of two or three years exists to diagnose “early” pancreatic cancer (i.e., Stage 0 or I) [60].

Chromothripsis, a recently identified phenomenon, is a catastrophic event causing tens to thousands of chromosomal rearrangements. Faced with hundreds of DNA breaks, the cell's DNA repair machinery attempts to rescue the genome, but the result bears little resemblance to its original structure [61; 62]. This genomic disruption can drive the development of cancer through DNA copy number changes, including deletion of tumor suppressor genes and increased copy number (amplification) of oncogenes [61].

A 2016 study of more than 100 whole genomes from pancreatic cancer tumors found evidence of at least one chromothripsis event in 65% of tumors, and most copy-number changes seemed to occur after such catastrophic genetic events. With evidence of chromothripsis in some PDACs and nongradual tumorigenesis that defies the established mutational sequence, a punctuated equilibrium model was proposed, dividing tumor development into two major events [63]:

- A cancer-initiating event: PDAC pre-neoplasms acquire extensive mutation burden but remain non-invasive over a prolonged preneoplastic phase.
- A cataclysmic cancer-transforming event: Chromothripsis induces DNA copy number changes, creating genomic instability and generating invasive clones with rapid dissemination and colonization of distant sites. Why chromothripsis occurs in PDAC is not yet understood.

Non-Genetic Mechanisms

Rather than being uniformly aggressive, PDAC demonstrates clinical (e.g., variable patient survival) and disease (e.g., variable chemotherapy sensitivity) heterogeneity [64; 65]. The first whole-genome description of PDAC in 2008 prompted great effort to advance a patient-tailored precision medicine approach that could better address this heterogeneity. Genetic alterations and molecular subtypes in PDAC were characterized and published. PDAC was shown mutationally dominated by the four driver genes and homogeneous. In general, the findings importantly informed the biology and familial predisposition of PDAC.

However, by 2019 it was apparent that PDAC disease heterogeneity cannot be explained by genetic mutations alone, and non-genetic mechanisms, including epigenetics and the tumorigenic microenvironment, were the path forward [21; 56; 59; 62; 64; 65; 66; 67].

PANCREATIC CANCER SUSCEPTIBILITY SYNDROMES AND MUTATIONS

Category	Specific Syndromes and Germline Mutations
Gastrointestinal tract cancers	Lynch syndrome, also termed hereditary nonpolyposis colorectal cancer (<i>MLH1</i> , <i>MSH2</i> , <i>MSH6</i> , <i>PMS2</i>) Peutz-Jeghers syndrome (<i>STK11/LKB1</i>) Familial adenomatous polyposis (<i>APC</i>)
Solid tumor cancers	Hereditary breast/ovarian syndrome (<i>BRCA1/2</i> , <i>PALB2</i>) Familial atypical multiple mole melanoma syndrome (<i>CDKN2A</i>) Li-Fraumeni syndrome (<i>TP53</i>)
Chronic pancreatitis-associated syndromes	Hereditary pancreatitis (<i>PRSS1</i> , <i>SPINK1</i>) Cystic fibrosis (<i>CFTR</i>)
Neurodegenerative disease	Ataxia-telangiectasia (<i>ATM</i>)
<i>Source: [48; 54]</i>	

Table 3

Epigenetic Factors

Broadly speaking, epigenetic changes influence gene expression, without altering the DNA sequence, through modifications of DNA or chromatin structures [4]. In PDAC, these include: DNA methylation and non-coding RNAs (ncRNAs).

Gene expression in PDAC can be silenced through non-mutational inactivation by aberrant promoter methylation, including the driver gene *p16/CDKN2A* [49]. Aberrant ncRNA expression plays a considerable role in initiation, proliferation, and chemo-resistance of PDAC. Oncogenic microRNA-21 promotes both cell proliferation and apoptosis and targets negative regulators of *KRAS*, which further enhances signaling by this oncogene [50; 54].

Pancreatic Tumor Microenvironment

Pancreatic cancer tissue is comprised of PDAC cells and dense fibrotic stromal (stellate) cells. The stroma consists of extracellular matrix and non-neoplastic (e.g., fibroblastic, vascular, immune) cells [3]. Also described as PDAC fibrosis, the stroma makes up most of the tumor mass. Its importance beyond a physical barrier to drug penetration was not historically considered. Recognized only recently, the entire neoplastic tissue, both tumor cells and stroma, create a pancreatic tumor microenvironment that crucially facilitates PDAC growth, survival, and treatment failure [21; 51; 68].

Pancreatic cancer progresses in tandem with a stromal reaction, characterized by extensive deposition of extracellular matrix, recruitment and activation of cancer-associated fibroblasts, and high interstitial fluid pressures that compress blood vessels, causing hypoperfusion, hypovascularity, and hypoxia [21; 69]. Extracellular matrix remodeling biomechanically induces intracellular signaling and tumor-stellate cell crosstalk. PDAC cells signal to stellate cells and recruit macrophages and immune suppressor cells. In turn, stellate cells secrete factors that promote PDAC cell proliferation and migration and suppress apoptosis [51]. Biochemical activation of signaling pathways that regulate PDAC cell survival and metastasis

promotes tumor growth, immunosuppression, disease progression, epithelial-mesenchymal transition (a key step of the metastatic cascade) and invasive potential, and chemotherapy resistance [3; 21; 69].

Exosomes (a macromolecule involved in RNA degradation) released by PDAC cells accumulate in other tissues to create a premetastatic niche by activating stellate cells and inducing remodeling of the host extracellular matrix, which facilitates cancer cell invasion and growth [59; 69].

HEREDITARY PDAC

In addition to the somatic mutations driving pancreatic tumorigenesis in all PDACs, specific germline variants also contribute to PDAC in some patients [48]. In many of these germline mutations, the oncogenic mechanism involves inactivation of DNA damage repair genes [49].

There are two broad categories of inherited risk for PDAC [26; 70; 71]:

- Genetic predisposition or hereditary pancreatic cancer: Germline mutations in PDAC susceptibility genes are present.
- Familial pancreatic cancer: Familial clustering of PDAC (i.e., at least one pair of affected first-degree relatives) without known germline mutations

Sporadic PDAC is when both factors are absent. However, mutations in known pancreatic cancer susceptibility genes are found in 5% to 10% of patients with apparently sporadic pancreatic cancer.

Inherited Cancer Susceptibility Syndromes and Germline Mutations

Several genetic syndromes are associated with specific genetic alterations with an increased risk for pancreatic cancer (Table 3) [48; 54]. Germline mutations in familial atypical multiple mole melanoma syndrome (*CDKN2A*) and Li-Fraumeni syndrome (*TP53*) are core gene drivers in sporadic PDAC.

PANCREATIC CANCER RISK IN PREDISPOSITION AND INHERITED CANCER SYNDROMES				
Syndrome	Gene(s)	Risk of PDAC		Other Cancers
		Relative	Lifetime	
General population	-	1	0.5%	-
Hereditary breast/ovarian cancer	<i>BRCA1</i>	2 to 3	1.2% to 2%	Breast, ovarian, prostate
	<i>BRCA2</i>	3.5 to 10	2% to 10%	
	<i>PALB2</i>	15	5% to 10%	
Familial atypical multiple mole melanoma	<i>CKDN2A</i>	13 to 36	10% to 30%	Melanoma
Peutz-Jeghers	<i>STK11</i>	75 to 125	11% to 66%	GI, lung, breast, reproductive
Hereditary nonpolyposis colon cancer (Lynch II)	<i>MLH1, MSH2, MSH6</i>	8 to 10	3.7% to 10%	Colorectal, ovary, uterine, upper GI, urinary tract
Li-Fraumeni	<i>TP53</i>	7	unknown	Breast, brain, adrenal
Familial adenomatous polyposis	<i>APC</i>	4.5	Less than 5%	Colon, upper GI, thyroid, brain
Ataxia telangiectasia	<i>ATM</i>	8 to 9	1% to 5%	Breast, prostate
Hereditary pancreatitis	<i>PRSS1, SPINK1</i>	50 to 82	25% to 44%	-
Cystic fibrosis	<i>CFTR</i>	5	Less than 5%	-
Familial pancreatic cancer ^a	1 first-degree relative	4.6	-	-
	2 first-degree relatives	6.4	-	-
	3 first-degree relatives	32	-	-

^aRisk determined by number of affected first-degree relatives rather than specific gene.

Source: [49; 54; 70; 73]

Table 4

Peutz-Jeghers syndrome is caused by germline inactivation of *STK11*, a tumor suppressor gene. Somatic *STK11* mutations are observed in approximately 4% of pancreatic cancers, suggesting *STK11* inactivation plays a role in both sporadic and familial forms [49].

Familial Pancreatic Cancer

An estimated 10% to 15% of all pancreatic cancers are attributable to genetic causes. Pancreatic cancer aggregates in some families; 5% to 10% of individuals with pancreatic cancer have a family history of the disease [26; 70; 72]. Familial pancreatic cancer represents 90% of all hereditary PDAC cases. The relative risk of PDAC increases with the number of affected first-degree relatives.

A specific gene defect responsible for familial pancreatic cancer has not been identified, but a rare autosomal-dominant gene may be responsible, putting 0.4% to 0.7% of the population at risk for developing PDAC [26; 70; 72]. Details about the relative and lifetime risks of PDAC, and the other prevalent cancers associated with specific germline mutations in cancer susceptibility syndromes and familial pancreatic cancer, are summarized in **Table 4**.

PANCREATIC CANCER SCREENING

With the low population incidence of PDAC (lifetime risk: 1.3%), absence of biomarker screening targets, and high cost of sensitive imaging methods, the U.S. Preventive Services Task Force recommended against screening for pancreatic cancer in asymptomatic adults in 2019, reaffirming its previous conclusion in 2004 [74]. As population screening to achieve earlier detection and intervention of PDAC is not currently feasible, other approaches for this objective have been identified.

In Australia, public awareness campaigns have highlighted the often vague symptoms of PDAC and encouraged individuals to seek medical attention early. Underscoring this point, one study found that many people who were ultimately diagnosed with PDAC were falsely reassured by the subtle, intermittent nature of their symptoms over the preceding months [75; 76].

As a relatively rare cancer, many primary care providers will only see a PDAC case every few years, making it imperative to elevate awareness of early PDAC signs and symptoms among these professionals. A retrospective case-control study in primary care found that patients sought medical attention 18 times on average in the period preceding their pancreatic cancer diagnosis. PDAC was associated with 11 alarm symp-

**INTERNATIONAL CANCER OF THE PANCREAS SCREENING (CAPS) CONSORTIUM
CONSENSUS ON SCREENING FOR PANCREATIC CANCER IN PATIENTS
WITH INCREASED RISK FOR FAMILIAL PANCREATIC CANCER**

What is the goal of pancreatic surveillance?		
The primary goal is to prevent the emergence of and death from pancreatic cancer by identifying and treating stage I pancreatic cancer (resected with negative margins) and pancreatic cancer precursor lesions with high-grade dysplasia (PanIN or IPMN).		
Who should be screened?		
All patients with Peutz-Jeghers syndrome (carriers of a germline <i>LKB1/STK11</i> mutation) All carriers of a germline <i>CDKN2A</i> (p16) mutation Carriers of a germline <i>BRCA2</i> , <i>BRCA1</i> , <i>PALB2</i> , <i>ATM</i> , <i>MLH1</i> , <i>MSH2</i> , or <i>MSH6</i> gene mutation with at least one affected first-degree relative Individuals with at least one first-degree relative with pancreatic cancer who in turn also has a first-degree relative with pancreatic cancer (familial pancreatic cancer kindred)		
At what age^a should pancreatic surveillance begin?		
Familial pancreatic cancer kindred	Start at 50 or 55 years of age, or 10 years younger than the youngest affected blood relative	
Mutation carriers	For <i>CDKN2A</i> and Peutz-Jeghers syndrome, start at 40 years of age	
	For <i>BRCA2</i> , <i>ATM</i> , <i>PALB2</i> , <i>BRCA1</i> , and <i>MLH1/MSH2</i> , start at 45 or 50 years of age, or 10 years younger than the youngest affected first-degree relative	
What tests and indications?		
Indication	Interval	Test(s)
Routine	At baseline and during follow-up	MRI/MRCP and endoscopic ultrasound Fasting blood glucose and/or HbA1c
Concerning abnormalities for which immediate surgery is not indicated	After 3 to 6 months	Repeat follow-up testing
No abnormalities or only non-concerning abnormalities (e.g., pancreatic cysts without worrisome features)	After 12 months	Repeat follow-up testing
If concerning features on imaging	Upon indication	Serum CA 19-9
Solid lesions of ≥ 5 mm Cystic lesions with worrisome features Asymptomatic main pancreatic duct strictures (with or without mass)	Upon indication	Endoscopic ultrasound-guided FNA
Solid lesions, regardless of size Asymptomatic main pancreatic duct strictures of unknown etiology (without mass)	Upon indication	CT
Positive FNA and/or a high suspicion of malignancy on imaging	Upon indication	Surgery ^b
^a Age to initiate surveillance depends on gene mutation status and family history. There is no consensus on the age to end surveillance.		
^b When surgery is indicated, it should be oncologic radical resection at a specialty center.		
CA 19-9 = carbohydrate antigen 19-9; CT = computed tomography; FNA = fine-needle aspiration; HbA1c = hemoglobin A1c; IPMN = intraductal papillary mucinous neoplasm; MRI/MRCP = magnetic resonance imaging/magnetic retrograde cholangiopancreatography = PanIN: pancreatic intraepithelial neoplasia.		
Source: [70; 71]		Table 5

toms; back pain, lethargy, and new-onset diabetes were unique features of PDAC [75; 77].

Specific screening efforts in PDAC have focused on identifying high-risk individuals [48]. In 2020, the International Cancer of the Pancreas Screening (CAPS) Consortium updated its consensus recommendations for the management of individuals with increased risk of pancreatic cancer based on family history or germline mutation status [71]. For selected high-risk individuals, pancreatic surveillance is recommended to detect and resect early pancreatic cancer and its high-grade precursors (**Table 5**). No consensus was reached on whether surveillance should be performed for hereditary pancreatitis.

However, it is important to remember that among patients with PDAC unselected for their family history of pancreatic cancer who had a germline susceptibility gene mutation, only 10% of these patients had a family history of pancreatic cancer, and most did not have a cancer family history to suggest an inherited cancer syndrome. Because family history remains one of the best predictors of future pancreatic cancer risk, routine gene testing of patients with newly diagnosed PDAC and their families may yield significant clinical benefits [78].

Genetic counseling of patients before and after any genetic testing is essential, to provide understanding and reassurance and to avoid harm. A challenge to less restrictive testing of patients with new PDAC is there are not enough genetic counselors to provide this service; this shortage of expertise applies to other cancers as well [78].

GERMLINE AND SOMATIC TESTING AND MOLECULAR ANALYSIS

With strong consensus that benefits outweigh harms, in 2018 the ASCO recommended germline genetic testing for patients with PDAC, even if family history is unremarkable, if an informative result could directly benefit the patient or their family members [73]. This stance was adopted in 2020 by the NCCN. Consensus has subsequently expanded.

All patients with pancreatic cancer should have germline testing and gene profiling offered as quickly as possible after diagnosis; the implications for first-line therapy and beyond are significant [79; 80]. The 2020–2021 ASCO and NCCN recommendations are for all patients with PDAC to receive germline genomic testing using comprehensive gene panels for hereditary cancer syndromes, and targeted (somatic) profiling of tumor tissue using next-generation sequencing [10; 11]. Patients with locally advanced or metastatic PDAC should have available tumor tissue tested for DNA mismatch repair deficiency (dMMR) and microsatellite instability–high (MSI-H) status. It is also recommended that these patients undergo testing for actionable somatic mutations, including fusions (*ALK*, *NRG1*, *NTRK*, *ROS1*), mutations (*BRAF*, *BRCA1/2*, *HER2*, *KRAS*, *PALB2*), and mismatch repair deficiency (dMMR).

CLINICAL EVALUATION OF PANCREATIC CANCER

Most pancreatic cancers (approximately 75%) originate in the head of the pancreas and typically metastasize to regional lymph nodes first, then to the liver. PDAC can also directly invade surrounding visceral organs (e.g., duodenum, stomach, colon); metastasize to any surface in the abdominal cavity via peritoneal spread where development of ascites carries an ominous prognosis; or spread to the skin as painful nodular metastases. By the time of diagnosis, 85% to 90% of patients have locally advanced tumors that have involved retroperitoneal structures, spread to regional lymph nodes, or metastasized to the liver or lung [2; 13; 24; 81].

Early-stage pancreatic cancer is notoriously difficult to diagnose. The most common symptoms in a series of patients diagnosed with PDAC were fatigue (86%), weight loss (85%), anorexia (83%), abdominal pain (79%), epigastric pain (71%), jaundice (56%), nausea (51%), diarrhea (44%), pruritus (32%), and steatorrhea (25%) [82].

Abdominal pain, jaundice, and weight loss are nonspecific, subtle in onset, and easily attributed to other processes. Unless the healthcare provider has a high index of suspicion for the possibility of underlying pancreatic carcinoma, this can make it difficult to know when to escalate a workup, as PDAC lacks a specified diagnostic algorithm [2; 24].

Development of abdominal pain, jaundice, or weight loss in the context of newly diagnosed diabetes, family history of PDAC, or history of pancreatitis should trigger inclusion of PDAC in the differential diagnosis [2]. Furthermore, past three-year onset of diabetes or ongoing hyperglycemia with significant weight loss and decreasing serum lipids should be considered a potential PDAC, even if abdominal pain or jaundice are absent, with urgent referral a priority.

As noted, pancreatic cancer-associated diabetes and pancreatic cancer cachexia are distinct paraneoplastic syndromes with clinical parameters that may alert attentive clinicians to pursue an appropriately aggressive workup [47]. The lethality of pancreatic cancer merits such an approach despite the absence of formal diagnostic guidelines in this area.

NEUROPSYCHIATRIC SYMPTOMS AND PANCREATIC CANCER

Depression is reported to be more common in patients with pancreatic cancer than with other abdominal tumors. In some patients, depression may be the most prominent presenting symptom, possibly secondary to delayed diagnosis. In addition, although patients may not communicate it to their families, they are often aware that a serious illness of some kind is occurring in them [24]. The risk of suicide among male patients with PDAC is almost 11 times higher than the

general male population. Patients who underwent resection are more likely to commit suicide, specifically in the early postoperative period [83].

The association between mood disorders, fatigue, and PDAC has been assumed secondary to the psychosocial impact of diagnosis, loss of independence, and treatment toxicity [2]. However, it is now clear that PDAC has independent detrimental effects on the brain. These symptoms, often present before a diagnosis, are collectively the greatest drivers of declines in health-related quality of life and are independently predictive of survival. Evidence points to neuroinflammatory processes and the need to rethink PDAC as a systemic disease [2].

FAMILY HISTORY

The importance is emphasized of taking a thorough family history when seeing a new patient with pancreatic cancer. A family history of pancreatitis, melanoma, and pancreatic, colorectal, breast and ovarian cancers should be noted [11].

If a cancer syndrome is identified, at-risk relatives should be offered genetic counseling. With or without a known syndrome, individuals with a suspicious family history should be advised on risk-reducing strategies, including smoking cessation and weight loss. The possibility of screening for pancreatic and other cancers should be discussed.

Referral for genetic counseling should be considered for patients diagnosed with pancreatic cancer, especially those with a family history of cancer or who are young, those of Ashkenazi Jewish ancestry, or for whom a hereditary cancer syndrome is suspect. A free pancreatic cancer risk prediction tool, PancPRO, is available and may help determine risk [11].

COMMON PRESENTING SYMPTOMS AND SIGNS

Some, but not all, initial symptoms of PDAC result from a mass effect, such that pancreatic tumor location influences the stage of disease progression when symptoms appear [13].

Abdominal Pain

Abdominal pain is the most common symptom, usually insidious in onset and often present for one to two months at the time of presentation, the pain is often severe, and unrelenting in nature. The typical gnawing, visceral quality of pain is generally epigastric, radiating to the sides and/or straight through to the back; some patients may describe the pain as originating in the back. Nighttime pain is often the predominant complaint. Some patients note increased pain after eating and worsened pain when lying flat [24; 81]. Rarely, acute pain develops when an episode of acute pancreatitis results in tumor occlusion of the main pancreatic duct [84].

While roughly one-third of patients may not have pain at the time of initial presentation, all patients will develop pain at some point [24]. Pancreatic cancer is one of the most painful malignancies, and effective pain control is extremely important [85]. This issue will be discussed in detail later in this course.

Jaundice

The most characteristic sign of tumor in the pancreatic head is obstructive jaundice, for which patients may seek medical attention before their tumor grows large enough to cause abdominal pain (and thus, a somewhat better prognosis). These patients usually notice a darkening of their urine and/or lightening of their stools before they or their families notice the change in skin pigmentation. Jaundice secondary to a tumor in the body or tail of the pancreas typically occurs at a later stage and may be secondary to liver metastases of PDAC [2; 84].

Pruritus can accompany and often precedes obstructive jaundice. If present, it is often the patient's most distressing symptom [24].

Significant Weight Loss

A characteristic feature of pancreatic cancer, significant weight loss may be related to cancer-associated anorexia and/or subclinical malabsorption from pancreatic exocrine insufficiency caused by pancreatic duct obstruction. Nausea and early satiety from gastric outlet obstruction and delayed gastric emptying from the tumor can contribute to weight loss [24]. Significant weight loss is a symptom of cachexia.

Cachexia

Pancreatic cancer cachexia is a multifactorial paraneoplastic syndrome characterized by a loss of skeletal muscle mass, commonly associated with adipose tissue wasting and anorexia, fatigue, and reduced exercise tolerance. Cachexia develops in approximately 80% of patients with PDAC, in whom the syndrome is typically present at diagnosis and responds poorly to therapeutic interventions [47; 86].

Pancreatic cancer leads to the development of cachexia through a combination of distinct factors that explain its high prevalence and clinical importance in this disease [86]:

- Systemic factors, including metabolic changes and pathogenic signals related to PDAC tumor biology
- Factors resulting from the disruption of the digestive and endocrine functions of the pancreas
- Factors related to the close anatomic and functional connection of the pancreas with the gut

Additional Symptoms

The initial assessment can uncover additional diagnostic clues. Undiagnosed diabetes leads to symptoms of glucose intolerance (e.g., polyuria, polydipsia). PDAC can interfere with production of digestive enzymes by the pancreas (pancreatic exocrine insufficiency) and with the ability to break down food and absorb nutrients (malabsorption) in some patients. This malabsorption causes bloating, gas, and a watery, greasy, and/or foul-smelling diarrhea, leading to weight loss and vitamin deficiencies [81].

While long-standing diabetes is a risk factor for later development of PDAC, new-onset hyperglycemia or diabetes has been identified in the majority of patients at diagnosis of otherwise asymptomatic PDAC. Deregulation in glucose homeostasis is often accompanied by changes in subcutaneous adipose tissue. Both represent paraneoplastic syndromes caused by the underlying PDAC [2].

This research is among the most important knowledge advances in PDAC in the past decade. In addition to metabolic deregulation, the pre-diagnostic soft tissue changes and symptoms of cachexia have profound implications for screening, early diagnosis, treatment selection, and patient prognosis [2].

Tumors can also grow locally into the duodenum (proximal for the head of the pancreas, distal for the body and tail of the pancreas) and result in an upper gastroduodenal obstruction [13]. Tumor in the body or tail of the pancreas may cause splenic vein obstruction, resulting in splenomegaly, gastric and esophageal varices, and gastrointestinal hemorrhage [81].

PHYSICAL EXAMINATION

Clinical signs of PDAC during physical examination include jaundice, pruritus, steatorrhea, and vascular issues [2; 24; 82; 84]. Healthcare professionals can usually recognize clinical jaundice when total bilirubin reaches 2.5–3 mg/dL. Patients and their families do not usually notice clinical jaundice until total bilirubin reaches 6–8 mg/dL. Patients with jaundice may have a palpable gallbladder (i.e., Courvoisier sign). As noted, patients with clinical jaundice may have skin excoriations from unrelenting pruritus. If the pancreas has lost the ability to secrete fat-digesting enzymes or if the main pancreatic duct is blocked, steatorrhea will develop.

Migratory thrombophlebitis (i.e., Trousseau syndrome) and venous thrombosis may be present, reflecting the hypercoagulable state that frequently accompanies pancreatic cancer. Thromboembolic events (both venous and arterial) are especially prevalent in advanced disease, and thromboembolic complications occur more commonly with tumors in the pancreatic tail or body.

Multiple arterial emboli resulting from nonbacterial thrombotic endocarditis may be the presenting sign of PDAC. Marantic endocarditis (also known as nonbacterial thrombotic endocarditis) may develop in patients with pancreatic cancer and possibly mimic subacute bacterial endocarditis.

METASTATIC DISEASE

Metastatic disease most commonly affects the liver, peritoneum, lungs, and less frequently, bone [24; 84]. Patients presenting with or developing advanced intra-abdominal disease may have ascites, a palpable abdominal mass, hepatomegaly from liver metastases, or splenomegaly from portal vein obstruction. Subcutaneous metastases (termed Sister Mary Joseph nodules) in the paraumbilical area signify advanced disease; pancreatic

cancer is the origin of a cutaneous metastasis to the umbilicus in 7% to 9% of cases [24; 84]. A metastatic mass in the rectal pouch may be palpable on rectal examination (Blumer shelf). As a metastatic node, left supraclavicular lymphadenopathy may be palpable, while other nodes in the cervical area may also be involved.

LABORATORY TESTING

Routine laboratory tests are often abnormal but nonspecific for PDAC. Common abnormalities include an elevated serum bilirubin and alkaline phosphatase levels, and presence of mild anemia [84].

Patients presenting with jaundice or epigastric pain should be evaluated with complete blood count, blood chemistry panel, and liver function tests to help assess the extent of cholestasis (bilirubin), liver metastasis (alkaline phosphatase), hepatitis (aminotransferases), and nutritional status (albumin, prealbumin). With epigastric pain, serum lipase should be measured to evaluate for acute pancreatitis [2].

DIFFERENTIAL DIAGNOSIS

Differential diagnosis before imaging and biopsy includes acute/chronic pancreatitis, cholangitis, cholecystitis, choledochal cyst, peptic ulcer disease, cholangiocarcinoma, and gastric cancer [85]. Unlike pancreatic exocrine tumors, the symptoms of pancreatic neuroendocrine tumors are distinctly related to excessive secretion of hormones such as insulin, glucagon, gastrin, somatostatin, and vasoactive peptide, resulting in hypoglycemia, hyperglycemia, and GI disturbances such as peptic ulcer and diarrhea.

THE DIAGNOSTIC AND STAGING WORKUP

It is not possible to reliably diagnose a patient with pancreatic cancer based on symptoms and signs alone. Abdominal imaging is used in the diagnostic and staging workup of a patient with suspected PDAC. Additional testing is based on the initial findings, the patient's clinical presentation and risk factors [2].

Accurate PDAC detection and staging at the time of presentation carries substantial implications for appropriate recommendation to patients of the most suitable treatment option, thus maximizing the survival benefit for patients in whom complete resection can be achieved and minimizing the morbidity from unnecessary laparotomy or major surgery in patients with high risk of residual disease following resection. The accuracy critically depends on the appropriate imaging protocol and radiologist experience [2; 87]. As such, decisions about diagnosis, resectability, and management of pancreatic cancer should involve multidisciplinary consultation at high-volume centers [11].

IMAGING

Multidetector Computed Tomography

Multidetector computed tomography (MDCT) angiography with intravenous (IV) contrast is the preferred imaging for initial evaluation of suspected PDAC. The Pancreatic CT Protocol standardizes its use, making MDCT highly accurate for assessing tumor extent, vascular invasion, and distant metastases [11; 16; 88; 89]. The NCCN recommends that MDCT angiography should also cover the chest and pelvis for complete staging [11].



The American Society of Clinical Oncology recommends a multiphase computed tomography (CT) scan of the abdomen and pelvis using a pancreatic protocol or magnetic resonance imaging (MRI) be performed for all patients with pancreatic cancer to assess the anatomic relationships of the primary tumor and to assess for the presence of intra-abdominal metastases.

(<https://ascopubs.org/doi/10.1200/JCO.19.00946>. Last accessed August 19, 2021.)

Strength of Recommendation/Level of Evidence:
Strong/high

MDCT is 77% accurate in predicting resectability and 93% accurate in predicting unresectability [85]. MDCT may be superior to magnetic resonance imaging/magnetic resonance cholangiopancreatography (MRI/MRCP) in vascular enhancement of a PDAC, the most important parameter of resectability. However, MDCT is inferior to MRI/MRCP in depicting isodense tumors or tumors smaller than 1.5 cm in size [54].

Magnetic Resonance Imaging/Magnetic Resonance Cholangiopancreatography (MRI/MRCP)

Abdominal MRI/MRCP with IV contrast also employs a standard multiphase protocol in PDAC, with efficacy comparable to MDCT in preoperative evaluation and assessment of vascular invasion. The sensitivity of MRI/MRCP in detecting liver metastases is nearly 100% (vs. 80% with MDCT) [81; 85].

Selection of initial MDCT or MRI/MRCP is typically based on local availability and expertise [81; 85]. Following initial MDCT, MRI/MRCP is used when PDAC is highly suspected but negative on MDCT, for characterizing small or indeterminate pancreatic and hepatic tumors, and in patients with severe allergy to iodinated IV contrast material used in MDCT [54; 81; 85].

Endoscopic Retrograde Cholangiopancreatography (ERCP)

With endoscopic retrograde cholangiopancreatography (ERCP), contrast dye is injected into the biliary ducts and pan-

creatic duct with an endoscope, and the level of obstruction is delineated. In some case, placement of a biliary stent can help relieve symptoms of jaundice [85]. Patients with obstructive jaundice may have ERCP as the first diagnostic procedure [81].

Ultrasonography

Transabdominal ultrasonography is useful in initial screening of patients who present with possible obstructive jaundice and can rapidly and accurately assess for biliary obstruction. However, definitive diagnosis requires other imaging [24].

Endoscopic ultrasonography is superior to MDCT in detecting solid pancreatic lesions less than 2 cm in size, with accuracy of about 92% [54]. Endoscopic ultrasonography-guided fine-needle aspiration (FNA) also allows for tissue sampling at the time of endoscopic ultrasonography diagnosis [24].

With the restricted field of view, endoscopic ultrasonography is complimentary to MDCT, but it should be used before other imaging options if no pancreatic mass is evident on MDCT. Endoscopic ultrasonography is also valuable in detecting tumor involvement of blood vessels or lymph nodes [11; 89].

Positron-Emission Tomography (PET)

Positron-emission tomography (PET) imaging alone does not offer added advantages to MDCT. Combining PET with CT (PET/CT) is a more recent development that may enhance the detection of occult metastases in pancreatic cancer. The NCCN guidelines consider PET/CT an evolving technology; its role in the diagnosis of PDAC is not yet established [11].

BIOPSY

A positive biopsy is not needed in patients with resectable PDAC before undergoing surgery; biopsy may result in seeding, interfere with definitive surgery, and needlessly delay surgical resection if nondiagnostic [11]. However, histologic confirmation of a pancreatic cancer diagnosis is required in some situations, and endoscopic ultrasonography-guided FNA biopsy is the best modality for obtaining a tissue diagnosis [84].

A pathologic diagnosis is indicated to confirm PDAC in locally advanced or metastatic disease, before neoadjuvant therapy, and in atypical presentations in which differential diagnosis is needed with other pancreatic masses (e.g., pancreatitis, lymphoma, tuberculosis). If a biopsy does not confirm malignancy, it should be repeated at least once [16].

The difficulty of diagnosing PDAC in patients with underlying chronic pancreatitis is noteworthy. In such cases, all typical imaging methods may show abnormalities that do not differentiate between PDAC and chronic pancreatitis, and carbohydrate antigen 19-9 (CA19-9) may be similarly elevated in pancreatitis. These patients may require combined multiple imaging modalities, close follow-up, serial imaging studies, and in some cases, empiric resection to diagnose an underlying pancreatic carcinoma [24].

CARBOHYDRATE ANTIGEN 19-9 (CA19-9)

CA19-9 is a sialylated Lewis A blood group antigen, commonly expressed and shed in benign and malignant pancreatic and biliary disease. Although unsuitable for asymptomatic screening, CA19-9 is the most clinically useful biomarker in PDAC, with good sensitivity (79% to 81%) and specificity (82% to 90%) in symptomatic patients. A normal serum level is 37 U/mL [90].

Preoperative CA19-9 provides important prognostic information. Levels <100 U/mL imply likely resectable disease, while levels >100 U/mL suggest unresectability or metastatic disease. Fewer than 4% of patients with levels >300 U/mL have resectable tumors [24; 90].

In one study, patients with preoperative CA19-9 levels <37 U/mL showed longer median survival (22 to 40 months) than patients with levels >37 U/mL (7 to 30 months). Post-treatment changes (two to five weeks post-resection; six to eight weeks post-chemotherapy) from baseline may predict overall survival [90; 91].

Post-operative CA19-9 levels of <37 U/mL, <200 U/mL, and >500 U/mL were associated with three-year survival rates of 49%, 38%, and 0%, respectively. Post-chemotherapy CA19-9 decreases of $\geq 20\%$ predicted prolonged disease-free survival and overall survival [90; 91].

Limitations

Around 5% to 10% of the population lacks the enzyme necessary to produce CA19-9; monitoring pancreatic cancer with this marker will not be possible in these individuals [24]. Biliary obstruction also stimulates the secretion of CA19-9. Hyperbilirubinemia is associated with elevated CA19-9 and false positivity in patients with obstructive jaundice. Following the treatment of obstruction, re-evaluation of CA19-9 should improve its diagnostic utility [92].

The NCCN recommends measurement of serum CA19-9 levels after neoadjuvant treatment, prior to and immediately following surgery before adjuvant therapy, and in surveillance. The importance is stressed of obtaining CA19-9 immediately before a therapeutic intervention to have an accurate baseline from which to follow response [11].

THE STAGING WORKUP

When a mass lesion of the pancreas is detected on MDCT (with or without additional imaging), it is reasonable to conclude that a neoplasm is present and is most likely malignant PDAC. After a probable diagnosis of pancreatic cancer is made, the next step is the staging evaluation to establish disease extent and resectability. Unlike many other cancers, imaging is the primary means through which the stage of pancreatic cancer is determined [11].

Using initial MDCT (with or without additional imaging), two different systems are involved [11; 93]:

- American Joint Committee on Cancer (AJCC) TNM staging system, to assess tumor status/extent (T), lymph nodes (N), and metastasis (M)
- NCCN guideline to characterize resectable, borderline resectable, or locally advanced disease

TNM Staging

The AJCC system (**Table 6**) is used for staging PDAC in two contexts [16; 94]:

- Clinical staging of all patients with imaging assessment of tumor size and extension, nodal involvement, and distant disease spread
- Pathologic staging of tissue specimens obtained during resection for presence of viable tumor cells

Clinical staging identifies the primary tumor and its vessel involvement, enlarged or suspicious lymph nodes, and metastatic disease sites. TNM staging provides important prognostic information (**Table 7**), but does not assess whether the PDAC tumor is amenable to surgical resection [54; 94].

Resectability Assessment

Complete resection is the only potentially curative treatment for PDAC, but fewer than 20% of patients presenting with PDAC have localized and easily resectable tumors, and non-curative resections provide no survival benefit. Thus, accurate assessment of resectability is crucial [24; 87; 89].

The NCCN guideline classes PDAC resectability into the following clinical stages [11]:

- Stage 1: Resectable
- Stage 2: Borderline resectable (i.e., tumors that are involved with nearby structures so as to be neither clearly resectable nor clearly unresectable with a high chance of removal of all macroscopic disease)
- Stage 3: Locally advanced (i.e., tumors that are involved with nearby structures to an extent that renders them unresectable despite the absence of metastatic disease)
- Stage 4: Metastatic (i.e., non-resectable)


Localized PDAC falls on a spectrum from high to low resectability, determined by the extent of vessel contact and whether the involvement is arterial or venous (**Figure 1**) [11; 54; 84; 87; 89; 95]. Major peripancreatic vessels include the superior mesenteric vein and artery, portal vein, common hepatic artery, and celiac artery. Tumor contact can be characterized as encasement (≥ 180 degrees of the vessel circumference), abutment (<180 degrees of the circumference), or direct involvement (absence of fat plane between tumor and vessel).

AMERICAN JOINT COMMISSION ON CANCER EXOCRINE PANCREATIC CANCER TNM STAGING	
Category	Criteria
Primary tumor (T)	
TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
Tis	Carcinoma in situ, including high-grade PanIN (PanIN-3) and IPMN, ITPN, or MCN with high-grade dysplasia
T1	Tumor ≤ 2 cm in greatest dimension
T1a	Tumor ≤ 0.5 cm in greatest dimension
T1b	Tumor > 0.5 and < 1 cm in greatest dimension
T1c	Tumor 1–2 cm in greatest dimension
T2	Tumor > 2 and ≤ 4 cm in greatest dimension
T3	Tumor > 4 cm in greatest dimension
T4	Tumor involves the celiac axis, superior mesenteric artery, and/or common hepatic artery, regardless of size
Regional lymph nodes (N)	
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in one to three regional lymph nodes
N2	Metastasis in four or more regional lymph nodes
Distant metastasis (M)	
M0	No distant metastasis
M1	Distant metastasis
IPMN = intraductal papillary mucinous neoplasm; ITPN = intraductal tubulopapillary neoplasm; MCN = mucinous cystic neoplasm; PanIN = pancreatic intraepithelial neoplasia	
Source: [93]	Table 6

AMERICAN JOINT COMMISSION ON CANCER ANATOMIC STAGE/ PROGNOSTIC GROUPS FOR EXOCRINE PANCREATIC CANCER			
Stage	T	N	M
A	T1	N0	M0
IB	T2	N0	M0
IIA	T3	N0	M0
IIB	T1–T3	N1	M0
III	Any T	N2	M0
	T4	Any N	M0
IV	Any T	Any N	M1
Source: [93]	Table 7		

In the past, vascular infiltration by PDAC was considered unresectable, but surgical advances have increased the number of patients with initial borderline resectable or locally advanced disease who can undergo resection. In general, venous abut-

ment or encasement is usually borderline resectable as long as the venous segment is reconstructable. Arterial reconstruction is substantially more difficult and risky than venous reconstruction with comparable tumor contact.



Based on PDAC clinical status of resectable, borderline resectable, locally advanced, or metastatic disease, additional considerations and therapeutic approaches will be undertaken. The time-urgency between the first availability of full imaging findings, multidisciplinary evaluation, the diagnostic and staging workup, discussion with the patient of available treatment options, and treatment initiation cannot be overstated in this aggressive malignancy.

TREATMENT APPROACHES FOR PANCREATIC CANCER

As mentioned, the initial imaging workup of PDAC confirms the diagnosis, searches for evidence of metastases, and classifies nonmetastatic PDAC into resectable, borderline resectable, or locally advanced disease based on the involvement of surrounding arterial (superior mesenteric artery, common hepatic artery, and celiac axis) and venous (superior mesenteric vein or portal vein) structures, and other nearby organs and lymph nodes [96].

On average, 10% to 20% of patients initially present with “up-front” resectable PDAC. However, an increasing number of patients with initial borderline resectable or locally advanced disease are eligible for surgical resection as a result of neoadjuvant (i.e., before resection) therapies which may downstage the tumor, and advances in surgical technique, such as venous reconstruction in a vascular infiltration formerly considered unresectable [2].

In all therapeutic decisions, multidisciplinary collaboration to formulate treatment planning and disease management that incorporates patient preferences and available support, their comorbidity profile, symptom burden, and performance status should be the standard of care [6; 7; 10].

PATIENT FUNCTIONAL STATUS

Performance status is an important indicator of general well-being and the ability to perform activities of daily living in

patients with cancer and is frequently assessed in both clinical and research settings. Performance status is repeatedly shown to predict important clinical outcomes, including quality of life, chemotherapy toxicity, response to chemotherapy, terminal illness, progression-free survival, and overall survival in patients with cancer [97].



According to the ASCO, the baseline performance status, symptom burden, and comorbidity profile of a person diagnosed with potentially curable pancreatic cancer should be carefully evaluated.

(<https://ascopubs.org/doi/10.1200/JCO.19.00946>. Last accessed August 19, 2021.)

Strength of Recommendation/Level of Evidence:
Strong/high

The Karnofsky Performance Status tool has been used for this purpose, but PDAC guidelines and randomized controlled trials now solely employ the Eastern Cooperative Oncology Group Performance Status (ECOG) scale (**Table 8**) [97]. For instance, some chemotherapies are indicated solely for patients with good ECOG performance status (0 or 1).

Baseline functional status and comorbidity profile should be carefully evaluated, because both have major implications for a person’s ability to tolerate therapy. Performance status is consistently identified as a prognostic factor for people with pancreatic cancer. It is also an important determinant in treatment selection; some patients with up-front resectable PDAC may be physically weakened by weight loss and cachexia to an extent that places them at high risk of serious complications or mortality from definitive surgery. Performance status also helps predict chemotherapy toxicity, which can determine the treatment approach for patients with performance status 0 to 1 (multi-agent regimens) or performance status ≥ 2 (e.g., single-agent gemcitabine) [8].

EASTERN COOPERATIVE ONCOLOGY GROUP (ECOG) PERFORMANCE STATUS SCALE	
Score	Definition
0	Fully active No performance restrictions
1	Strenuous physical activity restricted Fully ambulatory and able to carry out light work
2	Capable of all self-care but unable to carry out any work activities Up and about >50% of waking hours
3	Capable of only limited self-care Confined to bed or chair >50% of waking hours
4	Completely disabled Cannot carry out any self-care Totally confined to bed or chair
5	Deceased
<i>Source: [98]</i>	

Table 8

Similarly, the comorbidity profile can influence the choice of chemotherapy, such as avoiding fluoropyrimidine-based regimens in patients with a known history of uncontrolled coronary artery disease. Nonetheless, performance status and comorbidities alone should not be used simply to rule in or out patients for treatment. For instance, disease control of comorbidities, such as controlled type 2 diabetes, can indicate that patient benefit from treatment may outweigh risks associated with poorly controlled comorbid diabetes [8].

RECOMMENDED TREATMENT OPTIONS BY CLINICAL STAGE

Treatment approaches for PDAC include surgical resection, chemotherapy, radiation therapy, and combined regimens (chemoradiation therapy). Chemotherapy is the backbone of pancreatic cancer treatment; most patients present with disease too advanced to benefit from surgery or resection alone may be insufficient to provide a substantive survival advantage over best supportive care. Chemotherapy and radiation therapy also have a role in palliation, as will be discussed in a later section [99].

Curative surgical approaches for resectable pancreatic cancer are well-established. In contrast, the pace of new U.S. Food and Drug Administration (FDA) approvals and/or phase III evidence continue to make chemotherapy, molecular-targeted therapy, radiation, and chemoradiotherapy approaches a fluid, evolving area, requiring frequent updating and revisions in multidisciplinary clinical practice guidelines for pancreatic cancer treatment. Many potential treatment approaches lacking phase III or prospective evidence are being addressed, with publication of trial results awaited [2].

Resectable or Borderline Resectable PDAC

For patients with resectable or borderline resectable PDAC, neoadjuvant therapy consists of chemotherapy with or without radiation therapy before radical pancreatic resection [99]. Radical pancreatic resection may include Whipple procedure (pancreaticoduodenal resection) or total pancreatectomy when necessary for adequate margins. Distal pancreatectomy is indicated for tumors of the body and tail of the pancreas.

Following resection, patients may receive postoperative chemotherapy or postoperative chemoradiation therapy (typically fluorouracil [5-FU] chemotherapy and radiation therapy) [99].

Locally Advanced PDAC

Chemotherapy with or without targeted therapy is recommended for patients with locally advanced PDAC [99]. For patients without metastatic disease, this should be followed by chemoradiation therapy. If removal is a possibility, radical pancreatic resection may be attempted. Palliative surgery options include surgical biliary and/or gastric bypass, percutaneous radiologic biliary stent placement, or endoscopic biliary stent placement.

Metastatic or Recurrent PDAC

Treatment of metastatic or recurrent PDAC is limited to chemotherapy with or without targeted therapy [99]. Palliative approaches should be used whenever available and feasible to improve patient comfort and quality of life.

RESECTION OF PANCREATIC CANCER

Selecting patients for surgery should be based on the probability of cure as determined by resection margins. Other factors include comorbidities, overall performance status, and age. Pancreaticoduodenectomy and distal and total pancreatectomy are curative resection options based on the location, size, and locally invasive aspects of the tumor. Each has its own set of perioperative complications and risks, which should be considered by the surgical team and discussed with the patient [24].

Mortality rates from resection have fallen significantly, but morbidity remains common and interferes the delivery of adjuvant therapy in up to 40% of patients. The NCCN recommends that patients seek out high-volume centers performing more than 15 to 20 resections annually, with multidisciplinary expertise to optimize their treatment plan and increase opportunities for clinical trial participation [2].

The only curative treatment for PDAC is radical surgery, but potential cure is only possible with a microscopically negative resection margin (R0). Macroscopic (R2) and microscopic (R1) margin infiltration have survival trends similar to patients without surgery. R0 is a minimum >1 mm distance of viable tumor cells from the resection margin, R1 is ≤1 mm distance. A retrospective analysis of 44,852 patients with PDAC reported median survival of 19.7 months following R0, 14.3 months following R1, and 9.8 months with R2 resections compared with 10.3 months without surgery [100]. An incomplete tumor

resection imposes morbidity risks without benefit to the patient, and the aim of resection is to obtain microscopically negative margins (R0) [101].

Tissue specimens obtained during resection are examined. During resection, lymphadenectomy is performed, including at least 15 lymph nodes, which are likewise examined as part of pathologic staging [16].

With surgical advances and greater use of adjuvant therapies, long-term cancer survival outcomes following resection were anticipated to improve over time [102]. However, in 1,147 pancreatic resections performed over three decades at the Memorial Sloan Kettering Cancer Center, a lack of progress in long-term survival was reported. Although patients treated between 2000 and 2009 had lower rates of operative mortality and greater one-year survival, for patients treated in the 1980s, 1990s, and 2000s, the median survival was 23.2, 25.6, and 24.5 months, respectively [103]. The five-year survival rates were 17%, 20%, and 8%, respectively. These data underscore the need for earlier detection and more effective systemic therapies [102].

Approaches

Pancreaticoduodenectomy (Whipple Procedure)

Used for tumors in the pancreatic head or periampullary region, the conventional Whipple procedure involves removal of the pancreatic head, duodenum, gallbladder, and the antrum of the stomach, with surgical drainage of the distal pancreatic duct and biliary system, usually through anastomosis to the jejunum. The primary reason for removing so much of the intra-abdominal structures is that they all share a common blood supply [24; 102].

The former high morbidity and mortality rates of Whipple have declined with the greater experience of a more limited number of surgeons who regularly perform the procedure in high-volume centers [102]. Common morbidities include delayed gastric emptying in roughly 25% of patients, which may require nasogastric decompression and a longer hospital stay. Pancreatic anastomotic leak can be treated with adequate drainage. Postoperative abscesses are not uncommon [24].

With operative mortality associated with Whipple decreasing from around 25% in the 1970s to less than 2% at high-volume centers in the 2010s, the focus has shifted from surviving the operation to surviving the cancer [104].

Distal Pancreatectomy

Distal pancreatectomy is a procedure for tumors in the pancreatic body or tail. It has a lower mortality than standard Whipple, but its use in curative resection is limited; with tumors in this location seldom causing bile duct obstruction, most patients present at a later stage with unresectable disease. The procedure involves resection of the distal pancreas containing the tumor with splenectomy and over-sewing of the

distal pancreatic duct. Complications involve pancreatic stump leak, hemorrhage, and endocrine insufficiency. Laparoscopic exploration should precede attempted resection, because occult peritoneal metastases are common [16; 24].

Total Pancreatectomy

Total pancreatectomy, the least commonly performed procedure with the highest associated mortality (8.3%), may be needed to achieve an R0 resection margin for tumors in the neck of the pancreas, especially with extension into the body or tail, and in multifocal PDAC. Total pancreatectomy may be an option to pancreatic anastomosis in highly selected patients with a high-risk pancreas (small pancreatic duct) and obese patients with pancreatic fat infiltration. The metabolic consequences of permanent exocrine insufficiency and diabetes have a detrimental impact on quality of life and long-term survival [16; 24; 102].

Vascular Resection

Vascular involvement has traditionally been a formal contraindication to resection. With recent advances, venous resection and reconstruction can achieve R0 resection with similar overall survival and morbidity compared to surgery without venous resection. However, arterial resection during Whipple is associated with increased mortality and morbidity (bowel ischemia, hemorrhage, thrombosis) and is generally not recommended [16].

Progress in neoadjuvant therapies may downstage some tumors with arterial invasion to borderline resectable or resectable disease, making resection more achievable. Despite these advancements, it is currently accepted that arterial reconstruction is only appropriate in highly selected patients in high-volume centers with surgeons who are familiar with the advanced techniques required for reconstruction [16].

Total pancreatectomy should be considered in patients with locally advanced tumors who undergo pancreatectomy with arterial resection and reconstruction [16].

Biliary Drainage

In most patients with jaundice, early resection without biliary drainage is preferred. Preoperative drainage is indicated in patients with cholangitis or with obstructive jaundice scheduled for neoadjuvant therapy. Endoscopic retrograde placement of a fully covered metal stent is preferred. Endoscopic ultrasonography-guided stent placement is an effective and safe alternative [16].

CHEMOTHERAPIES IN PANCREATIC CANCER

As mentioned, the backbone of PDAC treatment is chemotherapy. Most patients present with advanced disease, and even those who undergo resection will require adjuvant chemotherapy. Chemotherapy is also used as neoadjuvant therapy and in metastatic disease with first-line or second-line indications [11].

Until recently, chemotherapies found effective in other GI cancers were applied to patients with advanced PDAC; the few agents showing any response became adjuvant therapies in localized PDAC. The near-futility in effective chemotherapy and redundancy in agents used in localized and metastatic PDAC reflects the pathologic complexity of this cancer and its profound resistance to cytotoxic therapies [2].

Since 2010, chemotherapy effectiveness has improved with the introduction of combination regimens, the identification of patients in whom mutational status conferred improved response to existing chemotherapies, and the introduction of novel compounds explicitly targeting mutational-related advanced PDAC.

FDA-Approved Chemotherapies in PDAC

In addition to single chemotherapy agents, the FDA has approved regimens of these agents, including FOLFIRINOX (consisting of folinic acid [also referred to as leucovorin], fluorouracil [5-FU], irinotecan [IRN], and oxaliplatin [OX]) (Table 9) [3; 24; 80; 99]. Available chemotherapies are associated with acute and delayed toxicities, some of which can be dose-limiting (Table 10). Table 11 summarizes the 2021 NCCN guideline for chemotherapy and chemoradiotherapy in PDAC.



According to the American Society of Clinical Oncology, all patients with resected pancreatic adenocarcinoma who did not receive preoperative therapy should be offered six months of adjuvant chemotherapy in the absence of medical or surgical contraindications. The mFOLFIRINOX regimen is preferred in the absence of concerns for toxicity or tolerance.

(<https://ascopubs.org/doi/10.1200/JCO.19.00946>. Last accessed August 19, 2021.)

Strength of Recommendation/Level of Evidence:
Strong/high

Fluoropyrimidines

Fluorouracil is a fluorinated (fluoro)-pyrimidine antimetabolite that inhibits thymidylate synthase and interferes with RNA synthesis and function, with some effect on DNA.

Capecitabine is an oral fluoropyrimidine that undergoes hepatic hydrolysis to form fluorouracil. The final enzyme, thymidine phosphorylase, is present at higher levels in tumor tissue, providing better selectivity and tolerability.

Gemcitabine is a pyrimidine antimetabolite that inhibits DNA polymerase and ribonucleotide reductase, which in turn inhibit DNA synthesis, blocks DNA replication and several forms of DNA repair [3; 24; 80; 99].

Erlotinib

Erlotinib is a human epidermal growth factor receptor type 1/epidermal growth factor receptor (HER1/EGFR) tyrosine kinase inhibitor. EGFR is expressed on the cell surface of normal cells and cancer cells. Erlotinib inhibits intracellular phosphorylation, which prevents further downstream signaling, resulting in cell death [3; 24; 80; 99].

Paclitaxel

Paclitaxel protein bound is a microtubular inhibitor (albumin-conjugated formulation) and a natural taxane that prevents depolymerization of cellular microtubules, which results in DNA, RNA, and protein synthesis inhibition [3; 24; 80; 99].

Irinotecan Liposomal

Irinotecan and its active metabolite SN-38 bind reversibly to the topoisomerase-1 DNA complex and prevent re-ligation of the single-strand breaks, leading to exposure time-dependent double-strand DNA damage and cell death. Irinotecan liposomal is used in combination with fluorouracil and leucovorin [3; 24; 80; 99].

DNA Damage Repair Mutational Status and Targeted Therapies

Platinum agents (e.g., cisplatin, oxaliplatin) and olaparib are recommended in patients with mutation in DNA damage repair (DDR) genes by the NCCN. DDR mutations are present in up to 24% of PDACs, most commonly *BRCA1/2* and *PALB2*. Germline *BRCA1/2* mutations (gBRCAm) affect approximately 7% of patients with PDAC [108]. DDR genes encode for proteins in the homologous repair pathway and DNA double-stranded break repair; thus, mutations may be more sensitive to further DNA damage [99].

Cisplatin inhibits DNA synthesis by the formation of DNA cross-links; denatures the double helix; covalently binds to DNA bases; and disrupts DNA function. Oxaliplatin is an alkylating agent. Following intracellular hydrolysis, the compound binds to DNA, forming cross-links that inhibit DNA replication and transcription, resulting in cell death [24; 99].

PDACs with DDR mutations demonstrate improved responses to platinum-based therapies, and patients with advanced PDAC showed significantly improved median overall survival (22 months vs. 9 months) compared with nonplatinum therapy [96].

Poly (ADP-ribose) polymerase (PARP) inhibition has been posited to act synergistically with *BRCA1/2* mutations by inhibiting single-stranded break repair, causing an accumulation of DNA damage and tumor-cell death [99; 109]. Olaparib is a PARP inhibitor FDA-approved for PDAC with gBRCAm as maintenance therapy to sustain a progression-free state during platinum-based chemotherapy in metastatic PDAC [96].

CHEMOTHERAPY PROTOCOLS IN PANCREATIC CANCER			
Drug	Dose and route	Administration	Given on days
Gemcitabine Indication: Nonmetastatic PDAC Cycle length: 4 weeks (once weekly for 3 weeks, then 1 week off)			
Gemcitabine	1,000 mg/m ² IV	Dilute in 250 mL NS (concentration ≤40 mg/mL), administered over 30 minutes.	Days 1, 8, and 15
Gemcitabine and capecitabine (GemCap) Indication: Adjuvant therapy Cycle length: 28 days Duration: 6 months			
Gemcitabine	1,000 mg/m ² IV	Dilute in 250 mL NS (concentration ≤40 mg/mL), administered over 30 minutes.	Days 1, 8, and 15
Capecitabine ^a	830 mg/m ² per dose oral	Twice daily (total 1,660 mg/m ² per day), 12 hours apart. Swallow with water within 30 minutes post-meal.	Days 1 through 21
Modified FOLFIRINOX Cycle length: 14 days			
Oxaliplatin ^b	85 mg/m ² IV	Dilute in 500 mL D5W, administer over 2 hours (before leucovorin). Shorter schedules (e.g., 1 mg/m ² per minute) appear safe.	Day 1
Leucovorin	400 mg/m ² IV	Dilute in 250 mL normal saline or D5W, administer over 2 hours (after oxaliplatin).	Day 1
Irinotecan ^c	150 mg/m ² IV	Dilute in 500 mL normal saline or D5W, administer over 90 minutes concurrent with the last 90 mins of leucovorin infusion, in separate bags, using Y-line connection.	Day 1
Fluorouracil	2,400 mg/m ² IV	Dilute in 500–1,000 mL 0.9% normal saline or D5W, administered as continuous IV infusion over 46 hours. ^d	Day 1
FOLFIRINOX Indication: Metastatic PDAC Cycle length: 14 days			
Oxaliplatin ^b	85 mg/m ² IV	Dilute in 500 mL D5W, administer over 2 hours (before leucovorin). Shorter schedules (e.g., 1 mg/m ² per minute) appear safe.	Day 1
Leucovorin	400 mg/m ² IV	Dilute in 250 mL normal saline or D5W, administer over 2 hours (after oxaliplatin).	Day 1
Irinotecan ^c	150 mg/m ² IV	Dilute in 500 mL normal saline or D5W, administer over 90 minutes concurrent with the last 90 mins of leucovorin infusion, in separate bags, using Y-line connection.	Day 1
Fluorouracil	400 mg/m ² IV bolus	Give undiluted (50 mg/mL) as a slow IV push over 5 minutes (immediately after leucovorin).	Day 1
Fluorouracil	2400 mg/m ² IV	Dilute in 500–1,000 mL 0.9% normal saline or D5W, administer as continuous IV infusion over 46 hours (immediately after IV bolus). ^d	Day 1
^a Capecitabine is contraindicated in patients with known DPD deficiency. ^b Many centers routinely infuse oxaliplatin via central venous line because of local pain with infusion into a peripheral vein ^c Consider a lower dose of irinotecan with poor performance status. ^d To accommodate an ambulatory pump for outpatients, can be administered undiluted (50 mg/mL) or the total dose diluted in 100–150 mL normal saline.			
Source: [98; 105]			Table 9

ACUTE AND DELAYED CHEMOTHERAPY TOXICITIES ^a		
Agent	Acute Toxicities	Delayed Toxicities
Fluorouracil	Nausea and vomiting Diarrhea	Oral and GI ulcers Bone marrow depression Diarrhea (especially with leucovorin) Neurologic defects, usually cerebellar Cardiac arrhythmias Palmar-plantar erythrodysesthesia (hand-foot syndrome)
Capecitabine	Nausea and vomiting	Hand-foot syndrome Diarrhea Stomatitis Dermatitis Bone marrow depression Hyperbilirubinemia
Gemcitabine	Fatigue Nausea and vomiting Fever	Bone marrow depression Edema Pulmonary toxicity
Irinotecan	Diarrhea	Diarrhea Leukopenia
Oxaliplatin	Peripheral sensory neuropathy Pharyngolaryngeal dysesthesias Paresthesias	Bone marrow depression Diarrhea Persistent neuropathy
Paclitaxel	Hypersensitivity reactions	Bone marrow depression Peripheral neuropathy Alopecia Arthralgias
^a Dose-limiting toxicities are bold-faced.		
Source: [106; 107]		Table 10

The NCCN expands the use of olaparib to PDAC with gPALB2m. There are calls to expand these agents to PDACs with somatic DDR mutations [108].

Other FDA-Approved Targeted Therapies

The approved indications for the following agents are biomarker-defined, rather than by tumor site (e.g., pancreatic).

Pembrolizumab

Pembrolizumab is indicated in patients with microsatellite-instability-high (MSI-H) or dMMR mutations. Immune checkpoint inhibitors (ICIs) have efficacy in solid tumors with a high tumor mutational burden, and MSI-H or dMMR mutation solid tumors are associated with high tumor mutational burden. The ICI pembrolizumab is an anti-programmed death receptor-1 antibody that releases inhibition of the immune response, improving antitumor immunity [11; 96].

Pembrolizumab is approved for any solid tumor with MSI-H or dMMR mutation that progresses during treatment without any satisfactory alternative treatment options [11; 96]. This

agent represented the first FDA approval (in 2017) with a biomarker-defined indication (i.e., agnostic of cancer site) [107]. Although this mutation is present in only about 1% of PDAC tumors, up to 83% of patients with dMMR PDAC respond to pembrolizumab [110].

Larotrectinib and Entrectinib

Larotrectinib and Entrectinib are neurotrophin receptor kinase (NTRK) inhibitors approved (in 2018 and 2019) for advanced, morbid, or unresectable solid tumors with NTRK fusion mutations, found in less than 1% of PDACs [96].

The mutation product, TRK fusion protein, activates mitogen activated protein kinase-extracellular regulated kinase and phosphoinositide3 kinase-serine threonine signaling pathways, implicated in the oncogenesis of pancreatic cancer [96]. The NCCN recommends larotrectinib and entrectinib as first-line and subsequent treatment options for patients with NTRK gene fusion-positive locally advanced or metastatic PDAC [11].

NCCN TREATMENT SUMMARY FOR PDAC		
Strength of Recommendation/ Evidence	Regimen	Notes ^a
Adjuvant stage 1 (resectable)		
Category 1	Gemcitabine Gemcitabine/capecitabine 5-FU/leucovorin	-
Category 2a	5-FU continuous infusion Chemoradiation	Chemoradiation should follow induction chemotherapy, with or without subsequent chemotherapy
Category 2B	Capecitabine	-
Neoadjuvant stage 1/2 (resectable or borderline resectable)		
Category 2A	Gemcitabine/paclitaxel NAB	-
Category 2B	Gemcitabine/cisplatin ^b FOLFIRINOX Chemoradiation	-
Stage 3 (locally advanced)		
Category 1	Gemcitabine	Preferred for patients with poor ECOG PS (≥ 2)
Category 2A	Gemcitabine/paclitaxel NAB Gemcitabine/erlotinib Gemcitabine/cisplatin ^b Gemcitabine/capecitabine Gemcitabine fixed-dose rate FOLFIRINOX Chemoradiation	Fixed-dose rate gemcitabine is a category 2B recommendation for patients with poor ECOG PS (≥ 2) Chemoradiation should follow induction chemotherapy, with or without subsequent chemotherapy
Category 2B	Gemcitabine/docetaxel/capecitabine Capecitabine 5-FU continuous infusion FOLFOX	-
Stage 4 (metastatic)		
Category 1	Gemcitabine Gemcitabine/paclitaxel NAB (preferred) Gemcitabine/erlotinib FOLFIRINOX (preferred)	-
Category 2A	Gemcitabine/cisplatin ^b Gemcitabine/capecitabine Gemcitabine fixed-dose rate Olaparib Pembrolizumab (for MSI-H or dMMR tumors only) Larotrectinib (for <i>NTRK</i> -positive only)	Fixed-dose rate gemcitabine is a category 2B recommendation for patients with poor ECOG PS (≥ 2) Olaparib for maintenance therapy only in <i>BRCA1/2</i> or <i>PALB2</i> mutated stage 4 disease without progression after 4 to 6 months of first-line platinum-based therapy
Category 2B	Gemcitabine/docetaxel/capecitabine Capecitabine ^c 5-FU continuous infusion ^c FOLFOX Entrectinib (for <i>NTRK</i> -positive only)	-

Table 11 continues to next page.

NCCN TREATMENT SUMMARY FOR PDAC (Continued)		
Strength of Recommendation/ Evidence	Regimen	Notes ^a
Second-line therapy		
Category 1	Gemcitabine ^{c,d} 5-FU/leucovorin/irinotecan ^d	-
Category 2A	Gemcitabine fixed-dose rate	Fixed-dose rate gemcitabine is a category 2B recommendation for patients with poor ECOG PS (≥ 2)
Category 2B	Capecitabine ^{c,e} 5-FU continuous infusion ^{c,e}	-
Strength of Recommendation Definitions		
Category	Definition	
1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.	
2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.	
2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.	
^a ECOG performance status (PS) 0/1 only, unless noted. ^b In <i>BRCA1/2</i> or <i>PALB2</i> mutations only. ^c Poor ECOG PS (≥ 2) only. ^d If prior non-gemcitabine-based therapy. ^e If prior gemcitabine-based therapy.		
Source: [11]		

Table 11

Chemotherapy Efficacy: Localized Disease

A variety of data on chemotherapy efficacy are available, allowing for comparison of available agents in specific patient populations (**Table 12**). However, the terminology used can be confusing. Disease-free survival and progression-free survival are synonymous terms, and choice of the term used in this section will reflect the reference material. This is also the case with median survival and median overall survival. Unless noted otherwise, all patient outcomes are reported as median figures.

The CONKO-001 trial established gemcitabine as standard adjuvant chemotherapy. In this study, 354 patients were randomized to receive gemcitabine or observation after resection and followed a median 136 months. Gemcitabine led to a 24% improvement in overall survival, a 10.3% absolute improvement in 5-year survival (20.7% vs. 10.4%), and a 4.5% improvement in 10-year survival (12.2% vs. 7.7%), compared to observation [111; 112].

The ESPAC-3 trial showed the importance of completing the full post-resection adjuvant chemotherapy course (six cycles) in extending median overall survival of these patients compared with those not completing chemotherapy (28.0 months vs. 14.6 months) [96].

ADJUVANT CHEMOTHERAPY TRIALS IN RESECTABLE PDAC		
Phase III trial (Year)	Chemotherapy Comparison	Median Survival (months)
ESPAC-1 (2004)	5-FU vs. observation	21 vs. 15.5
CONKO-001 (2013)	Gemcitabine vs. observation	22.8 vs. 20.2
ESPAC-3 (2012)	Gemcitabine vs. 5-FU/leucovorin	46 vs. 39
ESPAC-4 (2017)	Gemcitabine/ capecitabine vs. gemcitabine alone	28 vs. 25.5
PRODIGE 24 (2018)	Modified FOLFIRINOX vs. gemcitabine	54.4 vs. 35
APACT (2019)	Gemcitabine/ paclitaxel vs. gemcitabine alone	40.5 vs. 36.2
5-FU = 5-fluorouracil.		
Source: [2]		

Table 12

FIRST-LINE CHEMOTHERAPY TRIALS IN METASTATIC PDAC

Phase III Trial (Year)	Chemotherapy Comparison	Median Survival (Months)
Cullinan (1985)	5-FU vs. 5-FU/doxorubicin vs. 5-FU/doxorubicin/mitomycin	5.5 vs. 5.5 vs. 4.5
Burris (1997)	5-FU vs. gemcitabine	4.4 vs. 5.6
Tempero (2003)	Gemcitabine vs. gemcitabine fixed dose rate	5 vs. 8
Heinemann (2006)	Gemcitabine ± cisplatin	6.0 vs. 7.5
NCIC-CTG PA.3 (2007)	Gemcitabine ± erlotinib	5.9 vs. 6.2
Cunningham (2009)	Gemcitabine ± capecitabine	6.2 vs. 7.1
CALGB 80303 (2010)	Gemcitabine ± bevacizumab	5.9 vs. 5.8
SWOG S0205 (2010)	Gemcitabine ± cetuximab	5.9 vs. 6.3
PRODIGE 4 (2011)	Gemcitabine vs. FOLFIRINOX	6.8 vs. 11.1
MPACT (2013)	Gemcitabine ± nab-paclitaxel	6.7 vs. 8.5

Source: [2]

Table 13

A continuation, ESPAC-4, found adding another fluoropyrimidine-based agent (capecitabine) to gemcitabine was superior to gemcitabine alone in median survival (28.0 months vs. 25.5 months) and five-year survival (28.8% vs 16.3%). A synergistic effect between gemcitabine and capecitabine on the DNA thymidylate enzyme was suggested [96].

PRODIGE-24 randomized 493 patients (ECOG performance status ≤ 1) with resected PDAC to modified FOLFIRINOX or gemcitabine for 24 weeks. At median 33.6 month follow-up, the disease-free survival with modified FOLFIRINOX was 21.6 months, compared with 12.8 months with gemcitabine [113]. Grade 3/4 toxicities were more frequent with mFOLFIRINOX (75.9%) than gemcitabine (52.9%). Nonetheless, the median 54.4-month overall survival with resection followed by mFOLFIRINOX is the longest survival reported to date with phase III results [5; 114].

Tolerance of adjuvant therapy remains a limitation, and patients commonly receive less than 50% of the planned dose, reflecting exposure to significant chemotherapy-related toxicity in patients experiencing substantial post-resection morbidity [2].

Chemotherapy Efficacy: Advanced/Metastatic Disease

First-Line Chemotherapy in Metastatic PDAC

5-FU has been used in pancreatic cancer treatment since the 1950s. Patients with advanced PDAC typically show response rates greater than 20% and median survival of 2.5 to 6 months [24; 80].

In 1997, gemcitabine replaced 5-FU as first-line treatment in metastatic PDAC by improving one-year survival rates (18% vs. 2%) and median overall survival (5.65 months vs. 4.41 months) [32]. Subsequently, numerous attempts to improve gemcitabine efficacy in metastatic PDAC have involved add-

ing another cytotoxic drug [2; 96]. Some show marginal but statistically significant improvements in median survival over gemcitabine alone (**Table 13**).

The NCIC CTG PA.3 trial found a nonmeaningful clinical improvement with gemcitabine/erlotinib over gemcitabine alone in median overall survival (6.24 months vs. 5.91 months). Despite FDA approval for locally advanced/metastatic PDAC, the clinical impact of this modest gain with increased toxicity can be questioned [32; 96].

PRODIGE 4/ACCORD 11 demonstrated that patients with advanced PDAC and ECOG performance status ≤ 1 had better outcomes with FOLFIRINOX than gemcitabine in median overall survival (11.1 months vs. 6.8 months) and progression-free survival (6.4 months vs. 3.3 months). Following these findings, FOLFIRINOX became standard first-line therapy for candidate patients [2].

FOLFIRINOX was associated with more toxicities, but the six-month degradation in quality of life was better in FOLFIRINOX than gemcitabine (31% vs. 66%). Improved cancer control with FOLFIRINOX may be due to the inclusion of irinotecan, which has activity against PDAC and synergistic activity when given prior to 5-FU [96].

Finally, the MPACT study demonstrated an improvement of 1.8 months in both median overall survival and median progression-free survival with gemcitabine plus nab-paclitaxel versus gemcitabine alone, leading to another first-line option for metastatic PDAC [96].

Second-Line Chemotherapy in Metastatic PDAC

Second-line therapy primarily consists of doublet therapy using the alternative pyrimidine backbone to what was used in the first-line setting. In 2016, the NAPOLI-1 trial demonstrated that after progression on a first-line gemcitabine-containing

regimen for metastatic PDAC, 5-FU/leucovorin plus nanoliposomal irinotecan improved overall survival from 4.2 months (with 5-FU/leucovorin alone) to 6.1 months. As with nab-paclitaxel, improving the delivery of traditional chemotherapies may lead to more effective treatments for individuals with pancreatic cancer [32].

The POLO trial examined targeted maintenance therapy in a biomarker-selected population. In patients with metastatic PDAC harboring germline *BRCA1/2* mutations who had not progressed on first-line platinum-based chemotherapy, those randomized to olaparib had improved median progression-free survival (7.4 months compared with 3.8 months with placebo), but olaparib did not improve median overall survival [109]. The median duration of response to olaparib was 6 months, but was more than 24 months in a subset of patients (23%), which is exceptional in metastatic PDAC [108].

In second-line chemotherapy after progression on a first-line regimen, there is considerable heterogeneity in the survival of patients, and predicting which patients will benefit is not established. The decision to pursue second-line chemotherapy should be individualized and based on the patient's goals and preferences. Factors influencing the choice of second-line therapy include the regimen used for first-line therapy, performance status and comorbidity, and mutation status [106].

RADIATION THERAPY FOR PANCREATIC CANCER

In addition to resection and chemotherapy, treatment of patients with PDAC may include radiation therapy or chemoradiotherapy. Unlike chemotherapy, the role of radiation therapy in the treatment of PDAC is uncertain. Radiation therapy is not a stand-alone treatment in local PDAC but is sequenced with chemotherapy as chemoradiotherapy.

Earlier adjuvant radiation therapy trials demonstrated an overall survival and disease-free survival benefit, but subsequent European chemoradiation studies showed negative findings [12]. Technical advances suggest increasing promise with radiation therapy, but multi-institutional randomized trials in PDAC have lagged [12].

Stereotactic body radiation therapy has promising local control and quality of life, and is being evaluated for locally advanced and borderline resectable PDAC. However, adjuvant stereotactic body radiation therapy remains investigational with high toxicity risk and is only recommended as part of a clinical trial [12].

In the absence of phase 3 trials directly comparing neoadjuvant treatment approaches with or without radiation, adjuvant and neoadjuvant chemoradiation in PDAC awaits definitive evidence. Several such trials are in progress [2; 12]. In particular, RTOG 0848 is expected to definitively clarify the role of post-resection radiotherapy [115].

Nonetheless, the prospective cohort and retrospective evidence suggestive of decreased local recurrence and disease progression is sufficient for ASTRO, the NCCN and ASCO to recommend radiation therapy. Standard radiation prescriptions in the neoadjuvant setting consist of daily treatments over the course of five or six weeks to a total dose of 50–54 gray (Gy) [2].



Following surgical resection of pancreatic cancer, adjuvant conventionally fractionated radiotherapy with chemotherapy in select high-risk patients (i.e., positive lymph nodes and margins regardless of tumor location within the pancreas) is conditionally recommended by the American Society for Radiation Oncology.

(<https://www.practicalradonc.org/cms/10.1016/j.prro.2019.06.016/attachment/0e8abbe7-fcc6-4c5d-8b46-e81e636ce080/mmc1.pdf>. Last accessed August 19, 2021.)

Strength of Recommendation/Level of Evidence:
Conditional/low

The type and duration of chemotherapy given with radiation therapy for pancreatic cancer depends on the clinical stage, setting (neoadjuvant or adjuvant), performance status, and comorbidities. Patients with favorable performance status (0 or 1) are typically offered FOLFIRINOX prior to radiation therapy. Patients who are elderly or have a poor performance status (≥ 2) are typically offered gemcitabine or gemcitabine/nab-paclitaxel prior to radiation therapy. The duration (two to six months or longer) depends on patient tolerance and tumor response (i.e., no evidence of progression on chemotherapy). Common dose-limiting toxicities are diarrhea, neuropathy, and hematologic [12].

NEOADJUVANT THERAPY

Preoperative, or neoadjuvant, therapy is a major paradigm shift in treatment for patients with localized PDAC that offers the potential to lengthen survival while sparing patients unnecessary treatment-related morbidity using available treatments [116]. The rationale for neoadjuvant therapy differs somewhat by disease stage and clinical features.

Neoadjuvant therapy is recommended in upfront resectable disease with high-risk features of dissemination. This includes tumors in pancreas body and tail or >3 –4 cm, ascites, large regional lymph nodes, CA19-9 levels $>1,000$ U/mL, severe weight loss, and extreme pain. For these patients, staging laparoscopy is recommended to identify liver and peritoneal metastases missed by MDCT in assessing resectability, with endoscopic ultrasonography-guided biopsy [7; 11; 15]. The next step is systemic neoadjuvant therapy (i.e., chemotherapy), post-neoadjuvant therapy CA19-9, and MDCT with contrast to reassess resectability (with some limitations). If R0 resection is

feasible and there is no evidence of metastatic disease, surgery should be attempted [7; 11; 15].

In general, neoadjuvant therapy for patients who are candidates for resection is controversial [116]. Some oncology groups do not recommend neoadjuvant therapy in upfront resectable disease (except with high-risk features) until better evidence is available, but this stance has become less tenable as additional evidence supporting efficacy becomes available [7; 13; 15].

Even in patients with anatomically localized disease based on imaging and after complete resection with R0 margins, the high rates of distant failure after surgery for resectable PDAC indicates most patients already have systemic disease at the time of diagnosis. Current imaging fails to accurately assess the true burden of disease, missing occult metastases and under-staging patients [116].

Given this reality, systemic therapy is crucial, but many patients do not receive adjuvant therapy after resection. The high complication rates and potentially prolonged recovery with resection results in 25% to 50% of patients not receiving postoperative therapy [116]. However, systemic neoadjuvant therapy allows patients to receive therapy when they have better performance status and before the potential development of postoperative complications [116].

Neoadjuvant therapy also tests the tumor biology. Patients with aggressive tumors that progress and/or metastasize during neoadjuvant therapy are spared a futile operation. Due to their performance status, patients who do poorly on systemic neoadjuvant therapy would likely do poorly with surgery, resulting in mortality or serious perioperative morbidity precluding adjuvant therapy. Neoadjuvant therapy allows patients with resectable tumors who are poor surgical candidates time to medically and/or physically optimize before surgery.

Neoadjuvant therapy is not without its drawbacks. Eligibility for neoadjuvant therapy requires a tissue diagnosis, but the dense PDAC tumor stroma impedes tissue confirmation in approximately 15% of patients [116]. Further, neoadjuvant therapy means delaying surgery, with the possibility for local progression during neoadjuvant therapy into unresectable PDAC [15]. However, local progression almost always occurs concomitantly with development of systemic disease [116]. Essentially, better evidence is needed. Until phase III results are available, the poor outcomes of conventional treatment sequencing argue for the need for neoadjuvant therapy.

Borderline resectable pancreatic cancer is a recognized indication for neoadjuvant therapy, as this approach may shrink and make tumors more amenable for surgical resection with fewer complications and increased chance of R0 resection. Neoadjuvant therapy may minimize early non-detectable microscopic metastases, decrease lymph node involvement, and improve overall survival and outcomes [96].

Upfront Resectable/Borderline Resectable Tumor and Neoadjuvant Therapy

The NCCN recommends neoadjuvant therapy for patients with resectable or borderline resectable tumors. Treatment at or coordinated through a high-volume center is preferred, when feasible, and participation in a clinical trial is encouraged. The preferred neoadjuvant options are FOLFIRINOX with or without subsequent chemoradiation, or gemcitabine plus albumin-bound paclitaxel with or without subsequent chemoradiation [11]. For patients with *BRCA/PALB2* mutations, the preferred regimen is gemcitabine plus cisplatin (two to six cycles) with or without subsequent chemoradiation [11].

ASTRO guidelines for neoadjuvant chemoradiation specify a radiation dose of 4,500–5,040 cGy in 180–200 cGy fractions [12]. They recommend delivery of radiation therapy following two to six months of chemotherapy.

Locally Advanced Pancreatic Cancer and Neoadjuvant Therapy

Locally advanced pancreatic cancer accounts for 30% of newly diagnosed cases. With local involvement of adjacent critical blood vessels and presence of occult micrometastatic disease, locally advanced pancreatic cancer is generally considered surgically unresectable and incurable, and the standard of care is similar to metastatic disease [2].

However, the increased use of preoperative multiagent chemotherapy followed by chemoradiation has significantly expanded the pool of patients with locally advanced pancreatic cancer eligible for resection with curative intent, significantly improving the resectability and overall survival of these patients [117].

In a single-institution phase II trial, 49 patients with locally advanced pancreatic cancer received eight cycles of FOLFIRINOX followed by 50.4 Gy of photon radiation with capecitabine and losartan. Of these patients, 39 were brought to the operating room, 34 (69%) had their cancer removed, and of these, 30 patients (88%) had an R0 resection. Among patients who underwent resection, median progression-free survival and overall survival were 21.3 and 33 months, respectively, versus the 11- to 12-month historical overall survival [118].

Neoadjuvant therapy is associated with a downstaging-to-resection rate of greater than 30% in selected patients with locally advanced pancreatic cancer, with survival comparable to or better than initially resectable disease. For patients with arterial involvement, arterial divestment shows a lower morbidity and mortality rate than arterial resection and reconstruction [117].

Post-Neoadjuvant Therapy Restaging Evaluation of Resectability

Following neoadjuvant therapy, a restaging evaluation with pancreatic protocol MDCT is required to image tumor shrinkage and rule out local progression for resectability. However, post-neoadjuvant therapy imaging is not a reliable indicator of

resectability due to its inability to distinguish post-treatment fibrosis from residual viable tumor [117]. Post-neoadjuvant therapy CA19-9 levels are predictive of tumor regression and should be used to guide decisions about suitability for surgical exploration for resection. Diagnostic laparoscopy should be routinely used to minimize nontherapeutic surgery rates [117].

Adjuvant Chemotherapy in Patients with Resected PDAC After Neoadjuvant Therapy

After resection of pancreatic cancer following neoadjuvant FOLFIRINOX, the benefit of adjuvant chemotherapy on overall survival is unclear. Although randomized controlled trial confirmation is needed, a 2020 multicenter, retrospective study provided informative results [119]. Of 520 patients (median age: 61 years; 53.7% male) who received a median of six neoadjuvant cycles of FOLFIRINOX, 343 (66.0%) received adjuvant chemotherapy. Adjuvant chemotherapy was FOLFIRINOX for 68 patients (19.8%), gemcitabine-based chemotherapy for 201 (58.6%), capecitabine for 14 (4.1%), a combination or other agents for 45 (13.1%), and unknown for 15 patients (4.4%). The median overall survival was 38 months after diagnosis and 31 months after surgery. No survival difference was found for patients who received adjuvant chemotherapy compared with those who did not (29 months in both groups).

In multivariable analysis, the interaction of lymph node stage with adjuvant therapy was statistically significant. In patients with pathology-proven node-positive disease, adjuvant chemotherapy was associated with improved overall survival (26 months vs. 13 months). For those with node-negative disease, adjuvant chemotherapy was not associated with improved survival (38 months vs. 54 months). These results suggest that adjuvant chemotherapy after neoadjuvant therapy FOLFIRINOX and resection of pancreatic cancer was associated with improved survival only in patients with pathology-proven node-positive disease [119].

LOCALLY ADVANCED PANCREATIC CANCER

Neoadjuvant therapy increasingly shows the ability to downstage locally advanced pancreatic cancer into resectable tumor, but until such approaches are employed beyond specialized PDAC research centers, most of these patients will remain unresectable [2].

Chemotherapy selection for patients with locally advanced pancreatic cancer is largely based on extrapolation from studies in metastatic PDAC. However, the natural history of locally advanced pancreatic cancer is less predictable than metastatic disease [120]. In an important autopsy study, 28% of patients with locally advanced pancreatic cancer at initial diagnosis died with localized disease only, from complications of locally destructive tumor growth [120]. Also noted, not all isolated metastases at initial diagnosis are harbingers of widespread metastatic disease, nor the greatest threat to patient survival compared with the primary tumor or cachexia [17].

In patients with locally advanced pancreatic cancer, even with progression, treatment should not simply mirror that in metastatic disease. Rather, it should be based on the pattern of progression (locoregional vs. disseminated), prior chemotherapy and/or radiation, and sequence of therapy (as well as performance status and comorbidity). For example, if a patient with locally advanced pancreatic cancer and a history of only chemotherapy as prior treatment later develops locoregional progression, radiation may be the appropriate modality [8].

Fluoropyrimidines and gemcitabine are the most commonly used agents in adjuvant chemoradiotherapy trials of locally advanced pancreatic cancer. These studies suggest that as a radiosensitizer, capecitabine is a well-tolerated regimen with comparable or superior outcomes compared with low-dose gemcitabine [8].

There is a potential role for maintenance capecitabine or gemcitabine-based chemoradiotherapy in improving quality of life for patients with locally advanced pancreatic cancer and stable disease after 12 weeks of induction gemcitabine/capecitabine chemotherapy [8].

In contrast to conventionally fractionated chemoradiotherapy, there is growing interest in using induction chemotherapy for systemic control, followed by a short course of stereotactic body radiotherapy early during treatment with minimum disruption to systemic therapy. This could be particularly beneficial to patients with predominant local symptoms [8].

The ASCO guidelines for patients with locally advanced pancreatic cancer include several strong recommendations related to chemoradiotherapy or stereotactic body radiation therapy [2; 8]. Specifically, it states that chemoradiotherapy or stereotactic body radiation therapy may be offered upfront rather than chemotherapy [8]. This approach is recommended for patients with local progression but no metastases, performance status ≤ 2 , and favorable comorbid profile. It should also be offered to patients with response to an initial six months of chemotherapy or with stable disease who develop chemotherapy toxicities that are intolerable or cause a decline in performance status [8]. If patients respond or their disease has at least stabilized after six months of induction chemotherapy, chemoradiotherapy or stereotactic body radiation therapy may be offered as an alternative to continuing chemotherapy alone [8].

For patients with unresectable or locally advanced pancreatic cancer, definitive conventionally fractionated or dose-escalated radiation therapy with chemotherapy is used. For patients without systemic progression after four to six months (or longer) of chemotherapy, ASTRO recommends definitive radiation therapy [12]. The preferred dose is 5,040–5,600 cGy in 175–220 cGy fractions.

Local Ablative Radiation

With surgical resection considered the only potentially curative option but most patients harboring unresectable PDAC tumor, nonoperative local treatment options that can provide a similar benefit are needed. Emerging radiation techniques that address organ motion have enabled curative radiation doses delivered in patients with inoperable disease [121].

In one 2021 report, patients with locally advanced pancreatic cancer were treated with hypofractionated ablative radiation therapy, using respiratory gating, soft tissue image guidance, and other methods to address organ motion and limit the dose to surrounding luminal organs [121]. At baseline, 119 patients with locally advanced pancreatic cancer and median CA19-9 level >167 U/mL received four months of induction chemotherapy, followed by ablative radiation therapy. The median overall survival from diagnosis and ablative radiation therapy were 26.8 and 18.4 months. The 12- and 24-month overall survival following therapy were 74% and 38%, and the 12- and 24-month cumulative incidence of locoregional failure were 17.6% and 32.8% [121]. Postinduction CA19-9 decline was associated with improved locoregional control and survival. Grade 3 upper GI bleeding occurred in 10 patients (8%), with no grade 4 to 5 events. This cohort study of patients with inoperable locally advanced pancreatic cancer found that ablative radiation therapy following multiagent induction therapy was associated with durable locoregional tumor control and favorable survival [121].

METASTATIC DISEASE

Systemic chemotherapy can benefit patients with metastatic PDAC by improving disease-related symptoms and survival compared with best supportive care alone, but patients should understand that chemotherapy is palliative and not curative [80].

First-line chemotherapy in metastatic PDAC is highly consistent in clinical practice guidelines from ASCO, NCCN and ESMO. Treatment selection is based on PDAC mutation status, serum total bilirubin level, ECOG performance status, comorbidity profile, patient preference and a support system for aggressive medical therapy, and access to chemotherapy port and infusion pump management services for FOLFIRINOX or mFOLFIRINOX.

The initial chemotherapy selection for germline or somatic *HRR* gene mutation is a platinum-based chemotherapy regimen. For those with performance status ≤ 1 and serum bilirubin less than 1.5 times upper limit of normal, FOLFIRINOX or mFOLFIRINOX is preferred. Gemcitabine plus cisplatin can be used and probably has similar benefit. For patients with performance status 2, comorbidity that precludes intensive therapy, or a serum bilirubin more than 1.5 times upper limit of normal despite stenting, FOLFOX is preferred over FOLFIRINOX.

After at least 16 weeks of initial platinum-based chemotherapy without disease progression, chemotherapy should be discontinued and maintenance therapy with olaparib initiated for those with germline *BRCA* or *PALB2* mutation. For advanced PDAC with somatic (i.e., non-germline) *BRCA* or *PALB2* mutation, the benefit of olaparib maintenance therapy is not known and is under investigation.

For patients with an unknown (pending) *HRR* status, waiting until the germline or somatic mutation status is known is not recommend, given the rapidity of progression in most patients with newly diagnosed metastatic PDAC. These patients should be treated like *HRR* mutation carriers until results of genetic testing are available [80].

Patients with performance status ≤ 1 , serum bilirubin less than 1.5 times upper limit of normal, and favorable comorbidity, FOLFIRINOX is preferred, with gemcitabine plus nabpaclitaxel a potentially less toxic alternative. Patients with serum bilirubin more than 1.5 times upper limit of normal despite placement of a stent should receive FOLFOX rather than a gemcitabine-containing regimen, because gemcitabine is hepatically metabolized and associated with greater toxicity with hepatic impairment. For patients with performance status 2, favorable/adequate comorbidity, and serum bilirubin level less than 1.5 times upper limit of normal, gemcitabine monotherapy is suggested; gemcitabine/capecitabine is another option.

Highly selected patients with performance status 2 due to heavy tumor burden should be treated with gemcitabine plus nabpaclitaxel, owing to its higher response rate. Dose and schedule adjustments should be made to minimize toxicities. In patients with performance status ≥ 3 or poorly controlled comorbidity (regardless of histology or *BRCA/PALB2* mutation status), systemic chemotherapy should only be offered on an individualized, case-by-case basis; supportive care should be emphasized.

PALLIATION AND SYMPTOMATIC MANAGEMENT

At diagnosis, the median survival for patients with locally advanced, unresectable pancreatic cancer is 8 to 12 months; with metastatic disease, this decreases to 3 to 6 months. For patients with locally advanced and metastatic disease, systemic chemotherapy can improve survival. In the best outcomes to date, FOLFIRINOX demonstrated an 11.1-month median survival [122].

Patients receiving chemotherapy often report better overall quality of life, but extended survival with chemotherapy may not reduce symptom burden. Because the pancreas is located in the central abdomen at the root of the mesentery, most patients suffer from a significant symptom burden and frequently require medical attention and hospitalization for symptom management. Typical patients will require numerous

interventions targeting pain, anorexia and weight loss, depression and anxiety, biliary obstruction, gastric outlet obstruction, ascites, and venous thromboembolism [122].

All patients with newly diagnosed PDAC should have a full assessment of symptom burden, psychological status, and social supports as early as possible. Regardless of cancer stage and patient prognosis, early introduction to expert palliative and supportive care improves the social, psychological, and physical well-being of patients; decreases the intensity of medical interventions at the end of life; and ultimately improves survival [2].

Palliative care is an interdisciplinary specialty that is focused on preventing and relieving suffering, and supporting the best possible quality of life for patients and their families facing serious illness, such as pancreatic cancer. Palliative care specialist clinicians provide in-depth pain and symptom management, communication regarding goals of care, and coordinated care across settings and over time. Palliative care aims to relieve suffering in all stages of disease and can be provided in tandem with curative or life-prolonging treatments [122].

When initiated early in the disease course, palliative care improves clinical, quality of care, and survival outcomes. Furthermore, multiple studies have shown that palliative care services improve patients' symptoms, allow patients to avoid hospitalization and to remain safely and adequately cared for at home, lead to better patient and family satisfaction, and significantly reduce prolonged grief and post-traumatic stress disorder among bereaved family members. Palliative care also lowers costs and reduces rates of unnecessary hospitalizations, diagnostic and treatment interventions, and nonbeneficial intensive care when patients are near the end of life [122].

VENOUS THROMBOEMBOLISM PROPHYLAXIS

Pancreatic cancer is one of the highest-risk malignancies for venous thromboembolism (VTE), which includes deep venous thrombosis (DVT), pulmonary embolism, and visceral portal or superior mesenteric vein thrombi. The incidence of VTE is four- to seven-fold higher in PDAC. The risk is highest in the first three months after diagnosis; chemotherapy further increases the risk. In PDAC, VTE is strongly associated with higher short- and long-term mortality and high risk of recurrent VTE [122].

All patients should be educated about warning signs and symptoms of VTE. Physical examination of the legs for asymmetric pitting edema, erythema, and warmth is crucial in each office visit, and the threshold to perform a CT angiogram with tachycardia or pleuritic chest pain present should be extremely low [122].

Routine anticoagulation for primary VTE prevention is not indicated in ambulatory outpatients with pancreatic cancer and no other VTE risk factors [122]. In a patient with PDAC and documented VTE (symptomatic or incidentally found), early initiation of anticoagulation is the standard approach, and lifelong therapy should be considered. The decision to

continue anticoagulation should be balanced against bleeding risk, cost of therapy, quality of life, life expectancy, and patient preference. Low-molecular-weight heparin or oral rivaroxaban, apixaban, or edoxaban is preferred to vitamin K antagonist or unfractionated heparin for long-term anticoagulation [122].

PERI-PANCREATIC COMPLICATIONS

Bile Duct Obstruction

Endoscopic retrograde stenting is superior to surgical or percutaneous approaches to address bile duct obstruction because of a more favorable adverse event rate. Self-expandable metal stents are preferred over plastic stents in patients with a life expectancy of more than three months in terms of patency duration, less therapeutic failure and need for reintervention, lower cholangitis incidence, and better patient quality of life. Patency rates between covered and uncovered metal stents are not significantly different [16]. Endoscopic ultrasonography-guided biliary drainage is an alternative if endoscopic biliary stent placement is unsuccessful or technically not feasible.

Gastric Outlet Obstruction

In patients with gastric outlet obstruction, endoscopic duodenal stenting allows a quick resumption of oral intake, with a low complication rate and a short recovery period. However, the need for reintervention is higher after duodenal stenting compared with that of palliative surgery. Endoscopic ultrasonography-guided gastrojejunostomy is an effective and safe alternative to surgery [16].

Ascites

Ascites in patients with metastatic PDAC may be due to peritoneal metastases. In patients with locally advanced tumors, ascites may be caused by portal vein thrombus if the tumor compresses the portal vein locally [122].

Patients with malignant ascites from pancreatic cancer can experience abdominal discomfort, nausea, vomiting, and dyspnea from the pressure of the fluid against the anterior abdominal wall and diaphragm. For most patients, survival is short, and the focus is symptom control. Symptom relief from intermittent paracentesis tends to be short-lived, and the procedure must be repeated for symptom relief. If reaccumulation requires more than once-weekly paracentesis, placement of a long-term drainage catheter is an option; complication rates are higher with indwelling catheters. Diuretics such as spironolactone and furosemide decrease the absorption of water and sodium in the kidneys and may provide some symptomatic relief [122].

PAIN CONTROL INTERVENTIONS

Pancreatic cancer is one of the most painful malignancies [85]. All patients with locally advanced and metastatic pancreatic cancer should be offered aggressive treatment of pain [8]. Adequate control of pain may be unsatisfactory due to significant variation in local practice [123].

Pain is often the major presenting symptom of the disease and can be a significant feature of advanced pancreatic cancer. Patients describe a gnawing mid-epigastric pain, which radiates bilaterally under the ribs and into the mid-back, owing to the proximity of pancreatic tumors to the celiac plexus. All patients should have the level of pain and degree of pain relief from analgesics addressed at every visit [122].



The ASCO recommends that patients with metastatic pancreatic cancer should be offered aggressive treatment of the pain and symptoms of the cancer and/or the cancer-directed therapy.

(<https://ascopubs.org/doi/10.1200/JCO.20.01364>. Last accessed August 19, 2021.)

Strength of Recommendation/Level of Evidence:
Strong/intermediate

Pharmacotherapy

The mainstay of pain management is opioid therapy, and palliation of pain can often be successfully achieved by opioid analgesics alone [122]. Patients with moderate-to-severe pain should receive doses adequate to provide relief. Concern about addiction should not be a barrier to effective pain control; even with dose escalation, addiction is seldom a problem in patients with PDAC and the risk is lower than generally assumed in non-malignant pain [81; 123]. Given the ongoing concerns regarding opioid misuse in the United States, drug diversion may be a consideration. Accordingly, patients should be advised on safe storage strategies and disposal of any discontinued opioid or other controlled substance prescriptions to minimize diversion.

For patients with persistent nausea and vomiting for whom taking oral medications is difficult, pain control may be achieved using transdermal patches when adipose tissue is sufficient for transdermal absorption [122]. When pain is constant rather than intermittent, long-acting oral (e.g., morphine, oxycodone, oxymorphone) or transdermal (e.g., fentanyl, buprenorphine) preparations may work better [81]. Breakthrough pain can be treated with rapid-onset transmucosal or intranasal fentanyl formulations. Methadone may be advantageous in many patients and can be used in small doses as add-on to existing opioid treatment. Methadone should only be prescribed by clinicians who are familiar with the complex pharmacology and adverse effect profile of this opioid [123].

Laxatives should be considered for all patients on opioid analgesia for PDAC pain, because constipation is a nearly universal side effect. There is considerable individual variation in both efficacy and side effects. Not all patients benefit from or tolerate opioids. A trial of an alternative opioid may also be indicated. Cases of poor pain control or intolerable pain may benefit from continuous opioid infusion via epidural or intrathecal

catheters [81; 123]. Adjunctive treatments, such as cannabinoids, ketamine, clonidine, benzodiazepines, anti-psychotics, gabapentin, pregabalin, nortriptyline, or duloxetine, warrant consideration [122].

Near the end of life, pain management for advanced and terminal PDAC can become very challenging, and an interdisciplinary approach including palliative care specialists is needed. It is important wherever possible to consider the preferences of the patient. A range of supportive care measures can be offered, including intensive home support, home care with parenteral opioids, patient-controlled analgesia, and palliative sedation [123].

Celiac plexus neurolysis offers medium-term relief, but other procedures (e.g., splanchnicectomy) are also available. Adjunctive treatments for pain, depression, and anxiety as well as radiotherapy, endoscopic therapy, and neuromodulation may be required. Palliative chemotherapy may provide pain relief as a collateral benefit [123].

Celiac Plexus Neurolysis

Neurolytic procedures reduce pain by destruction of the afferent pathways from the pancreas to the brain. One of the most commonly used procedures is celiac plexus neurolysis.

The celiac plexus is a dense network of nerves that innervates the upper abdominal organs. Pain may be relieved by inhibiting synaptic pathways within the plexus by chemical destruction of the pathways and ganglia using dehydrated alcohol. Celiac plexus neurolysis is performed under endoscopic ultrasonography guidance [122].

Celiac plexus neurolysis improves analgesia and quality of life and decreases opioid requirements. The analgesic effect seems to vanish after eight weeks, and in most patients, pain recurs after three months. Repeated celiac plexus neurolysis benefits about 30% of patients and is normally not offered [123].

Splanchnic Nerve Neurolysis

Splanchnicectomy may disrupt more nerve pathways than celiac plexus neurolysis and is a better option when there is a large mass in the region of the celiac plexus. Splanchnicectomy is seldom performed in patients with PDAC despite some evidence of long-lasting pain relief and few complications in observational series, possibly because the expertise is not widely available [123].

Radiation Therapy

External beam radiation therapy with or without concomitant chemotherapy may also significantly alleviate pain due to local invasion of pancreatic cancer, frequently with improvement in cachexia and obstructive symptoms. However, it may take several weeks to achieve its maximal effect. When pain is caused by liver or bone metastases, patients may benefit from radiation therapy [16; 122].

CACHEXIA, WEIGHT LOSS, AND NUTRITIONAL COMPROMISE

Nutritional compromise in PDAC is common, but the underlying pathologies are diverse [2]. Nausea, caused both by the primary disease process and its associated chemotherapy, is most effectively treated with serotonin-3 receptor antagonists and atypical antipsychotics (e.g., olanzapine), with some emerging evidence suggesting efficacy with cannabinoids. Loss of appetite, even in the absence of overt nausea, is frequently reported by patients, and this symptom is driven by central pathways that are largely distinct from those that produce nausea.

Malabsorption secondary to pancreatic exocrine deficiency degrades nutritional status. Pancreatic enzyme-replacement therapy helps to stabilize weight loss and also improves quality of life by decreasing gastrointestinal symptoms. Malabsorption from biliary obstruction is a complication found in up to 90% of patients with PDAC. Similar to the replacement of pancreatic enzymes, the treatment of biliary obstruction improves symptoms beyond its effects on digestion, including anorexia, pruritus, and fatigue.

Collectively, careful attention to the nutritional status of patients with PDAC improves both their survival and quality of life. Early and regular involvement of nutrition experts in their care is recommended [2; 124].

Cancer-Related Anorexia/Cachexia Syndrome (CACS)

A constellation of disproportionate loss of lean body mass, weight loss, muscle wasting, adipose tissue reprogramming, and anorexia, cancer-related anorexia/cachexia syndrome (CACS) is more frequent in patients with PDAC than in any other malignancy due to the complex metabolic profile of pancreatic cancer [2]. In a study of 390 patients with advanced cancers, the rate of cachexia was highest in PDAC (89%), followed by gastric cancer (76%) and esophageal cancer (53%) [125].

Unlike simple starvation, which is characterized by a caloric deficiency that can be reversed with appropriate feeding, the weight loss of cachexia cannot be adequately treated with aggressive feeding [126]. The physical impact of CACS contributes to decreased patient quality of life, treatment response, and survival due to gross alterations in protein metabolism, increased oxidative stress, and systemic inflammation. The psychological impact also contributes to decreased quality of life for both patients and their families [125].

In CACS, an abnormally accelerated resting energy expenditure increases muscle protein breakdown and lipolysis, which seems related to activation of cytokines (e.g., tumor necrosis factor- α , interleukin 6 and 1 beta), and tumor-derived, potentially cachexia-inducing factors that target skeletal muscle gene products [122; 126].

Potentially Beneficial Agents

Cachexia in itself does not respond to nutritional support. There are no FDA-approved medications for treatment of CACS, and positive pharmacotherapy response in patients with anorexia associated with non-malignant disease has been difficult to translate into benefit for patients with cancer [127; 128].

Many agents have been evaluated for the treatment of CACS, but only corticosteroids (e.g., dexamethasone) and progesterone analogs (e.g., megestrol acetate) have a proven benefit in the anorexia associated with this syndrome [122]. Selection is based on life expectancy and assessment of risks versus benefits. Dexamethasone is suggested for patients for whom only weeks of therapy are anticipated, while megestrol acetate or medroxyprogesterone acetate (another progesterone analog) are suggested for patients with longer life expectancies [126].

A phase III study randomized 190 patients with advanced cancer and anorexia to megestrol acetate (480 mg/day), dexamethasone (4 mg/day), or placebo for up to four weeks. Differences in primary endpoint (at least 25% improvement in appetite) between megestrol (79.3%), dexamethasone (65.5%), and placebo (58.5%) were non-significant. Hyperglycemia and deep vein thromboses were more frequent with dexamethasone than megestrol or placebo. No other differences from placebo were found [127].

In this trial, the higher rate of deep vein thromboses with dexamethasone was unexpected. Megestrol acetate is associated with thromboembolic events and is contraindicated in patients with VTE. Dexamethasone has the potential to reduce cancer-related fatigue and elevate mood, at the significant cost of accelerating catabolic effects on muscle [127]. The primary benefits associated with these drugs are increased appetite and weight gain, not improved survival, and both drugs are associated with potential harms [122].

Mirtazapine is well-known for promoting weight gain. A placebo-controlled randomized trial found that appetite scores increased similarly with mirtazapine (15 mg at night) and placebo during the 28-day study. Mirtazapine was associated with significantly less increase in depressive symptoms and higher prevalence of somnolence than placebo, but no other differences were found [128].

The evidence of benefit in patients with CACS is inconclusive for androgens and selective androgen receptor modulators, anamorelin, cyproheptadine, long-chain omega-3 fatty acids, vitamins, minerals, and other dietary supplements, nonsteroidal anti-inflammatory drugs (NSAIDs), thalidomide, and combination approaches [126]. However, a trial of low-dose olanzapine (5 mg/day) is reasonable, particularly for patients who have concurrent nausea and/or vomiting unrelated to chemotherapy or radiation therapy [126].

Cannabis and Cannabinoids

In the cannabis plant, delta9-tetrahydrocannabinol (THC) and cannabidiol (CBD) are the best-characterized therapeutic constituents. Pharmaceutical cannabinoid products containing THC (dronabinol), a THC analog (nabilone), or THC:CBD in an oromucosal spray (nabiximols, investigational) were examined for efficacy in CACS and palliative care in two meta-analyses [126].

Unfortunately, no benefit beyond placebo was found for pharmaceutical cannabinoid products in CACS, despite their superior weight gain and appetite effects in patients with advanced HIV [129]. Cancer patients with more than 30% decrease in pain with cannabinoids compared with placebo approached significance [129].

In both meta-analyses, available studies of smoked cannabis in CACS did not meet evidence thresholds and were excluded. This limits the ability to inform real-world clinical practice, where patient preference, self-titration to tolerability/effect, access, and other factors favor smoked/vaped cannabis over single-molecule pharmaceutical cannabinoids [130].

Counseling and Support

The substantial loss of body mass can cause significant distress to patients. Although advanced cachexia is irreversible, palliating anorexia in patients with advanced cancer is best approached by focusing on stimulating appetite, supporting each person's food preferences, and avoiding prescriptive dietary advice [127].

Providing education to patients and their caregivers is crucial. The objective is to promote a shared understanding about changed goals of care, and to help reduce the distress caused by reduced oral intake [127].

Family members in particular can require educational intervention, as their distress may manifest in attempts to pressure or coerce the patient into increased feeding. Key points to discuss with patients and their family members, related to interactions about nutrition and eating near the end of life, include the following [131]:

- Loss of appetite is common in patients with advanced cancer and may be the result of the cancer process itself.
- Trying to force a patient to eat is usually counterproductive, potentially leading to increased nausea/vomiting.
- In most patients with advanced cancer and cachexia, providing additional calories by feeding tubes and/or intravenously does not improve outcomes.

- Trying to make a patient eat, when they have marked appetite loss, can lead to decreased social interactions and increased patient distress regarding interactions with caregivers (including stories of patients, in their dying days, pretending to be asleep when relatives visit, so that the relatives do not try to make them eat something).

Caregivers should be advised that it may be best to listen to and support the patient in a variety of other ways (such as giving the patient a massage or applying a lip moisturizer) instead of trying to talk them into eating more. Referral to a registered dietitian may provide patients and caregivers with additional opportunities to discuss concerns and challenges related to nutrition, appetite, and meal planning.

Diabetes Mellitus in PDAC

The presence of diabetes has been associated with higher mortality in patients with PDAC; corticosteroids can induce or exacerbate diabetes in these patients. For patients with PDAC-related diabetes, nutritional management by an experienced dietitian is essential [16]. Metformin or insulin is used as a first-line therapy. Insulin is often the preferred agent because of its efficacy, flexibility, and safety.

Careful monitoring of plasma glucose levels two hours after meals is widely recommended. The limited literature on this topic recommends maintaining blood glucose levels to avoid hypoglycemia and reduce symptoms of hyperglycemia.

Pancreatic Exocrine Insufficiency and Pancreatic Enzyme Replacement Therapy (PERT)

A contributory factor to extreme weight loss may be pancreatic exocrine insufficiency, which leads to maldigestion, fat malabsorption, and steatorrhea. The main clinical manifestation is weight loss and malnutrition, and nonspecific symptoms such as abdominal cramping, flatulence, and urgency to defecate. Fat malabsorption does not become evident until pancreatic lipase secretion falls below 10% of normal levels [122].

Pancreatic exocrine insufficiency results from loss of pancreatic parenchyma and/or tumor obstruction of the main pancreatic duct, and can occur after surgery or irradiation. The characteristic fatty stools associated with steatorrhea (loose, greasy, foul-smelling) may not be evident because patients tend to limit fat ingestion [122].

Pancreatic exocrine insufficiency is very frequent (>90% with tumors in the pancreatic head), and is associated with higher mortality in patients with unresectable PDAC. Pancreatic enzyme replacement therapy (PERT) improves survival in these patients [16]. Given its high incidence, diagnostic testing is not necessary. Patients suspected of fat malabsorption should be treated empirically with oral PERT [122].

The classical approach to patients with pancreatic exocrine insufficiency was restricting fat intake (<20 gm/day) in an attempt to reduce steatorrhea. However, this further restricts the intake of fat-soluble vitamins, which are already malabsorbed in patients with pancreatic exocrine insufficiency, and is not recommended. Frequent low-volume meals and avoidance of foods that are difficult to digest (e.g., legumes) are generally recommended [122].

Pancreatic exocrine insufficiency is treated with capsules of porcine pancreatic enzymes (pancrelipase). There are a number of commercial products available, and the amount of enzyme per capsule varies [81]. Doses are in United States Pharmacopeia (USP) units or International Units (IU); 90,000 USP is equivalent to 30,000 IU [122]. A healthy pancreas produces about 900,000 USP of lipase in response to a meal. Sufficient fat absorption can be maintained at around 10% of normal capacity; thus, roughly 90,000 USP per meal is needed. Because non-resected patients retain some pancreatic function, a starting dose of 75,000 USP with main meals and 25,000 with snacks should suffice in reducing steatorrhea and preventing weight loss. Enzymes are most effective when taken across the course of a meal. Following Whipple, patients will require 90,000 USP with meals and 45,000 USP with snacks [124].

Acidic gastric pH is normally neutralized by pancreatic bicarbonate secretion, which is absent in many patients with PDAC, especially following Whipple resection. Acid-suppressing therapy with a proton pump inhibitor is often required, as failure to neutralize gastric acid inactivates the enzymes [16; 124].

Despite recommendation from expert groups, including the NCCN, evidence suggests PERT is underutilized. This was examined in a large commercially insured U.S. population from 2001–2013. Among patients with PDAC (32,461), 1.9% had diagnostic testing for exocrine insufficiency, 21.9% filled a prescription for PERT, and 5.5% were prescribed an adequate dose (defined as $\geq 120,000$ USP lipase daily) [132].

Testing and appropriate dosing is infrequent and inconsistent in an insured U.S. population. Efforts are needed to educate medical providers on the best practices for managing exocrine pancreatic insufficiency in these patients [132].

CONSIDERATIONS FOR NON-ENGLISH-PROFICIENT PATIENTS

For patients who are not proficient in English, it is important that information regarding all aspects of their care (including diagnostic procedures and treatment options) and palliative care resources be provided in their native language, if possible. When there is an obvious disconnect in the commu-

nication process between the practitioner and patient due to the patient's lack of proficiency in the English language, an interpreter is required. Interpreters can be a valuable resource to help bridge the communication and cultural gap between patients and practitioners. Interpreters are more than passive agents who translate and transmit information back and forth from party to party. When they are enlisted and treated as part of the interdisciplinary clinical team, they serve as cultural brokers who ultimately enhance the clinical encounter. In any case in which information regarding treatment options and medication/treatment measures are being provided, the use of an interpreter should be considered. Print materials are also available in many languages, and these should be offered whenever necessary.

CONCLUSION

PDAC is the most lethal solid malignancy, predicted to become the second leading cause of cancer death in the United States by 2030. The complexity of this aggressive cancer has been vexing to investigators and tragic for patients and their families. Major research efforts over the past 50 years have only marginally improved the five-year survival rate from 6% to 10.8%. The greatest gains—from resection of early-stage tumors—are the least likely to present at diagnosis. There is an urgent need to reduce PDAC incidence through primary and secondary prevention, and mortality by accelerating therapeutic development [133].

Until diagnostic or therapeutics breakthroughs arrive, novel uses of standard treatments (i.e., neoadjuvant therapy) show survival advantages for a greater number of patients. The longest survival reported by a phase III trial was published in 2018—a median 54.4 months in patients who received resection followed by mFOLFIRINOX [113]. Many novel treatments are in phase III trials. Additional approaches to manage morbidities and provide better palliative care are also needed. Cancer anorexia/cachexia is a high-priority area.

It is now clear that even early-stage PDAC is a systemic disease and that new-onset metabolic (e.g., diabetes, anorexia/cachexia, hyperglycemia) and neuropsychiatric (e.g., depression, fatigue) symptoms/syndromes are prodromal rather than comorbid or secondary. This recognition has also called for a re-thinking of pancreatic cancer from a more integrative, multi-system perspective [2].

Customer Information/Evaluation insert located between pages 88–89.

SELF-ASSESSMENT QUESTIONS

#90240 PANCREATIC CANCER

After reviewing the course, complete the following Self-Assessment Questions. Receive immediate feedback by reviewing the Study Guide provided on pages 69–70. Please record the number of questions you correctly answered on the Evaluation. The Evaluation must be completed in order to receive credit for this course.

This 10 contact hour activity must be completed by August 31, 2024.

- The median age at diagnosis of pancreatic cancer is**
 - 20 years.
 - 40 years.
 - 55 years.
 - 70 years.
- Of the following, which has the highest relative risk for pancreatic cancer?**
 - Obesity
 - Family history
 - Cigarette smoking
 - Chronic pancreatitis
- Which of the following statements regarding diet and pancreatic cancer risk is FALSE?**
 - High plasma 25-hydroxyvitamin D levels increase the risk for pancreatic cancer.
 - Pancreatic cancer incidence may be lower in persons with higher intake of fresh fruits.
 - Pancreatic cancer incidence may be lower in persons with higher intake of vegetables rich in folate and lycopenes.
 - There is some evidence that higher consumption of red/processed meat is associated with elevation in pancreatic cancer risk.
- Starting at 30 months prediagnosis, pancreatic ductal adenocarcinoma (PDAC) showed three distinct metabolic phases. Which of the following is NOT one of these phases?**
 - Cachexia
 - Pancreatitis
 - Pre-cachexia
 - Hyperglycemia
- The most common precancerous precursor lesions associated with pancreatic cancer are**
 - mucinous cystic neoplasms (MCN).
 - cervical intraepithelial neoplasia (CIN).
 - pancreatic intraepithelial neoplasia (PanIN).
 - intraductal papillary mucinous neoplasm (IPMN).
- Which of the following statements regarding pancreatic cancer screening is TRUE?**
 - Hereditary pancreatitis is a clear indication for annual pancreatic cancer screening and surveillance.
 - Population screening to achieve earlier detection and intervention of PDAC is federal policy in the United States.
 - The U.S. Preventive Services Task Force recommends against screening for pancreatic cancer in asymptomatic adults.
 - The International Cancer of the Pancreas Screening (CAPS) Consortium recommends pancreatic surveillance for all adults to detect and resect early pancreatic cancer and its high-grade precursors.
- All patients with pancreatic cancer should have germline testing and gene profiling offered**
 - only if requested by the patient.
 - after one year of unsuccessful therapy.
 - only if they have a positive family history.
 - as quickly as possible after diagnosis.
- Most pancreatic cancers (approximately 75%) originate in the**
 - liver.
 - tail of the pancreas.
 - body of the pancreas.
 - head of the pancreas.

9. In one study, the most common symptom of early-stage PDAC was
- A) nausea.
 - B) fatigue.
 - C) steatorrhea.
 - D) abdominal pain.
10. Which of the following should trigger inclusion of PDAC in the differential diagnosis?
- A) Ongoing hypoglycemia with significant weight gain
 - B) The constellation of abdominal pain, hyperkalemia, and weight loss
 - C) Longstanding hyperglycemia regardless of other symptoms or diagnoses
 - D) Development of abdominal pain, jaundice, or weight loss in the context of newly diagnosed diabetes, family history of PDAC, or history of pancreatitis
11. Which of the following psychiatric issues is 11 times higher in men with PDAC compared with the general population?
- A) Suicide
 - B) Attention deficit
 - C) Aggression and violence
 - D) Borderline personality disorder
12. Referral for genetic counseling should be considered for patients diagnosed with pancreatic cancer, especially those
- A) of Ashkenazi Jewish ancestry.
 - B) with a family history of cancer or who are young.
 - C) for whom a hereditary cancer syndrome is suspect.
 - D) All of the above
13. Which of the following is the most common sign/symptom in patients with pancreatic cancer?
- A) Jaundice
 - B) Polydipsia
 - C) Abdominal pain
 - D) Significant weight gain
14. Which of the following statements regarding pancreatic cancer cachexia is TRUE?
- A) It rarely develops in patients with PDAC.
 - B) It typically responds well to therapeutic interventions.
 - C) It is characterized by an increase in skeletal muscle mass.
 - D) It is commonly associated with adipose tissue wasting and anorexia.
15. Which of the following is a common site of metastases in patients with advanced pancreatic cancer?
- A) Liver
 - B) Brain
 - C) Colon
 - D) Bladder
16. All of the following are common laboratory abnormalities in patients with pancreatic cancer, EXCEPT:
- A) Mild anemia
 - B) Decreased serum amylase
 - C) Elevated serum bilirubin levels
 - D) Elevated alkaline phosphatase levels
17. What is the preferred imaging for initial evaluation of suspected PDAC?
- A) Ultrasonography
 - B) Endoscopic retrograde cholangiopancreatography (ERCP)
 - C) Multidetector computed tomography (MDCT) angiography with IV contrast
 - D) Magnetic resonance imaging/magnetic resonance cholangiopancreatography (MRI/MRCP)
18. In patients with PDAC, biopsy
- A) can interfere with definitive surgery.
 - B) is required before undergoing surgery.
 - C) does not result in seeding if proper protocols are followed.
 - D) will appropriately delay surgical resection if nondiagnostic.
19. Although unsuitable for asymptomatic screening, which biomarker is the most clinically useful in PDAC?
- A) Alpha-fetoprotein (AFP)
 - B) Carcinoembryonic antigen (CEA)
 - C) Carbohydrate antigen 125 (CA-125)
 - D) Carbohydrate antigen 19-9 (CA19-9)
20. Patients with a pancreatic tumor 1–2 cm in greatest dimension would be staged
- A) Tis.
 - B) T1c.
 - C) T2.
 - D) T3.

Self-Assessment questions continue on next page →

21. According to the NCCN guideline, PDAC tumors that are involved with nearby structures would be classed as
- A) Resectable
 - B) Localized and complicated
 - C) Metastatic (non-resectable)
 - D) Borderline resectable or locally advanced
22. A patient with pancreatic cancer who is capable of all self-care but unable to carry out any work activities would receive an Eastern Cooperative Oncology Group Performance Status (ECOG) score of
- A) 1.
 - B) 2.
 - C) 3.
 - D) 4.
23. Which of the following is considered a curative treatment for PDAC?
- A) Chemotherapy
 - B) Radical surgery
 - C) Combination chemotherapy and radiation therapy
 - D) There is no potentially curative treatment for PDAC.
24. The conventional Whipple procedure (pancreaticoduodenectomy) is used for tumors
- A) growing multifocally.
 - B) in the pancreatic body or tail.
 - C) in the pancreatic head or periampullary region.
 - D) in the neck of the pancreas, especially with extension into the body or tail.
25. The cycle length of FOLFIRINOX chemotherapy for pancreatic cancer is
- A) 7 days.
 - B) 14 days.
 - C) 1 month.
 - D) 3 months.
26. Which chemotherapy agent/regimen has the strongest recommendation and level of evidence for use in patients with stage 3 (locally advanced) PDAC?
- A) Gemcitabine
 - B) Capecitabine
 - C) FOLFIRINOX
 - D) 5-FU continuous infusion
27. ASTRO guidelines for neoadjuvant chemoradiation specify a radiation dose of
- A) 450–500 cGy in 18–20 cGy fractions.
 - B) 1,500–2,040 cGy in 80–120 cGy fractions.
 - C) 4,500–5,040 cGy in 180–200 cGy fractions.
 - D) 6,500–7,400 cGy in 280–300 cGy fractions.
28. To address bile duct obstruction in patients with PDAC, the preferred approach is
- A) ERCP.
 - B) cholecystectomy.
 - C) percutaneous approaches.
 - D) endoscopic retrograde stenting.
29. Which agents have proven efficacy in the treatment of anorexia associated with cancer-related anorexia/cachexia syndrome?
- A) Cannabinoids
 - B) Antidepressants
 - C) Insulin and ghrelin
 - D) Corticosteroids and progesterone analogs
30. When providing education to caregivers of patients with PDAC, which of the following points related to nutrition should be emphasized?
- A) Trying to force a patient to eat is usually counterproductive, potentially leading to increased nausea/vomiting.
 - B) Loss of appetite is common in patients with advanced cancer and may be the result of the cancer process itself.
 - C) Trying to make a patient eat, when they have marked appetite loss, can lead to decreased social interactions and increased patient distress regarding interactions with caregivers.
 - D) All of the above

STUDY GUIDE

COURSE #90240 PANCREATIC CANCER

This Study Guide has been included to provide immediate feedback about your achievement for the course learning objectives. The correct answer for the Self-Assessment Questions and their locations within the text are indicated below.

- The median age at diagnosis of pancreatic cancer is**
D) 70 years. (Epidemiology)
- Of the following, which has the highest relative risk for pancreatic cancer?**
D) Chronic pancreatitis (Epidemiology; Table 2)
- Which of the following statements regarding diet and pancreatic cancer risk is FALSE?**
A) High plasma 25-hydroxyvitamin D levels increase the risk for pancreatic cancer. (Epidemiology; Non-Genetic Risk Factors; Diet)
- Starting at 30 months prediagnosis, pancreatic ductal adenocarcinoma (PDAC) showed three distinct metabolic phases. Which of the following is NOT one of these phases?**
B) Pancreatitis (Epidemiology; Non-Genetic Risk Factors; Prediagnostic Metabolic and Soft Tissue Changes)
- The most common precancerous precursor lesions associated with pancreatic cancer are**
C) pancreatic intraepithelial neoplasia (PanIN). (Pathophysiology; Mutational Sequence of PDAC Development)
- Which of the following statements regarding pancreatic cancer screening is TRUE?**
C) The U.S. Preventive Services Task Force recommends against screening for pancreatic cancer in asymptomatic adults. (Pancreatic Cancer Screening)
- All patients with pancreatic cancer should have germline testing and gene profiling offered**
D) as quickly as possible after diagnosis. (Pancreatic Cancer Screening; Germline and Somatic Testing and Molecular Analysis)
- Most pancreatic cancers (approximately 75%) originate in the**
D) head of the pancreas. (Clinical Evaluation of Pancreatic Cancer)
- In one study, the most common symptom of early-stage PDAC was**
B) fatigue. (Clinical Evaluation of Pancreatic Cancer)
- Which of the following should trigger inclusion of PDAC in the differential diagnosis?**
D) Development of abdominal pain, jaundice, or weight loss in the context of newly diagnosed diabetes, family history of PDAC, or history of pancreatitis (Clinical Evaluation of Pancreatic Cancer)
- Which of the following psychiatric issues is 11 times higher in men with PDAC compared with the general population?**
A) Suicide (Clinical Evaluation of Pancreatic Cancer; Neuropsychiatric Symptoms and Pancreatic Cancer)
- Referral for genetic counseling should be considered for patients diagnosed with pancreatic cancer, especially those**
D) All of the above (Clinical Evaluation of Pancreatic Cancer; Family History)
- Which of the following is the most common sign/symptom in patients with pancreatic cancer?**
C) Abdominal pain (Clinical Evaluation of Pancreatic Cancer; Common Presenting Symptoms and Signs; Abdominal Pain)
- Which of the following statements regarding pancreatic cancer cachexia is TRUE?**
D) It is commonly associated with adipose tissue wasting and anorexia. (Clinical Evaluation of Pancreatic Cancer; Common Presenting Symptoms and Signs; Cachexia)

15. Which of the following is a common site of metastases in patients with advanced pancreatic cancer?
A) Liver (*Clinical Evaluation of Pancreatic Cancer; Metastatic Disease*)
16. All of the following are common laboratory abnormalities in patients with pancreatic cancer, EXCEPT:
B) Decreased serum amylase (*Clinical Evaluation of Pancreatic Cancer; Laboratory Testing*)
17. What is the preferred imaging for initial evaluation of suspected PDAC?
C) Multidetector computed tomography (MDCT) angiography with IV contrast (*The Diagnostic and Staging Workup; Imaging; Multidetector Computed Tomography*)
18. In patients with PDAC, biopsy
A) can interfere with definitive surgery. (*The Diagnostic and Staging Workup; Biopsy*)
19. Although unsuitable for asymptomatic screening, which biomarker is the most clinically useful in PDAC?
D) Carbohydrate antigen 19-9 (CA19-9) (*The Diagnostic and Staging Workup; Carbohydrate Antigen 19-9 (CA19-9)*)
20. Patients with a pancreatic tumor 1–2 cm in greatest dimension would be staged
B) T1c. (*The Diagnostic and Staging Workup; Table 6*)
21. According to the NCCN guideline, PDAC tumors that are involved with nearby structures would be classed as
D) Borderline resectable or locally advanced (*The Diagnostic and Staging Workup; The Staging Workup; Resectability Assessment*)
22. A patient with pancreatic cancer who is capable of all self-care but unable to carry out any work activities would receive an Eastern Cooperative Oncology Group Performance Status (ECOG) score of
B) 2. (*Treatment Approaches for Pancreatic Cancer; Table 8*)
23. Which of the following is considered a curative treatment for PDAC?
B) Radical surgery (*Treatment Approaches for Pancreatic Cancer; Recommended Treatment Options by Clinical Stage*)
24. The conventional Whipple procedure (pancreaticoduodenectomy) is used for tumors
C) in the pancreatic head or periampullary region. (*Treatment Approaches for Pancreatic Cancer; Resection of Pancreatic Cancer; Approaches; Pancreaticoduodenectomy (Whipple Procedure)*)
25. The cycle length of FOLFIRINOX chemotherapy for pancreatic cancer is
B) 14 days. (*Treatment Approaches for Pancreatic Cancer; Table 9*)
26. Which chemotherapy agent/regimen has the strongest recommendation and level of evidence for use in patients with stage 3 (locally advanced) PDAC?
A) Gemcitabine (*Treatment Approaches for Pancreatic Cancer; Table 11*)
27. ASTRO guidelines for neoadjuvant chemoradiation specify a radiation dose of
C) 4,500–5,040 cGy in 180–200 cGy fractions. (*Treatment Approaches for Pancreatic Cancer; Neoadjuvant Therapy; Upfront Resectable/Borderline Resectable Tumor and Neoadjuvant Therapy*)
28. To address bile duct obstruction in patients with PDAC, the preferred approach is
D) endoscopic retrograde stenting. (*Palliation and Symptomatic Management; Peri-Pancreatic Complications; Bile Duct Obstruction*)
29. Which agents have proven efficacy in the treatment of anorexia associated with cancer-related anorexia/cachexia syndrome?
D) Corticosteroids and progesterone analogs (*Treatment Approaches for Pancreatic Cancer; Cachexia, Weight Loss, and Nutritional Compromise; Cancer-Related Anorexia/Cachexia Syndrome (CACS); Potentially Beneficial Agents*)
30. When providing education to caregivers of patients with PDAC, which of the following points related to nutrition should be emphasized?
D) All of the above (*Treatment Approaches for Pancreatic Cancer; Cachexia, Weight Loss, and Nutritional Compromise; Cancer-Related Anorexia/Cachexia Syndrome (CACS); Counseling and Support*)

Pathophysiology: Muscles, Joints, and Connective Tissues

This course meets one hour of Pain Management education.

Audience

This course is designed for nurses in all practice settings.

Course Objective

As health care becomes more complex, it is essential that the theoretical concepts of the basis of illness (pathophysiology) be well understood. The purpose of this course is to reinforce the scientific rationales for the interventions nurses perform and the decisions nurses make as patients move through the ever-changing struggle with their illness.

Learning Objectives

Upon completion of this course, you should be able to:

1. Describe the structure and function of the muscles, joints, and connective tissues.
2. Discuss the pathophysiologic influences that may affect the muscles, joints, and connective tissues.
3. Outline the role of subjective data in completing a full nursing assessment of the muscles, joints, and connective tissues.
4. Describe objective data compiled during a nursing assessment of the muscles, joints, and connective tissues.
5. Identify imaging and diagnostic studies used in the identification and classification of muscles, joints, and connective tissues.
6. Discuss genetic conditions manifesting in the muscles and connective tissues.
7. Evaluate the presentation and differential diagnosis of inflammatory muscle and connective tissue disorders.
8. Describe the clinical presentation and treatment of immunologic disorders of the muscles and connective tissues.
9. Review the assessment and treatment of traumatic conditions of the muscles and connective tissue.
10. Discuss disorders of the joints with multifactorial origin.
11. Analyze the manifestations and therapeutic approaches for degenerative joint diseases.
12. Outline the presentation, treatment, and nursing considerations for patients with immunologic joint conditions, such as rheumatoid arthritis.
13. Compare and contrast the various joint diseases with an infectious origin.
14. Describe cancers of the joints, muscle, and connective tissues.
15. Evaluate the appropriate assessment and management of traumatic joint injuries.

Faculty

Jane C. Norman, RN, MSN, CNE, PhD, received her undergraduate education at the University of Tennessee, Knoxville campus. There she completed a double major in Sociology and English. She completed an Associate of Science in Nursing at the University of Tennessee, Nashville campus and began her nursing career at Vanderbilt University Medical Center. Jane received her Masters in Medical-Surgical Nursing from Vanderbilt University. In 1978, she took her first faculty position and served as program director for an associate degree program. In 1982, she received her PhD in Higher Education Administration from Peabody College of Vanderbilt University. In 1988, Dr. Norman took a position at Tennessee State University. There she has achieved tenure and full professor status. She is a member of Sigma Theta Tau National Nursing Honors Society. In 2005, she began her current position as Director of the Masters of Science in Nursing Program.

Faculty Disclosure

Contributing faculty, Jane C. Norman, RN, MSN, CNE, PhD, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Division Planner

Sharon Cannon, RN, EdD, ANEF

Senior Director of Development and Academic Affairs

Sarah Campbell

Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

Accreditations & Approvals



In support of improving patient care, NetCE is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American

Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Designations of Credit

NetCE designates this continuing education activity for 15 ANCC contact hours.

AACN Synergy CERP Category A.

Individual State Nursing Approvals

In addition to states that accept ANCC, NetCE is approved as a provider of continuing education in nursing by: Alabama, Provider #ABNP0353 (valid through 07/29/2025); Arkansas, Provider #50-2405; California, BRN Provider #CEP9784; California, LVN Provider #V10662; California, PT Provider #V10842; District of Columbia, Provider #50-2405; Florida, Provider #50-2405; Georgia, Provider #50-2405; Kentucky, Provider #7-0054 (valid through 12/31/2023); South Carolina, Provider #50-2405; West Virginia, RN and APRN Provider #50-2405.

About the Sponsor

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

Disclosure Statement

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

How to Receive Credit

- Read the following course.
- Complete the Self-Assessment Questions at the end of the course. (No test required.)
- Complete the Evaluation.
- Return your Customer Information/Evaluation and payment to NetCE by mail or fax, or complete online at www.NetCE.com/PA23.
- A full Works Cited list is available online at www.NetCE.com.



Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the study questions and course material for better application to your daily practice.

INTRODUCTION

Along with the bones, muscles, ligaments, tendons, cartilage, and the joints provide the body with a supportive framework that allows flexibility of movement and protects the internal organs. These tissues also give shape to the body and act partially as a storage and supply area for minerals. When the tissues are unable to perform their usual functions because of trauma or rheumatic, inflammatory, or degenerative conditions, a person's physical support, protection, mobility, and ability to carry out activities of daily living are affected.

MUSCLES, JOINTS, AND CONNECTIVE TISSUES: STRUCTURAL AND FUNCTIONAL INTER-RELATIONSHIPS

The musculoskeletal system is composed of many anatomical structures that work together to produce movement, support, and protection of the body and its parts. These structures include the bones and joints of the skeletal system; the skeletal muscles; and the tendons, ligaments, and other elements that connect these tissues. This course will focus on the components of the system excluding the bones.

STRUCTURE AND FUNCTION OF SKELETAL MUSCLES

Contraction of skeletal muscle is its primary function, with the intent of moving the bones of the skeleton. Bone serves as a lever, the joint serves as a fulcrum upon which the bone pivots, and the muscle provides the force that moves the lever. A second function of skeletal muscles is maintenance of body posture. A residual amount of contraction in the muscles, known as muscle tone, serves to keep the body erect. A third function is heat production. To combat hypothermia, small, rapid contractions of skeletal muscle (shivering) produce body heat [1].

Producing Skeletal Movement

A typical skeletal muscle is anchored at each end to bone by a tendon. The muscle often stretches across a joint. The muscle's attachment to the less movable bone is called its origin, and its attachment to the more movable bone is called its insertion. When the muscle contracts, one bone remains more or less stationary, forcing the other bone to move [2].

Most skeletal muscles work in groups. The prime mover is the muscle that contracts to produce the movement. Synergists are muscles that work with prime movers to assist in performing the movement. Antagonists are muscles that work opposite prime movers by relaxing during their contraction or by producing an opposite effect. For example, the arm is flexed by contracting the biceps brachia, which acts as the prime mover; at the same time, the triceps brachii on the opposite side of the humerus relaxes, acting as the antagonist (**Figure 1**). When the arm is extended, the roles of the biceps and triceps are reversed. An isotonic contraction occurs when a muscle shortens during contraction. An isometric contraction occurs when a muscle becomes tense while remaining the same length [2].

Skeletal Muscle Structure

Muscle—skeletal, smooth, and cardiac—is made up of elongated cells called fibers (**Figure 2**). The fibers contain strands of contractile protein called myofibrils that extend the length of the cell. At the neuromuscular junction, the chemical acetylcholine creates the stimulus for muscle-nerve conduction of movement. Skeletal muscle fibers are multinucleated, and their myofibrils have striations: light and dark bands perpendicular to the long axis of the cell. The dark bands (anisotropic or A strands) are composed of the protein myosin, and the light bands (isotropic or I strands) contain the protein actin. A sense fibrous line called the Z line crosses the center of each I band and divides the myofibrils into a series of repeating units called sarcomeres. The bands are visible to the unaided eye and give skeletal muscle its alternate name: striated muscle. Smooth and cardiac muscles are made up of uninucleated cells. They further differ from skeletal muscle in that smooth muscle has tapered fibers with no striations and cardiac muscle has branched fibers [2].

Muscle fibers are bound together by connective tissue into small bundles called fascicles, visible to the unaided eye. Fascicles are bound into larger bundles, which collectively form the muscle. The entire muscle is enclosed by a connective tissue covering called the epimysium, which is continuous with the connective tissue surrounding the fascicles and fibers. The epimysium is also continuous with the tendon or other connective tissue at attachment of muscle to bone. Thus, there is a continuous network of connective tissue extending from individual muscle fibers to the tendon. Blood vessels and nerves penetrate the connective tissue of the muscle, so muscle has sufficient blood supply to furnish nutrients and oxygen and to remove the waste products of muscular activity [2; 3].

OTHER CONNECTIVE TISSUE STRUCTURES


Tendons are cords of connective tissue that attach muscles to the periosteum of the bones. During muscle contraction, the muscle pulls the tendon, which pulls the bone to which it is attached, producing movement. Flexion, extension, adduction, and abduction are normal movements of muscles and bones.

Ligaments, made of fibrous connective tissue, connect bones to one another. They have the ability to stretch while providing stability. The knee joint, for example, is stabilized by ligaments, such as the anterior and posterior cruciate ligaments, which bind the femur to the tibia within the joint capsule, and by the medial and lateral collateral ligaments outside the joint capsule [4].

A bursa is a fluid-filled sac that facilitates motion of structures that move against each other. It can be found between skin and bone, muscle and bone, tendons and bone, ligaments and bone, and between muscles. The bursae function as padding between structures to reduce friction caused by moving parts [4].

Connective tissue, in the broad sense of the term, include all tissues made up of cells in a matrix, including bone, cartilage, blood, and lymph. However, the term is used in a more limited sense when discussing diseases of the connective tissues. In this sense, connective tissue means the binding and covering tissues of the body, including tendons, ligaments, muscle fascia, and the deep layers of the skin. This kind of connective tissue (sometimes called “connective tissue proper”) is essential in holding together all the components of the musculoskeletal system. Also included are intervertebral discs (or intervertebral fibrocartilage), which serve as “shock absorbers” to cushion the spine and help it move [4; 5].


MUSCLES OF THE UPPER EXTREMITY



Source: National Cancer Institute

Figure 1

STRUCTURE OF A SKELETAL MUSCLE



Source: National Cancer Institute

Figure 2

THE PROCESS OF SKELETAL MUSCLE CONTRACTION

Skeletal muscle contraction begins with the stimulus of a muscle fiber by a motor neuron. Every motor neuron ends in many fine branches, with each branch connecting with an individual muscle fiber. A group of muscle fibers activated by a single motor neuron is called a motor unit. Motor units range in size from a single muscle fiber in muscles controlling fine, skilled movements to over one hundred fibers in muscles involved in gross movements. All the fibers of a motor unit contract together when the neuron is stimulated [6]. There are two types of motor units in skeletal muscle, Type 1 and Type 2. Type 1 has a small cell diameter, with a high excitability and fast conduction velocity. It has an oxidative profile with moderate contraction velocity and low fatigability. There are few muscle fibers of this type. In contrast, Type 2 has a large cell diameter, with low excitability but a very fast conduction velocity. Type 2 fibers are numerous in quantity, with a glycolytic profile and high fatigability. The small motor units, with Type 1 (also known as “slow-twitch”) fibers, are recruited first and are frequently active, while the large motor units, with Type 2 (“fast-twitch”) fibers, are used infrequently, in forceful contractions. Maximal efforts, in which fast motor units are recruited, cannot be sustained because of the rapid depletion of glycogen.

When a nerve impulse reaches the end of a motor neuron, small vesicles in the ends of the nerve branches release acetylcholine, which increases the permeability of the muscle cell and causes an influx of calcium ions into the cell. The calcium ions cause structural changes in the myofilaments that allow them to slide past each other, causing contraction. The structural changes also allow breakdown of adenosine triphosphate (ATP) to adenosine diphosphate (ADP) to provide energy for the contraction. The muscle relaxes as a result of the action of the enzyme cholinesterase, which breaks down acetylcholine, allowing the muscle to return to its resting state [6].

At the beginning of muscle contraction, ATP is formed from creatine phosphate stored in the muscle. The supply of creatine phosphate is limited, however, and even with mild muscle activity, additional ATP must be formed from ADP. The energy for forming this additional ATP is supplied by respiration. The first step in respiration is glycolysis, or anaerobic respiration, which produces lactic acid and small amounts of ATP. Under normal conditions, the lactic acid is broken down further by aerobic respiration, which requires an oxygen supply. The final products of aerobic respiration are carbon dioxide, water, and large amounts of ATP [6].

During sustained strenuous exercise, the blood cannot supply enough oxygen to keep pace with glycolysis, and lactic acid accumulates in the muscle, causing an oxygen debt. Muscle contractions continue for a short time using the small amount of ATP produced by glycolysis, but soon the demand exceeds

the supply and the muscle is fatigued. The contractions decrease in strength and then stop. The pain of muscle fatigue is the result of accumulated lactic acid. Oxidation of excess lactic acid occurs after exercise, when the person breathes deeply to pay off the oxygen debt [6].

The effects of exercise on the body’s cells are significant. Physical activity increases the size and number of mitochondria, increases muscle’s ability to use fat as a source of energy, increases the size of muscle fibers, and increases the content of myoglobin in muscle fibers. Exercise also results in increased fat oxidation. All of these increases lead to hypertrophy of the muscle, which leads to an increase in strength of the muscle. The wasting of muscle due to lack of use is assessed as atrophy

PATHOPHYSIOLOGIC INFLUENCES AND EFFECTS

The primary function of the musculature and connective tissues of the body is to provide body movement. When disease or trauma alters the system, the individual’s ability to move and ambulate can be affected, which can profoundly affect a person’s lifestyle. Movement is often still possible, but not without pain or difficulty [7].

INFLAMMATION

Inflammation may occur in muscle or connective structures as a result of excessive or repeated strain or pathogenic invasion. Restricted motion and pain usually result. One such example is rotator cuff injury, when the patient is unable to abduct the arm because of pain and muscle spasms. Other connective tissues of the body may be affected by inflammation, resulting in changes in other organs as well as the musculoskeletal system. Many of these connective tissue disorders are believed to be associated with immune processes [7].

DEGENERATIVE CHANGES

The joint is the musculoskeletal structure most frequently influenced by degenerative disease. Changes are most often associated with aging, excess weight, trauma, and inflammatory conditions. In the presence of these factors, articular cartilage softens, thins, and ulcerates, and the joint surfaces become rough. There may be a narrowing of the joint space and swelling of adjacent soft tissue. The normal smooth-gliding joint action is diminished, and the periosteum becomes irritated by friction, stimulating the growth of bone spurs at the joint margins. The effects of this destruction include joint pain, stiffness, and joint deformity, which can result in slight to moderate limitation of movement. Crunching or grating sounds, called crepitus, may be heard upon movement [8; 9].

The intervertebral discs can also be affected by degeneration. The water content of the discs decreases with age, causing them to become thinner. The surrounding ligaments also change with age, so the disc becomes unstable. These changes along with increased bone resorption cause decreased height and

painless restriction of spinal movement in the elderly. In some cases, the condition becomes more severe, with pressure on nerves causing pain and neurologic deficits [8; 9].

Somewhat akin to degeneration is the process of atrophy. Muscle can atrophy as a result of disuse. As noted, the normal strain on muscles contributes to their development and to the maintenance of their size, shape, strength, and composition. Through disuse, muscle cells become reduced in size and weakened, and the muscle mass becomes more fibrous. Inactivity can also lead to joint contracture; the muscle fibers become shortened and fixed, and the joint's range of motion becomes limited. These conditions are reversible with the resumption of activity. However, contractures can progress to an irreversible state without treatment [8; 9].

INFECTION

Musculoskeletal structures, such as joints and bursae, can be infected by pathogens entering from penetrating wounds or via the circulation. Pain and restricted motion are common in these cases [10].

NEOPLASIA

Malignant neoplasms of the bone, muscle, and cartilage are called sarcomas. Cancer affecting the muscle is called rhabdomyosarcoma; chondrosarcoma originates in the bones but can extend to the cartilage [11]. Depending on the specific cancer and location, patients may experience a temporary limitation in mobility (e.g., following surgery for tumor removal) or permanent limitation due to extensive surgical intervention, such as amputation [11].

TRAUMA

Skeletal muscle can be injured by trauma. Fortunately, skeletal muscle fibers can regenerate, but when the damage is extensive, the fibers are replaced by scar tissue. Trauma to the musculoskeletal structures supporting the joints is common. Muscle fibers may be injured due to overuse, overstretching, forcible twisting and other abnormal movement. The fibers may be torn, or stretched too far, and joint surfaces may dislocate, that is, separate partially or completely. Associated blood vessels and nerves may be damaged in the process. Pain and limited motion are the result [12].

Direct muscle trauma, overuse, or exposure to high temperatures can induce rhabdomyolysis. Rhabdomyolysis is a complex syndrome involving the rapid dissolution of damaged skeletal muscle, resulting in the leakage of intracellular contents to such an extent that it results in organ (particularly kidney) damage.

RELATED INFLUENCES AND EFFECTS

NEUROLOGIC AND VASCULAR PROBLEMS

Neurologic and vascular problems can cause or contribute to connective tissue and muscle disorders. Because muscle functioning is the result of the combined effect of muscle

fibers and motor nerves, neurologic damage or interference can impair muscle functioning, causing atrophy and paralysis. Likewise, disruption of the vascular supply to these tissues can limit the nutrient and oxygen supply to cells and interfere with removal of cellular waste products. Prolonged interruption of circulation leads to necrosis [13; 14].

Connective tissue disorders can also give rise to neurologic or vascular problems, which may in turn cause further musculoskeletal damage. Pressure from bandages, traction devices, tumor growth, and poor positioning are a few problems that can hinder nerve and blood vessel functioning. Trauma to muscles causes edema and hemorrhage in soft tissues, increasing the pressure within a confined space. Pressure on nerves and blood vessels in the area can become so great as to produce irreversible necrosis of the muscle tissue. A permanently disabling contracture of the limb may occur, as well as loss of motor and sensory functioning [13; 14].

OCCUPATION AND LIFESTYLE

A person's occupation and lifestyle can contribute to alterations in the muscular and connective tissues. Interest in physical fitness has prompted many people to become active in athletic endeavors. Highly athletic activities, including weightlifting, distance running, and more intense sports, are associated with an increased risk for injury, particularly with improper conditioning and training [15].

Sport injuries can generally be categorized as acute or overuse. Acute injuries occur most often in contact sports and include strains, sprains, and dislocations. Overuse injuries are usually a result of repetitive motions or excessive intensity or duration of exercise. Acute injuries are typically traumatic (e.g., ligament tears), while the most common overuse injuries are tendinitis and osteoarthritis [15].

With muscle and connective tissue disorders, patients may be unable to continue their usual recreational activities. Further, roles within the family may change to accommodate impaired ability to conduct usual activities of daily living. Occasionally, it may be necessary to use assistive devices or to modify the environment, which requires a period of adjustment [15].

NURSING ASSESSMENT: ESTABLISHING THE DATA BASE

The nursing assessment of patients with muscle, joint, and/or connective tissue disorders requires special emphasis on the musculoskeletal, neurologic, and vascular systems [16].

SUBJECTIVE DATA

As part of any nurse assessment, patients should provide important information about what they are experiencing as a result of their conditions.

Pain

Pain, in some cases severe, is a common manifestation of joint, muscle, and connective tissue problems. Patients should be asked to describe their pain thoroughly, including location, intensity, quality, duration, radiation, precipitating factors, and successful relief measures. Some patients ache all over and should indicate each of the areas involved. Knowing the quality of pain may help pinpoint a specific problem, but the patient may require help in describing the pain. All these data are helpful in reaching a diagnosis [16].

Some patients experience pain so severe they cannot tolerate moving or being touched. Others have learned to live with chronic pain. It is important to pay attention to descriptions of pain that seem unusual or excessive for the patient's condition; such complaints warrant a thorough assessment. Changes in pain status may indicate a new or undiagnosed condition [16].

Paresthesia

Some patients with musculoskeletal conditions will experience paresthesia, such as tingling, numbness, and/or diminished or absent sensation. The affected area should be defined as precisely as possible. Paresthesia is an indication of a neurologic problem and requires an in-depth assessment [16].

Changes in Activities of Daily Living and Mobility

Nurses can obtain additional subjective data by asking the patient how the problem affects activities of daily living and mobility. Changes in normal activities may be from pain alone or from other effects of their illness, including fatigue, weakness, stiffness, or decreased mobility of a particular body part. Some patients may have abandoned activities or made adjustments to maintain independence. Patients should be encouraged to discuss their view of the situation to bring insights and misconceptions to the surface [16].

Assistive Devices

Patients should be asked about any assistive devices used to help maintain independence, including aids for walking, eating, dressing, bathing, or toileting. These may not be devices designed specifically for the tasks; some are creative and adaptive in finding new ways to meet their daily needs [16].

History of the Injury

Subjective data are particularly helpful in the case of injury when the patient can describe the traumatic event and the action taken. This information can help the healthcare team determine what tissues and structures were injured as well as anticipate potential problems.

OBJECTIVE DATA

Physical Assessment

Objective data include the results of physical assessment and of laboratory and other diagnostic tests. When assessing patients with musculoskeletal disorders, vital signs, posture, muscle strength and tone, ability to ambulate, and neurologic status should all be included in the patient assessment [16].

Vital Signs

Assessment of vital signs is of particular importance in cases of musculoskeletal trauma. Hyperthermia may accompany inflammation and is common with an infection. Observing respiration is essential when injury occurs to the face, neck, or chest. Patients with spinal or chest changes may also have abnormal respirations [16].

Inflammation and Swelling

Inflammation is an immune response to infection, physical trauma, or autoimmune reaction. Swelling occurs as inflammatory exudate forms to defend the tissues from the injury. Edema may also be present. Inspection and palpation are used when assessing patients for swelling and inflammation and comparing one extremity to the other for size, warmth, and erythema. A joint will appear swollen when there is an increase in synovial fluid or when blood or purulent material is present in the joint capsule. This swelling is known as effusion. Effusion in the knee is detected by displacing the fluid with an upward stroke along the medial side of the knee and then pressing on the lateral side. The fluid will return and form a bulge (the bulge sign).

It is important to be gentle when assessing inflamed areas because they are usually tender. It is best to start palpating at the distance from the obvious tender area and work toward it, letting patients know when and where they will be touched and reassuring them that the touch will be gentle [16].

Skin Integrity

Injury or disease processes may cause changes in the skin. Discoloration results when trauma to soft tissues causes ecchymosis (bruising). The skin may be broken or torn as result of injury. Describe any lesions completely: include the occasion, length, depth, and appearance of the involved tissue. If there is any drainage, describe the amount, color, type, and odor [17; 18].

Rashes are common in connective tissue disorders. Look for changes in the skin such as discoloration, dryness, scaliness, and lesions. With some types of arthritis, the hair, skin, and nails may show signs of changes. Discoloration, usually redness, may occur in the palms, over joints, and at the distal ends of the toes and fingers. Normal pigmentation may also be altered. Characteristic nodules may be noted when palpating and observing the skin [17; 18].

Structural Changes

Joints may be assessed for changes by observation and palpation. Heberden nodes may be noted on the distal interphalangeal joint of patients with osteoarthritis. Likewise, rheumatoid nodules may be noted near the joints of patients with rheumatoid arthritis, even in the absence of other signs. Joints may be compared bilaterally to assess symmetry, position, and changes in alignment [17; 19].

The curvature of the spine should be assessed to identify the presence of scoliosis (lateral curve), kyphosis (convex curve of the thoracic spine), or lordosis (concave curve of the lumbar spine). Patients with skeletal changes may shift another body part in the opposite direction to compensate for the imbalance; for example, the pelvis may tilt to compensate when one leg is shorter than the other [17; 18].

Range of Motion

Range of motion can be measured with an instrument called a goniometer. Placing the arms of the goniometer parallel to the axis of the bones that form the joint, the examiner measures the angle for the typical positions of the joint. The elbow's normal flexion, for example, is 160°, whereas its normal extension is 0°. To determine what is normal for a patient, compare a joint with an apparently impaired range of motion to the corresponding joint in the other extremity, if possible. Patients can have differences in range of motion for a variety of reasons, particularly as they age, so it is vital to assess typical range of motion on an individualized basis. Dexterity is usually assessed by asking the patient to pick up an object from a flat surface [17; 18].

If a patient is unable to move an extremity, range of motion may be determined through passive movement. Joints should not be moved beyond the point of comfort, and if possible, assessments should not include acutely inflamed joints, which may be tender [17; 18].

During the assessment of range of motion, note joint stiffness, instability, and changes. Bony crepitation may be heard or felt during movement when there is a rough surface of the articular cartilage or when broken bone ends rub together. A limitation of motion may be due to a contracture. Early detection of signs movement limitation can allow for the implementation of measures to improve range of motion and prevent further limitations [17; 18].

Posture

Observe the patient's standing posture for abnormalities. Posture can be affected by structural changes or differences, muscle weakness, or trauma. In addition, patients may hold themselves in positions that relieve or decrease pain. Patients should be observed for symmetry. Posture is also an indication of energy and muscle tone. Normally, posture is erect but not rigid [16].

Muscle Strength, Size, and Tone

Assessment of muscle strength, size, and tone can support the diagnostic process, but it can also provide information about the amount of assistance necessary for ambulation and participating in activities. Muscle strength is assessed by asking the patient to resist movements or to move against resistance [16].

Muscles should be observed and palpated bilaterally to check their size and asymmetry. If there seems to be a significant discrepancy in size, the limb circumferences should be measured [16].

Muscle tone is assessed by moving the extremities passively. While the patient is relaxed the examiner moves the extremity through the ranges of motion, noting resistance to movement. A muscle with diminished tone is described as flaccid. When the muscle is tight and tense from involuntary contraction, it is said to be spastic [16]. Function of the muscles depends on proper function of the nervous system. Muscle abnormalities noted during the assessment may be due to disorders of the nervous system rather than musculoskeletal disorders.

Ability to Ambulate

To assess the ability to ambulate, the patient should be asked to get up to walk across the room, turn around, and come back. Any difficulties rising to standing, starting or stopping walking, or turning should be noted. In a typical gait, the feet are 2–4 inches apart, and the body shifts from side to side about 1 inch. The posture is erect, with toes pointed straight ahead and shoulders in a straight line; the arms swing back and forth at the person's sides, and movement is smooth with good balance [16].

There can be a variety of irregularities in gait, with an equally diverse underlying etiology. A limp can occur from differences of leg length, joint motion, muscle strength, or other causes. The gait may appear stiff, unsteady, or wide-based; the feet may drag, or the steps may be very short. The body may lurch to the side as the individual shifts weight from one leg to the other. An irregular gate can cause fatigue because of the extra energy needed for walking. Ambulation may also be affected by pain, fear of falling, and loss of balance and coordination. As adults age, walking speed and balance may decrease. Steps may be short and shuffling, without the confidence and poise of youth [16].

DIAGNOSTIC STUDIES

Diagnostic studies provide information useful in diagnosing and following the course of the disease process.

Serum Enzyme Tests

Blood tests performed to detect presence of muscle disease measure levels of enzymes released when muscle tissues are destroyed or injured. These enzymes are creatine phosphokinase or creatine kinase, lactic dehydrogenase (LDH), and serum glutamic-oxaloacetic transaminase (SGOT), also known as aspartate amino-transaminase (AST). The same tests indicate cardiac muscle destruction in the patient with a myocardial infarction [20].

Serum Tests for Antibodies and Antigens

The antinuclear antibody (ANA) test is the most specific and sensitive test for lupus and is therefore the most commonly used autoantibody test. Ninety-seven percent of patients with lupus have a positive ANA blood test. The titer and patterns of the blood sample are reported. A titer greater than 1:80 is usually considered positive [21]. It is important to note that a positive ANA test is found in 97% of patients with lupus,

but alone, it does not indicate a conclusive diagnosis of lupus [21]. A positive ANA test, although not always found, satisfies one of the four typical clinical characterizations required for a definitive diagnosis of lupus. ANA tests may also be positive in patients with other connective tissue diseases, chronic infectious diseases, and autoimmune diseases [21].

The 2010 American College of Rheumatology (ACR) and European League Against Rheumatism (EULAR) joint working group recommends several laboratory tests for the diagnosis of rheumatoid arthritis, including rheumatoid factor, erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), and anti-cyclic citrullinated peptide (anti-CCP) antibody [22]. A positive rheumatoid factor is the most specific and sensitive laboratory marker of rheumatoid arthritis, as it is seen in about 70% to 80% of patients [23; 24; 27]. It is also present in many healthy individuals, patients with other rheumatic diseases, and individuals with chronic infections [26]. The anti-CCP antibody test is a specific blood test available for diagnosing rheumatoid arthritis and distinguishing it from other types of arthritis [24; 27]. The anti-CCP antibody test is a marker of anti-citrullinated protein antibody (ACPA) and is positive in about 80% to 90% of patients; it can also be present in other diseases, including active tuberculosis, and is especially useful in early synovitis. While rheumatoid arthritis differs from person to person, individuals with rheumatoid factor, the anti-CCP antibody, or subcutaneous nodules tend to have more severe forms of the disease [24; 26; 27]. However, biomarkers for the initial tissue processes that cause joint damage in rheumatoid arthritis lack prognostic accuracy and are therefore inadequate as stand-alone tests. As such, they are typically used to help rule out other causes of arthritis when a patient has clinical features of rheumatoid arthritis [28].

The presence of human lymphocyte antigen B27 (HLA-B27) is used to help diagnose or rule out ankylosing spondylitis and reactive arthritis. This antigen is present in 90% of those with these conditions, but it can also be found in those without pathology, so it is not diagnostic [20].

Serum Uric Acid

Serum uric acid is elevated during an acute episode of gout but may be normal during remission. Serum uric acid level is also used to assess kidney function [20].

Erythrocyte Sedimentation Rate and C-Reactive Protein

The ESR is a test in which the settling of red blood cells in uncoagulated blood is timed. This is a nonspecific test, and elevations in ESR are indicative of generalized inflammation. Changes in the ESR give an indication of improvement or worsening of the condition [20].

CRP is also associated with disease activity, and the CRP value over time correlates with radiographic progression in patients with rheumatoid arthritis [24; 26; 29]. ESR is typically ≥ 30 mm/hour, and CRP level is typically ≥ 0.7 pg/mL.

Synovial Fluid Analysis

In certain instances, clinicians may perform an arthrocentesis in order to differentiate rheumatoid arthritis from other arthropathies [30]. Findings from synovial fluid aspiration that support a diagnosis of rheumatoid arthritis include straw-colored fluid with a significant number of fibrin flecks, synovial fluid ability to clot at room temperature, and 5,000–25,000 white blood cells/mm³ ($5\text{--}25 \times 10^9/\text{L}$) with 85% polymorphonuclear leukocytes [23; 24]. In addition, bacterial cultures are negative, no crystals are present, and the synovial fluid glucose level is low [23; 24].

X-ray

Examination by x-ray helps diagnose joint problems; it also allows following of the progress of a condition and its response to treatment. X-rays are able to show joint changes, such as erosion of joint margins, joint space narrowing, bone spurs, loose bodies, and dislocation. Specific injuries to soft tissues such as tendons and ligaments do not show on x-rays, but soft tissue swelling may be obvious [20].



For patients with chronic extremity joint pain and suspected rheumatoid arthritis, the American College of Radiology recommends x-ray as the imaging study of choice for evaluation.

(<https://acsearch.acr.org/docs/3097211/> Narrative. Last accessed September 26, 2022.)

Strength of Recommendation: 9 (Usually appropriate)

Magnetic Resonance Imaging (MRI)

Magnetic resonance imaging (MRI) can produce a detailed and highly useful image of the joints and soft tissues. As such, it is usually the best option when evaluating major joints, the spine and the muscles, tendons, and ligaments of the extremities. MRI has a role in the diagnosis of a variety of musculoskeletal disorders, including osteoarthritis, back pain, tears in the connective tissues of the joints, congenital disorders of the joints, and occupational/sports-related injuries [31].

Musculoskeletal Ultrasound

Because it is readily available and avoids the use of radiation, ultrasonography is often a good option in the assessment of musculoskeletal disorders and injuries. Ultrasound allows for the visualization of joints, tendons, muscles, bursae, ligaments, cartilage, nerves, fascia, and related soft tissue and can have a role in diagnosis and/or evaluation of disease progression for a variety of conditions. The American Academy of Physical Medicine and Rehabilitation indicates that ultrasound is an essential component in the diagnosis of tendinopathies/tendon tears, nerve entrapments (e.g., carpal tunnel syndrome), and acute or chronic muscle injury [32]. It may also be involved in the evaluation of ligamentous injury and joint instability

syndromes, subluxations/dislocations, and fascia injury or inflammation. When joint aspiration is necessary, it may be guided by ultrasound, as may therapeutic injections.

Electromyogram

The electromyogram is a test to measure the electric currents produced by muscles, at rest and during contraction. Small needle electrodes are inserted into the muscle being tested and then connected by wires to an electromyography machine. Changes in muscle electrical activity may be helpful in diagnosing neuromuscular disease, and the test is particularly useful in differentiating muscular disease from neurologic disease [20].

Biopsy

Various biopsies may be performed on the musculoskeletal system. Skin samples, obtained by a punch biopsy, may be examined to diagnose certain connective tissue disorders. Muscle biopsies are usually operative procedures done to evaluate muscle disease. The synovial membrane can be biopsied, and analysis can be useful in diagnosing different types of arthritis. Buccal mucosa may be biopsied to help diagnose Sjögren syndrome, and the temporal artery may be biopsied to diagnose temporal arteritis [20].

SPECIFIC DISORDERS OF CONNECTIVE TISSUE AND MUSCLES

Injury to connective tissue and muscle may arise from congenital or acquired disease or from trauma. Diagnosis and treatment/management of these conditions are specific to the disorder.

GENETIC DISORDERS

Genetic disorders of connective tissue are structural connective tissue changes present at birth. Most of these disorders are transmitted by a single autosomal dominant gene. Although there are many congenital connective tissue disorders, most are rare; two more widely known conditions are Marfan syndrome and Ehlers-Danlos syndrome. The obvious manifestations of these disorders may not appear until the second decade of life or later [33].

Both syndromes are serious and require collaborative assessment and treatment by the entire healthcare team. Healthcare providers should gather careful family histories detailing the patterns of disease transmission so families can see the degree of risk [33].

Marfan Syndrome

Marfan syndrome is one of the most common inherited (autosomal dominant) disorders of connective tissue, occurring in 1 in every 10,000 to 20,000 individuals [34]. It is the result of mutations in the *FBN1* gene. *FBN1* mutations are associated

with a broad continuum of physical features ranging from isolated features of Marfan syndrome to a severe and rapidly progressive form in newborns.

Clinical Manifestations

There is wide variability in clinical symptoms in Marfan syndrome, with the most notable occurring in eye, skeleton, connective tissue, and cardiovascular systems. The most common symptom is myopia. Ocular problems are a result of defective supporting tissue of the lens, which can cause bilateral subluxation or total dislocation of the lens. The dislocation is usually upward, but slit-lamp examination is done to detect more subtle variations. Complications such as reduced visual acuity, uveitis, glaucoma, cataracts, and retinal detachment may also occur [33].

Cardiovascular complications of Marfan syndrome are potentially life-threatening and commonly involve the aorta. Marfan syndrome causes degeneration of the elastic fibers of the aortic media, which can lead to dissecting aneurysm. Aortic regurgitating may occur, producing a diastolic murmur. Mitral valve prolapse, thickening of the coronary arteries, conduction system abnormalities, and aortic coarctation have also been associated with this condition [33].

Echocardiogram is useful in following aortic and mitral valve abnormalities. Patients with valve involvement are at risk for endocarditis. These patients should be prescribed antibiotic prophylaxis for any dental work causing bleeding or for any other invasive procedures, to prevent bacteremia [33].

The most obvious skeletal manifestations in patients with Marfan syndrome are extreme height and long extremities. These patients are usually much taller than other members of their families and have excessively long arms and legs in relation to their bodies. The measurement from fingertip to fingertip with the arms outstretched is typically greater than the body height. Arachnodactyly (extremely long fingers) is commonly noted. The sternum may bulge outward (pectus carinatum, or pigeon breast), or it may be depressed (pectus excavatum, or funnel breast). If the chest differences are extreme, the echocardiogram becomes unreliable [33].

Kyphoscoliosis may be quite severe because of the weakness of the ligaments and other supporting connective tissues. Other skeletal manifestations include a long and narrow skull, with a high, arched palate, and flat feet. Joints and ligaments are hyperextensive, leading to recurrent dislocations of the knees and hips [33].

Therapeutic Measures

Therapeutic approaches in Marfan syndrome are directed toward the specific manifestations. Corrective lenses are almost universally necessary, and yearly ophthalmologic examinations aid in early detection of retinal detachment and lens dislocation [33].

Because cardiovascular problems are the major cause of mortality, most diagnostic and treatment efforts are directed here. Echocardiograms are done yearly, unless the diameter of the aorta exceeds the upper limits by 50%, in which case echocardiogram is performed every six months [33]. Beta blockers are used to decrease the stress on the aorta at the time of diagnosis or when there is progressive aortic dilatation. There is some evidence that angiotensin receptor blockers may be used, and clinical trials are underway to evaluate this use. Surgery to repair the aorta is done when the aortic diameter is greater than 5 cm in adults and older children, when the aortic diameter increases by 1.0 cm per year, or when there is progressive aortic regurgitation [34].

Kyphoscoliosis is the most deforming and disabling skeletal manifestation of Marfan syndrome. Patients should be examined biannually, and therapy (e.g., bracing, physical therapy, spinal fusion) should be initiated as soon as possible to prevent or slow further changes [33]. In more severe cases, the thoracic cavity in patients with kyphoscoliosis can be so reduced that cardiac and respiratory function are compromised. These patients are particularly susceptible to upper respiratory infections and should be treated aggressively if an infection occurs [33].

Prepubertal girls are often given estrogen and prepubertal boys given androgens to decrease height and help prevent kyphoscoliosis. While these hormones induce early epiphyseal closure, they also trigger the physical and psychosocial changes of puberty, which can create additional psychosocial stresses.

While Marfan syndrome is not always inherited, it is always heritable. Approximately 75% of cases are inherited, and the offspring of patients with Marfan syndrome have a 50% chance of developing the syndrome. In addition, patients with Marfan syndrome who become pregnant are at risk for potentially dangerous aortic changes resulting from cardiovascular overload and increased intra-abdominal pressure [33].

Specific Nursing Measures

The health history is extremely important in patients with congenital disorders such as Marfan syndrome. Particular attention should be paid to the patient's coping abilities in terms of living with a chronic disease that involves numerous changes in body image [7; 35].

At each visit, the patient should be thoroughly assessed, with particularly attention to the eyes, cardiovascular system, and musculoskeletal system [7; 35]. When examining the patient's eyes, look for tremor of the iris as it is moved horizontally. This is an indication of subluxation of the lenses. These patients may also have myopia and blue sclera (due to the presence of thin sclera through which the vessel-rich choroid can be seen).

Patients may display early diastolic murmurs of aortic regurgitation. This consists of a high-pitched blowing sound, heard best with the stethoscope over the second right or third left intercostal space. Increased pulse pressure and collapsing

(water-hammer) pulse may also be evident. Occasionally, a midsystolic click indicative of mitral valve prolapse may be auscultated.

Nursing interventions for these patients will focus on supportive symptomatic care and education needs. The nurse should be prepared to discuss the nature and course of the disease and the importance of genetic and pregnancy counseling. The patient should be urged to keep current with biannual exams. Patients should also be counseled to avoid trauma, including contact sports, and invasive surgical procedures (when possible) [7; 35]. They are also advised to avoid medications and foods that can lead to chronic increases in blood pressure and stretch the connective tissue in the cardiovascular system.

Ehlers-Danlos Syndromes

Ehlers-Danlos syndromes are a group of rare genetic disorders of connective tissue that affect the skin, joint, and hematopoietic systems. It is usually transmitted by an autosomal dominant gene, but it may also be recessive or an X-linked recessive gene [33].

Clinical Manifestations

The major manifestations of Ehlers-Danlos syndromes are fragile and increased elasticity of skin, hyperextensible joints, and fragility of blood vessel walls [33]. In the 2017 classification system, 13 types of Ehlers-Danlos syndrome were identified, including rarer forms [36]. They are generally organized according to the dominant system(s) involved, severity, and mode of transmission.

The skin of most patients with an Ehlers-Danlos syndrome is very smooth and hyperextensible; it can be pulled away from the body but returns to its original shape. Fragility and bruising are often evident. Minor cuts cause gaping wounds with little bleeding. Even the slightest trauma may cause purpura or hematomas that calcify, particularly over pressure points such as knees and elbows [33].

An unusually large range of joint movement (hypermobility) occurs in most forms of Ehlers-Danlos syndrome, and it is a hallmark feature of the hypermobile type. Dislocations, effusion, and hemarthrosis of the hip, patella, and shoulders may occur. Kyphoscoliosis, flat feet, and hyperextensible knees are often present. Thoracic changes are not as common but do sometimes occur, as does a forward slipping of the lower lumbar vertebrae (spondylolisthesis) [33].

The patient may have episodes of bleeding, including spontaneous epistaxis; bleeding into the joints (hemarthrosis); blood in the sputum (hemoptysis); dark, tarry stools (melena) indicating bleeding in the digestive tract; and bleeding gums. It is not known whether the abnormal bleeding is from weakness in blood vessel walls or abnormal interactions of platelets with collagen [33]. Patients with Ehlers-Danlos syndrome who become pregnant are at risk for uterine rupture.

Abnormalities of the heart and blood vessels occur in patients with the cardiac-valvular type. These include mitral valve prolapse, right bundle branch block, and other conduction abnormalities. Patients with this type of Ehlers-Danlos syndrome have friable arteries, increasing the risk for adverse events during invasive angiography [33].

Other manifestations of Ehlers-Danlos syndrome can include spontaneous bowel rupture, pneumothorax, and diaphragmatic hernias or diverticula. In rare instances, a patient may have glaucoma, retinal detachment, or corneal abnormalities [33].

Specific Nursing Measures

Care for patients with an Ehlers-Danlos syndrome is limited to symptomatic treatment and support; there is no curative treatment. The main concern is to protect the patient's skin and joints from cuts, bruises, and dislocations. At each visit, the patient should be assessed for bleeding gums, melena, hemoptysis, and nosebleeds. Inadequate wound healing or wound dehiscence after a surgical procedure should be noted. Assessment of the lungs for pneumothorax, particularly following surgery, is important [35].

As with any chronic condition, the nurse needs to teach patients and their families about the nature and course of the disease. The patient should also be referred to a genetic counselor, as there are varying modes of heritability. A patient with Ehlers-Danlos syndrome who becomes pregnant is at risk for abortion, preterm birth, exacerbation of joint problems, increased bruisability, abdominal hernia, and varicosities. Serious complications may arise with cesarean deliveries, because sutures do not hold well and wound dehiscence may result [7].

INFLAMMATORY DISORDERS

Many pathologic conditions involve inflammation of connective tissue. In this section, most of the inflammatory conditions are related to alterations in the immune system [37].

Bursitis, Tendinitis, and Tendinosis

Bursitis is an inflammation of the synovial membrane lining a bursa; tendinitis is an inflammation of a tendon. These inflammations may result from trauma, or they may be secondary to disease. Although both conditions are usually acute, they can become chronic and disabling with repeated injury or inadequate care [37]. Note that tendinitis is distinct from tendinosis, which is the result of a noninflammatory condition characterized by degeneration of the tendon in response to chronic overuse.

Bursitis and tendinosis develop from prolonged overuse of a particular muscle group that can eventually damage a bursa or tendon. Overuse may be due to repetitive work movements or to a sports activity. Because the vascular supply of tendons is poor, their healing is limited and inflammation can become chronic, resulting in tissue damage and persistent pain. Often,

the patient becomes unable to continue performing the movements that led to the condition, potentially impairing their ability to continue working.

Calcium deposits in tendons or bursae may also be the cause of inflammation. Tendon sheaths may become inflamed secondarily to systemic disease, such as gout, rheumatoid arthritis, or scleroderma [37].

Clinical Manifestations

The major symptom of bursitis/tendinitis/tendinosis is pain, often so severe that the patient is unwilling to move the affected part. Swelling may be present, and this alone may keep the patient from moving the joint. Any of the body's many bursae and tendons can become inflamed, but some joint areas are more commonly affected than others. Differential diagnosis of acute pain and erythema in joint areas should include infection, gout, and rheumatoid arthritis [37].

Bursitis and tendinitis/tendinosis of the shoulder involve the subacromial and subdeltoid bursa (different sections of the same large bursa) and the tendon of the supraspinatus muscle. The onset of bursitis or tendinitis in the shoulder usually follows activities involving repetitive movements of the whole arm, such as sanding, painting, sawing, throwing, or repeated lifting. Pain in the deltoid area increases when the patient lies on the shoulder or actively abducts the arm. A classic sign of bursitis/tendinitis/tendinosis of the shoulder is the "painful arc" between 80° and 120° of active arm abduction. The patient is often unable to support the weight of the arm at these angles. Further abduction causes no pain, and the examiner can perform assisted range of motion. If passive range of motion causes pain, capsulitis, rather than a periarticular disorder, is suspected [37].

Inflammation of the elbow region most often involves the olecranon bursa and the medial and lateral epicondyles. "Tennis elbow" is generally lateral epicondylitis, and "pitcher's elbow" is medial epicondylitis. These conditions cause pain that radiates from the elbow down to the forearm. The patient may drop heavy objects because of a feeling of decreased strength, although there is no real loss of strength or range of motion. Palpation of the involved epicondyle causes pain. Activities involving lower arm movement, such as tennis or hammering, may precipitate an attack. Olecranon bursitis usually is caused by leaning or falling on the elbow. There may not be severe pain, but swelling is often extensive [37].

Tenosynovitis involves inflammation of the tendon and tendon sheath and is also known as de Quervain tenosynovitis of the wrist [38]. When the tendons at the base of the thumb become irritated or inflamed this causes the tunnel around the tendon to swell and results in pain and difficulty grasping and holding objects. Overuse is the most common cause [38]. New repetitive activity, hormonal fluctuations associated with pregnancy and breastfeeding, and wrist fractures also are possible causes of de Quervain tenosynovitis [39].

Stenosing tenosynovitis, also referred to as “trigger finger,” occurs when the pulley/tendon relationship between the hand and fingers is restricted by thickening or swelling at the base of the fingers. This creates pain and a distinctive catching, popping, or locking action in the finger or thumb. A cycle of triggering, inflammation, and swelling is common. Like carpal tunnel syndrome, stenosing tenosynovitis has been associated with other health conditions, such as gout, diabetes, and rheumatoid arthritis. In many cases, the actual cause is not clear [40].

The most common inflammatory problem of the hip is trochanteric bursitis. Pain, which is distributed over the lateral aspect of the hip and thigh, may inhibit ambulation. An increase in pain is seen with abduction and internal rotation against resistance. The patient feels tenderness with palpation over the greater trochanter. Patients who have leg length discrepancy may develop this inflammation in the hip of the longer leg [37].

Four bursa in the knee can cause significant discomfort for the patient when inflamed [37]:

- Prepatellar bursa
- Superficial infrapatellar bursa
- Deep infrapatellar bursa
- Pes anserine bursa

Prepatellar bursitis (“housemaid’s knee”) results from the combined action of excessive kneeling and leaning forward, as when gardening. Superficial infrapatellar bursitis (“clergymen’s knee”) can result from excessive kneeling. Deep infrapatellar bursitis and pes anserine bursitis are secondary to excessive weight bearing or unusually strenuous exercise [37].

Achilles tendinitis is a painful inflammation of the tendon of the ankle with or without swelling. This injury often results from a single episode of overuse. It can also occur in runners who wear shoes with rigid soles. Recurrent episodes of Achilles tendinitis, when a patient resumes activity before complete healing has occurred, can result in progressive scar formation, which may require surgical repair [37].

Therapeutic Measures

The measures employed for relief of bursitis and tendinitis vary according to the patient’s age and the location, cause, and severity of the injuries. Recommendations usually include [37]:

- Short-term immobilization, particularly during differential diagnosis
- Ice packs applied to the affected area
- Physical therapy and structured exercise after the initial period of rest
- Anti-inflammatory medication

Occasionally, local corticosteroid injections are administered to the inflamed bursa or tendon area. While this approach is relatively widespread, it is not supported by well-designed systematic reviews [41].

Physical therapy and increasing return to activities is the best practice for these patients. Physical therapy consists of a four-step approach [42]:

1. Pain reduction and load management (isometric loading and avoiding positions of compression)
2. Isotonic loading (heavy-slow resistance through concentric-eccentric phases)
3. Energy-storage loading (plyometric loading)
4. Return to activity/sport

Exercise is crucial in the rehabilitation process, and active movement is started early. For example, in bursitis of any bursa of the knee, quadriceps-setting exercise is begun as soon as pain allows. When pain and tenderness have completely subsided, range of motion and full quadriceps activity are initiated. Physical therapists are often involved in designing and implementing exercises for patients, according to their individual needs. Occupational therapists may also participate if the nature of the problem involves a modification or change in job [15].

In some cases, fluid may be aspirated from the bursal space to relieve the symptoms. Any fluid obtained should be cultured and inspected. X-rays of joints are usually normal, but in some instances, calcium deposits can be identified as the precipitating factor. Arthrography is indicated in specific types of shoulder trauma to rule out any disruption of the joint capsule. Surgery is rarely used for bursitis or tendinitis unless rupture of the tendon occurs [37].

Specific Nursing Measures

Goals of nursing care are to relieve the patient’s pain, maintain maximum mobility, and prevent joint contracture. Assessment of pain and range of motion is important both initially and after treatment to measure improvements. Reassurance and support can contribute to the relief of pain, so it is helpful to assure the patient that the pain of bursitis or tendinitis/tendinosis is usually of short duration [35].

Instrumental to the success of treatment is comprehensive patient education. Patients should receive instruction on physical therapy exercises (including frequency), pain management techniques, and return to activities; written instruction should also be provided. If pain is relieved with pharmacotherapy, the patient may be tempted to use the affected area too soon. It is important to caution patients to refrain from early resumption of activity to avoid reinjury and/or the creation of scar tissue [35].

Polymyalgia Rheumatica

Polymyalgia rheumatica is an immune-mediated inflammatory disorder characterized by muscle stiffness, pain, and weakness around the neck, shoulders, and hip. While this is an inflammatory disorder, the cause of trigger is unclear; genetic, infectious (e.g., Epstein-Barr virus, parvovirus), and gut health-related etiologies have all been suggested, with varying levels of evidence [43]. The incidence increases with age, with the greatest incidence in White patients older than 50 years of age; the average age at diagnosis is 70 years.

Clinical Manifestations

As noted, the characteristic symptoms of polymyalgia rheumatica are pain and stiffness in the shoulders, neck, upper arms, and hip area. The pain and stiffness are usually worse upon waking in the morning or after resting, and usually last an hour or more. Patient may experience difficulty performing normal activities, including rising from bed or a chair, dressing, and brushing hair. Many patients will have difficulty raising their arms above the shoulders [44]. Less common signs and symptoms include flu-like symptoms (e.g., low-grade fever, weakness, loss of appetite, weight loss) and swelling of the wrists or joints in the hands. Onset of symptoms is typically over the duration of a few days but may be as short as overnight.

Diagnosis is typically based on the presence of elevated inflammatory markers, particularly ESR and immunoglobulin G (IgG). In addition, these patients will display a decreased number of circulating B cells compared with healthy adults [43].

A significant portion of patients with polymyalgia rheumatica are also diagnosed with giant cell arteritis, and research indicates the co-occurrence of these conditions is common even without the presence of symptoms [43].

Therapeutic Measures

The EULAR and the ACR have issued a joint guideline for the management of polymyalgia rheumatica [45]. The cornerstone of treatment is at least 12 months of glucocorticoid therapy. This typically consists of 12.5–25 mg prednisone, although a lower dose may be preferred in patients at risk for glucocorticoid-related adverse events (e.g., those with osteoporosis, glaucoma, diabetes). Drug therapy should be tapered up to effective dose and tapered down when discontinued.

Care of patients with polymyalgia rheumatica includes monitoring for and preventing (when possible) the adverse effects of long-term steroid therapy. This can include vitamin D and calcium supplementation as well as bisphosphonate prophylaxis for those at increased risk for fracture [43]. Because close monitoring is necessary, patient education should include the necessity for keeping all follow-up appointments.

IMMUNOLOGIC DISORDERS

The disorders in this section are believed to have an autoimmune etiology. As with many autoimmune disorders, there are a variety of potential initiating factors, including viral infections, genetic predisposition, and exposure to toxins [46].

Autoimmune disorders may be generally classified as organ-specific or generalized. Autoimmune connective tissue diseases are generalized, usually involving a progressive degradation of collagen in connective tissue throughout the body. Rheumatoid arthritis is among autoimmune disorders but will be discussed later in this course, because joint involvement is the major problem [46]. Some autoimmune disorders result in musculoskeletal manifestations but have an etiology in another body system. For example, fibromyalgia is characterized by widespread musculoskeletal pain and fatigue, but it is believed to be the result of nervous system dysfunction.

Autoimmune connective tissue disorders can be associated with significant morbidity mortality. However, early diagnosis and treatment have improved prognosis, though they remain chronic (incurable) conditions. Successful therapy for patients with autoimmune disease requires an interprofessional team approach in order to ensure the best outcomes for patients [46].

Familiarity with each disorder will prepare the nurse to be alert for manifestations, exacerbations, and patient education needs. Because patients are often prescribed several medications to help manage the disorder, nursing management often includes medication management. Comfort measures are another important aspect of nursing care during acute phases or exacerbations. Proper positioning, use of splints, and small comfort measures (e.g., backrubs, smoothing wrinkled sheets, creating a calm environment) all contribute to the patient's well-being [46].

The nurse will explain to patients how they can try to prevent exacerbations of specific manifestations of their disease and how to cope with them when they do occur. Prevention measures may include avoiding stress, cold, sun, or certain drugs [46].

Systemic Lupus Erythematosus

Four different forms of lupus have been identified: cutaneous lupus erythematosus, drug-induced lupus, neonatal lupus, and systemic lupus erythematosus (SLE) [47]. Cutaneous lupus mainly affects the skin. It is associated with chronic skin eruptions that, if left untreated, can lead to scarring and permanent disfigurement. Drug-induced lupus is associated with ingestion of various drugs that result in lupus-like symptoms. Neonatal lupus is a rare, non-systemic condition affecting infants of women with lupus. SLE, which affects multiple organ systems as well as the skin, is considered the most common of the four forms.

SLE, often referred to simply as lupus, is a chronic inflammatory autoimmune disorder of the connective tissue, primarily affecting the skin, joints, blood, and kidneys [47; 48; 49]. In this autoimmune disorder, antibodies are formed within the body that target healthy body systems, causing inflammation and structural changes. The word lupus means “wolf” in Latin, while erythematosus means “redness.” The disease is named for the characteristic red rash that appears on the face and is thought to resemble a wolf’s face [47; 49]. The term “lupus erythematosus” was coined in 1851 by Pierre Cazenave, a French dermatologist, but writings describing lupus date to ancient Greece [49; 50].

Lupus has been characterized as a multidimensional, unique, complex, challenging, unpredictable, and often elusive disease [47]. It is a non-organ-specific systemic disease with a varying prognosis that can be mild, serious, life-threatening, or even fatal. The disease is characterized by recurring remissions and exacerbations, often called flares, that occur most commonly in the spring and summer [48; 51]. Periods of remission vary considerably among those diagnosed with lupus [47].

The number of reported cases of lupus varies based on different sources; it is believed that there are at least 1.5 million affected individuals in the United States [52; 53]. More than 90% of SLE cases occur in women, with most women developing symptoms in their childbearing years (15 to 45 years of age) [54]. New diagnoses of lupus in women older than 45 years of age are uncommon [49]. SLE is most common among African Americans, with African American women having three times the incidence of White American women [54]. The incidence of lupus is also greater in Hispanic, Asian, and Native American women when compared to White women [55]. Statistics show that Black and Hispanic women tend to develop the disease at a younger age, are more likely to develop more serious complications (particularly cardiovascular complications and kidney disease), and tend to have a higher mortality rate from the disease as compared to White women [54].

The exact cause of lupus remains a mystery, but researchers believe that it results from multiple factors [49; 56]. Possible causes may be interrelated and include immunologic dysfunction, genetic factors, hormones, and environmental influences [50; 51].

Immune dysregulation, in the form of autoimmunity, is thought to be the prime cause of lupus. In patients with lupus, the body produces an accelerated inflammatory response, resulting in the production of autoantibodies, causing immune complexes (antigens combined with antibodies) [49; 56]. These autoantibodies and complexes assault the body’s own healthy cells and tissues [47; 49; 50; 51]. Symptoms of SLE are the result of the damage to the body’s tissues secondary to the immunologic response. One of the hallmark indicators of lupus is the formation of autoantibodies, and the presence of autoantibodies in the blood is a key factor to the diagnosis of lupus [47; 49; 51].

The strong hereditary component of lupus is supported by the fact that first- and second-degree relatives of patients with lupus are at a greater risk for developing lupus [57]. Estimates indicate that 5% to 13% of relatives will develop lupus, but only 5% of children whose mothers had lupus will develop the disease [57]. For those with a genetic predisposition, environmental factors may trigger lupus [47]. Environmental factors that may precipitate or exacerbate lupus include physical or emotional stress, streptococcal or viral infections, exposure to sunlight, immunizations (live vaccines), surgery, smoking, chemical agents (drugs, metals, or toxins), certain foods or supplements, and other environmental irritants [47; 50; 58]. Further, female sex hormones are believed to have a potential role, as women in their reproductive years are most susceptible to lupus.

Diagnosis

The diagnosis of lupus may be a challenge for the healthcare provider as well as the patient. In 2019, the EULAR and the ACR published updated classification criteria for lupus (**Table 1**) [59]. The EULAR/ACR criteria classifies a person as having lupus if they meet entry criterion of an ANA titer of >1:80, followed by additive weighted criteria (seven clinical and three immunologic) in which the patient must meet one clinical criterion and ≥ 10 points between the clinical criteria and immunologic criteria [59].

Clinical Manifestations

No two people with lupus will experience identical symptoms. The onset of lupus may be acute or insidious, vague, or even nonspecific. On average, individuals with lupus have symptoms of the disease for two to three years before a diagnosis is made [49]. Symptoms are the result of the inflammatory and immune response of the individual’s body to the disease process [49]. Repetitive cycles of exacerbations and remissions of symptoms are a hallmark of the lupus disease process.

Common symptoms of lupus include fever, weight loss, malaise, fatigue, skin rashes, polyarthralgia, vasculitis, Raynaud syndrome (discussed in detail later in this course), patchy alopecia (hair loss), and painless ulcers of the mucous membranes [51]. Fatigue is probably the most universal symptom, described as a persistent complaint of a paralyzing fatigue that normal rest may not relieve [47]. Vague symptoms of lupus include aching, fatigue, low-grade or spiking fever, chills, and malaise. Episodic fever is reported by more than 80% of all patients with lupus, with a low-grade fever most often noted [47]. Infection is certainly a major concern and is a potential symptom for patients with lupus. Those with lupus are more susceptible to opportunistic infections due to alterations in their hematologic system, especially in white blood cells. Women with lupus may also experience irregular periods or amenorrhea due to the disease process [47; 49].

CLASSIFICATION CRITERIA FOR THE DIAGNOSIS OF SYSTEMIC LUPUS ERYTHEMATOSUS		
Domain	Criteria	Weight
Entry Criterion		
Positive antinuclear antibody (ANA) titer	ANA titer of >1.80 on Hep-2 cells or an equivalent positive test (ever)	Must be positive to continue to additive criteria
Additive Criteria, Clinical		
Constitutional	Fever	2
Hematologic	Leukopenia	3
	Thrombocytopenia	4
	Autoimmune hemolysis	4
Neuropsychiatric	Delirium	2
	Psychosis	3
	Seizure	5
Mucocutaneous	Non-scarring alopecia	2
	Oral ulcers	2
	Subacute cutaneous OR discoid lupus	4
	Acute cutaneous lupus	6
Serosal	Pleural or pericardial effusion	5
	Acute pericarditis	6
Musculoskeletal	Joint involvement	6
Renal	Proteinuria >0.5 g/24h	4
	Renal biopsy Class II or V lupus nephritis	8
	Renal biopsy Class III or IV lupus nephritis	10
Additive Criteria, Immunology		
Antiphospholipid antibodies	Anti-cardiolipin antibodies	2
	OR	
	Anti-β2GPI antibodies	
	OR	
Complement proteins	Lupus anticoagulant	3
	Low C3 OR low C4	
	Low C3 AND low C4	
SLE-specific antibodies	Anti-dsDNA antibody	6
	OR	
	Anti-Smith antibody	

Source: [59]

Table 1

Skin rashes are very common among patients with lupus; approximately 80% of patients report skin involvement [47]. A red, raised rash over the nose and cheeks characterizes the classic “butterfly rash” of lupus. The butterfly rash is reported by 55% to 85% of all patients with lupus at some point during their disease process [47]. Discoid lupus lesions may also be seen. Ultraviolet light often aggravates skin eruptions, and approximately one-third of all patients with lupus are found to be photosensitive [47; 60]. Oral, nasal, and vaginal ulcers may occur. Conditions such as alopecia, pruritus, alteration in wound healing, and bruising are other common dermatologic symptoms.

Polyarthralgia (pain in multiple joints) occurs in more than 90% of lupus cases [47]. The joint pain associated with lupus is similar to that experienced by rheumatoid arthritis patients and is often called lupus arthritis. Most patients complain of

morning joint stiffness and pain. The pain is typically symmetrical, and joints may become tender, warm to the touch, and swollen. The dominant extremities are usually more inflamed. Joints commonly affected include the toes, ankles, fingers, wrists, elbows, and knees [61]. Joint pain is often one of the first and most common complaints of those with lupus and is often what initially brings them to a healthcare provider [50]. Additional musculoskeletal symptoms that may occur include subcutaneous nodules, tendonitis, tendon rupture, and carpal tunnel syndrome [47].

Anemia and cardiopulmonary abnormalities are relatively common among patients with SLE, affecting 50% of patients [47; 49; 62]. The most common cardiac complication of lupus is pericarditis, while pleurisy is the most common respiratory complication [47; 49].

Nervous system involvement secondary to lupus is common and can range from mild to severe. Central nervous system involvement may result in cognitive disorders, including confusion, fatigue, memory impairment, and difficulty in articulating thoughts [49]. Cognitive dysfunction is estimated to occur in up to 90% of patients with lupus and is not associated with lupus disease activity [63].

Renal damage is one of the most serious complications of lupus, often causing such symptoms as hematuria, proteinuria, urine sediment, cellular casts, urinary tract infections, and fluid/electrolyte imbalance. Renal involvement has the potential to cause renal failure, affecting up to 50% of patients [47]. Renal disease is a leading cause of death in patients with lupus [47].

Ophthalmic disease affects approximately 20% of patients with lupus [47]. Ophthalmic symptoms associated with lupus may include a lupus rash on the eyelids, conjunctivitis, dry eyes, glaucoma, and cataracts [47]. In severe cases, retinal exudates or blindness may occur.

Therapeutic Measures

There is currently no cure for lupus, and long-term disease management is required. Due to the variability of lupus symptoms, treatment protocols differ for each individual. The range of treatments, however, are increasing in number and becoming more effective; thus, the disease can be controlled reasonably well in most people. The ultimate goal of treatment is to suppress immune system abnormalities, prevent disease flares, and reduce inflammation and other complications secondary to lupus [51].

Treatment is based on such factors as symptoms and severity, overall general health, activity level, school and/or family schedule, age, family and social situations, other medical conditions, and financial and insurance considerations [50].

Although there is no cure for lupus, there are several types of drugs available to aid in the treatment and management of secondary symptoms. Among these drug classes are non-steroidal anti-inflammatory drugs (NSAIDs), corticosteroids, antimalarials, biologics, and immunosuppressives. In cases of severe lupus kidney disease not helped by pharmacologic intervention, dialysis or kidney transplant may be necessary.

Specific Nursing Measures

Nurses may see patients with SLE in both inpatient and ambulatory care settings. Discovering early symptoms and signs of exacerbations and complications is important in prolonging the life of patients with SLE. Carefully monitor all diagnostic study reports to remain well informed about the patient's progress [46].

Individuals diagnosed with lupus are encouraged to do all of the following [47; 49; 50; 51]:

- Get plenty of physical and emotional rest.
- Maintain a healthy diet.

- Establish an exercise regimen.
- Avoid sunlight.
- Seek prompt treatment of infection.
- Limit stress.
- Set realistic goals and priorities.
- Maintain effective communication with their healthcare providers.
- Develop a support system, including family, friends, healthcare professionals, community organizations, and organized support groups.
- Avoid triggering or aggravating factors.
- Seek regular health care.

Eight to 10 hours of sleep per night along with naps are recommended for patients with lupus. In addition, individuals with lupus should minimize stress to reduce emotional distress, as well as avoid direct prolonged sunlight, especially during the hours between 10 a.m. and 4 p.m. The use of a sunscreen with a sun protective factor (SPF) of 15 or greater that protects against both ultraviolet A and B rays is recommended along with protective clothing such as long sleeves and a hat [47]. Routine exercise is important to reduce fatigue and maintain joint mobility.

Social support can have a positive impact on individuals diagnosed with lupus. However, seeking and gaining social support can be difficult when one is experiencing a chronic illness such as lupus, because tremendous energy is necessary to maintain social networks [64]. Lupus symptoms, as well as treatment side effects, can present a challenge for individuals in maintenance of their pre-illness social relationships and activities. Furthermore, to gain necessary support, individuals with lupus should understand and then communicate to others what they need to assist them in managing their disease.

Keller noted similar findings in her research on social support and psychological distress in women with lupus. She concluded that “younger women with lupus were more psychologically distressed than older women with lupus and that women with shorter duration since diagnosis were more distressed” [65]. Keller also found that the perception of having social support and being satisfied with the social support were more important than the number of social supports [65]. Thus, perception of and satisfaction with social support has been found to reduce psychological distress.

One important potential source of assistance can be support groups. It has been noted that “participating in a support group can provide emotional assistance, boost self-esteem and morale, and help to develop or improve coping skills” [51]. Successful support groups can assist patients to gain insights into how to live with their lupus [66]. Darner found that women with lupus who had been diagnosed for longer periods of time had a healthier psychosocial adjustment [67]. Therefore, those newly diagnosed with lupus may require more support and interventions to aid in psychosocial adjustment.

Systemic Scleroderma

Systemic scleroderma, also called sclerosis, is an autoimmune connective tissue disorder that causes fibrous changes in the skin, synovium, and small arteries of the digits, as well as in various internal organs, most notably the esophagus, intestines, heart, lungs, kidneys, and thyroid. The disease occurs in various forms, ranging from a primarily skin condition (localized scleroderma) to the CREST (calcinosis, Raynaud phenomenon, esophageal dysmotility, sclerodactyly, and telangiectasia) syndrome, which is thought to be more benign, to involvement of visceral organs (systemic scleroderma). Some patients with mild-to-moderate types of scleroderma can progress to the visceral and more extensive cutaneous lesions associated with systemic scleroderma [46].

In all forms of the disease, there is vascular injury at the level of small arteries and capillaries, and the resulting decrease in circulation is the cause of the tissue changes. The precipitating factor for the onset of systemic scleroderma is not clear, although there is some evidence that genetic and environmental factors play a role. Silica and certain organic solvents are recognized as risk factors of occurrence of systemic scleroderma. In addition, the prevalence of the disease is 13 times higher in first-degree relatives of patients than in the general population [68]. The result is an activation of the immune system, causing blood vessel damage and injury to tissues that result in scar tissue formation and the accumulation of excess collagen.

There are no definitive tests to diagnose systemic scleroderma, and diagnosis is primarily based on clinical evaluation. Autoantibodies occur in this disorder, and the ESR may be elevated. As part of the diagnostic workup, the following tests may be performed [68]:

- Nailfold capillaroscopy
- Screening for antinuclear antibodies (mainly anti-centromere and anti-scl70/anti-topoisomerase antibodies)
- Transthoracic echocardiography
- High-resolution computed tomography (CT) of the chest
- Diffusing capacity of the lung for carbon monoxide and spirometry
- Hand x-ray
- Esophageal manometry

Clinical Manifestations

The most frequent presentation in systemic scleroderma is the clinical triad of skin changes, Raynaud phenomenon, and esophageal hypomotility. However, manifestations are often present in other organ systems, requiring continual monitoring [46].

The most typical changes in all types of scleroderma occur in the skin. Typically, skin changes begin with swelling of the hands and gradual thickening, tightening, and hardening of the skin of the fingers (sclerodactyly). The fingers become tapered and in severe cases claw-like, with impaired mobility. Ulcers may develop on fingertips and over knuckles as the skin becomes taught. Skin changes can progress proximally at a slow rate, eventually affecting the face. In these cases, the skin of the face becomes tight and shiny, with a loss of normal wrinkles and skin folds. The nose may become beaked, and sometimes radial furrowing is seen around the mouth. Patients may experience an impaired ability to fully open their mouths. In extreme cases, the face becomes expressionless [46].

Most patients with systemic scleroderma have Raynaud syndrome, and this is often the first symptom to appear. With Raynaud syndrome, there is diminished blood flow to the digits secondary to vasoconstriction of the digital arteries, typically triggered by cold, vasoconstriction drugs, or emotional states. The initial sign is digital pallor, which progresses to cyanosis, then to erythema on rewarming [46].

The patient may have pain and stiffness in both small and large peripheral joints. Occasionally, patients develop arthritis and synovial effusion. Contracture and atrophy of the fingers may eventually occur [46].

Hypomotility of the esophagus occurs in most patients with systemic scleroderma. This typically presents as gastroesophageal reflux, with resulting heartburn and stricture, and potentially difficulty swallowing. In some cases, patients require esophageal dilation. Gastrointestinal involvement can progress to the intestine and colon, with development of hypomotility of the small intestine and wide-mouth diverticula [46]. In patients with gastrointestinal involvement, impaired nutrition is common.

Systemic scleroderma can also cause cardiopulmonary problems. Dyspnea may develop as a result of pulmonary hypertension and interstitial fibrosis. The examiner may hear fine dry rales or crackles at the bases of the lungs, and spirometry is often abnormal. Manifestations involving the heart are primarily the result of lung complications, but dysrhythmias, conduction disturbances, pericarditis, and pericardial effusions uncommonly occur [46].

In some patients, the kidneys can be seriously affected, with malignant hypertension rapidly producing renal failure, the leading cause of death for these patients. High renin levels and proteinuria are signs of kidney involvement [46].

Hematologic problems, in addition to a mild normochromic, normocytic anemia, include vitamin B₁₂/folic acid deficiency anemia, which may occur secondary to bacterial overgrowth in an atonic small intestine. There is also a risk for gastrointestinal bleeding and resultant iron-deficiency anemia [46]. Other manifestations include thyroid disease, biliary cirrhosis, trigeminal sensory neuropathy, and Sjögren syndrome [46].

Therapeutic Measures

Treatment of systemic scleroderma is symptomatic and driven by the stage and organ involvement of the disease. In its 2017 guideline for the treatment of systemic scleroderma, the EULAR has established guidelines for the management of manifestations, organized by affected body system [69]. For patients with systemic scleroderma-associated Raynaud phenomenon, evidence supports nifedipine to reduce the frequency and severity of attacks. As such, oral nifedipine should be considered as first-line therapy. Phosphodiesterase-5 (PDE-5) inhibitors should also be considered [69]. For patients with severe disease who do not improve on oral therapy, intravenous iloprost is the recommended approach.

Intravenous iloprost is also recommended for patients with systemic scleroderma who experience digital ulcers [69]. PDE-5 inhibitors have been proven to expedite healing and prevent the development of digital ulcers and should be considered for these patients. Patients who do not respond to calcium channel blockers, PDE-5 inhibitors, or iloprost therapy, may be prescribed bosentan, which has been shown to reduce the number of new digital ulcers in patients with systemic scleroderma. Physical therapy for the hands is important to prevent contractures. For patients with Raynaud phenomenon, biofeedback is sometimes useful for controlling temperature in the hands and feet [46].

For patients whose systemic scleroderma is characterized by pulmonary arterial hypertension, EULAR recommends treatment with endothelin receptor antagonists (e.g., ambrisentan, bosentan, macitentan), PDE-5 inhibitors (e.g., sildenafil, tadalafil), or riociguat [69]. In cases of severe disease, intravenous epoprostenol is the first-line option. In cases of malabsorption by the small intestine, absorption often improves with the use of tetracycline, which destroys the bacterial overgrowth that occurs with hypomotility [46].

Hypertension is treated aggressively with angiotensin-converting enzyme (ACE) inhibitors to prevent irreversible renal damage [46]. The risk for scleroderma renal crisis is increased in patients taking glucocorticoids, and these patients should be closely monitored [69].

Arthritis responds to NSAIDs, and the dry eyes (sicca syndrome) of Sjögren syndrome are helped by artificial tears. A patient with dry mouth (xerostomia) should have frequent dental exams, because this condition predisposes patients to severe dental caries [46].

Specific Nursing Measures

Patients with known or suspected systemic scleroderma should be thoroughly assessed, including the skin, joints, and cardiovascular, pulmonary, and gastrointestinal status. The eyes and mouth should be evaluated for adequate lacrimal and salivary gland secretions. It is important to closely monitor

blood pressure and review laboratory results. Venipuncture in the antecubital area may be difficult because of skin changes; further, finger sticks should be avoided. If only a small amount of blood must be drawn, the earlobe may be the best site [35].

Patient education should include a clear explanation of the nature and course of systemic scleroderma, including signs of more serious involvement. For some patients, demonstration of range-of-motion exercises to prevent joint contracture may be warranted. Patients should be encouraged to use moisturizing lotions to decrease dryness [35].

Patients with Raynaud phenomenon are advised to avoid cold, ergotamine, and amphetamines. They should be cautioned to take precautions against cold weather, including the use of warm gloves and socks. The use of nicotine should be avoided, as it is associated with pronounced peripheral vasoconstriction, which markedly aggravates Raynaud syndrome [35].

Patients with esophageal dysmotility should be advised to eat small, frequent meals and to chew their food thoroughly; meals should be followed with water. Proton pump inhibitors (PPIs) and antacids after meals and at bedtime can help to help to relieve gastroesophageal reflux disease. Resting and sleeping with the head of the bed elevated may also help to relieve symptoms [35].

The face and hands often undergo considerable changes in scleroderma, which alters the patient's appearance and manual dexterity. The facial skin becomes taut, the nose may become beaked, and telangiectasias may appear on the face. Tapering of the fingers, with tightness of the overlying skin, occurs as flexion contractures may be present [35]. These physical changes may cause varying levels of disability, but they can also have a negative effect on the patient's self-esteem and self-worth. Referral to mental health care and participation in support groups can be helpful.

TRAUMATIC DISORDERS

Sprains and Strains

Traumatic injuries to the soft tissues surrounding joints—muscles, ligaments, and tendons—are called sprains and strains; chronic injury is joint instability. The acute injury may arise from blunt trauma to the muscle or joint; excessive exercise; or twisting, stretching, or forcible extension of a joint (e.g., “twisting” the ankle). Surgery is seldom needed unless complete rupture occurs, but the pain of such an injury can be severely limiting [70].

A sprain is an injury to a ligament caused by forcing a joint beyond its normal range of motion. The ligament may be stretched or actually torn. Sprains usually occur following a blunt blow during sports activities or falls. A strain is an injury to a muscle and/or tendon at any location from origin to insertion.

Strains are associated with excessive stretching of a muscle or muscle unit; they usually do not occur because of a blow or direct trauma. Poor conditioning, improper warm-up before activity, muscle fatigue or weakness, and strength imbalance can all contribute to muscle or tendon strain. Both strains and sprains have a high incidence of recurrence [70].

Clinical Manifestations

A sprain causes pain, swelling, local hemorrhage, spasm of the muscle that moves that joint, and disability. Pain occurs with passive movement of the joint, and there is intense pain over the involved ligament itself. Sprains are graded according to damage to the ligaments and the resultant joint instability [70]. A Grade I sprain is characterized by slight stretching and microscopic tearing of the ligament fiber, mild tenderness, and swelling around the joint. A Grade II sprain is identified by partial tearing of the ligament, moderate tenderness and swelling, and an abnormal looseness in the joint. The most severe is a Grade III sprain, which consists of a complete tear of the ligament, significant swelling and tenderness, and substantial instability.

The most common sprains affect the ankle and occur when inversion of the foot tears a ligament, usually the anterior talofibular ligament. Knee sprains cause swelling, hemarthrosis, significant decrease in range of motion, and joint laxity. Often the person hears a “pop” when the injury occurs and later describes the knee as feeling as if it is going to “give way.” The medial collateral ligament is most commonly involved [70]. Following the acute injury, patients are usually able to bear weight.

Strains cause pain, swelling, muscle spasm, and hemorrhage into the muscle. Discoloration and weakness may also be present. Pain increases with active flexion or passive stretching, which helps in differentiating strains from sprains. Strains are graded according to loss of muscle strength [70; 71]:

- Grade 1: A mild injury with no appreciable tissue tearing and no substantial (less than 5%) loss of function or strength
- Grade 2: A moderate injury with nearly half of muscle fibers torn, reduced strength, and some residual function
- Grade 3: A severe injury resulting from the complete rupture of the muscle, severe swelling and pain, and complete loss of function

Therapeutic and Specific Nursing Measures

Approaches to the treatment of strains and sprains are similar. Before initiating treatment, a thorough assessment and history to determine the nature and cause of the injury as well as any significant health problems that may influence the treatment. When a suspected strain or sprain occurs, the first-line treatment consists of five components known by the acronym PRICE:

- **Protection:** The affected joint or muscle should be covered to minimize the risk of additional traumatization.
- **Rest:** The patient should take steps to avoid use of the joint, tendon, or muscle to allow time for repair and healing.
- **Ice:** The application of cold will reduce pain and swelling (by causing vasoconstriction), and patients should be instructed to apply cold compresses up to several times per day, but to limit duration to 20 minutes or less.
- **Compression:** In order to reduce diapedesis and promote lymphatic drainage, the area may be bandaged. Patients should be instructed that wrappings should not be so tight as to restrict circulation.
- **Elevation:** The affected limb should be elevated to the level of the heart (or as close as possible) to promote venous return and reduce inflammation.

The PRICE regimen is usually continued for one week after injury, though there is some controversy about whether cold or heat is used after the first 24 hours. Cold is usually recommended for five to seven days because of its anti-inflammatory and analgesic effect. Then, wet heat may be used to aid in muscle relaxation and promote blood flow to the area [35].

With a second- or third-degree sprain, an x-ray should be taken to rule out fracture. Patients with sprains are usually immobilized for one week. When all pain on motion has ceased, patients can begin active range-of-motion and muscle-strengthening exercises. NSAIDs are the treatment of choice [35].

The PRICE regimen and NSAIDs are also appropriate for management of a strain. Emphasis is placed on prevention of recurrence through the use of muscle-strengthening and stretching exercises. Patients should be advised to engage in warm-up exercises before engaging in strenuous activity. For example, for patients with chronic ankle sprain/instability, slow stretching of the Achilles tendon daily can effectively reduce the incidence of recurrent sprain. Surgical intervention is recommended only in cases of complete muscle rupture [35].

Rhabdomyolysis

Rhabdomyolysis is a condition that develops as a result of the rapid dissolution of damaged or injured skeletal muscle [72]. Though not strictly a traumatic disorder, the most common cause of rhabdomyolysis is direct trauma to the skeletal muscle. However, any trigger of muscle destruction can theoretically result in rhabdomyolysis, and additional causes include infection, drugs/toxins, electrolyte disorders, endocrine disorders, extremes of body temperature, and excessive exertion [73].

As discussed, the function of skeletal muscle relies on ATP metabolism, electrolyte exchange, and intact myocytes. When these factors break down, the intracellular components of the muscle (e.g., electrolytes, creatine kinase, lactate dehydrogenase, myoglobin) are released into the body and enter the bloodstream. In more severe cases, this can lead to acute kidney injury, electrolyte imbalances, renal failure, and even death.

Clinical Presentation

The presentation of rhabdomyolysis is typically believed to consist of muscle pain, weakness, and discolored (reddish-brown) urine. Though this is considered the “classic” triad of symptoms, less than 10% of patients will present with all of these symptoms [73]. More than half of patients present only with myoglobinuria.

Diagnosis of rhabdomyolysis depends on detection of plasma creatine kinase. A diagnostic level has not been definitively identified, but most experts use a concentration five times the upper limit of the normal reference range (1,000 IU/L) [72].

Therapeutic Measures

Treatment of rhabdomyolysis focuses mainly on prevention of kidney damage and acute renal failure. Therefore, fluid therapy to increase urine output (and dilute urine) is the standard of care. The American Society of Nephrology has identified an ideal fluid regimen for these patients consisting of half isotonic saline (0.45%, or 77 mmol/L sodium), to which 75 mmol/L sodium bicarbonate is added [74]. At least 3–6 L should be administered per 24 hours; however, up to 10 L (or more) may be given if continuous supervision is possible. If necessary, 10 mL/hour of mannitol 15% may be added to further increase urine output. In cases that have already progressed to overt renal failure, extracorporeal blood purification is warranted [74].

In addition, supportive treatment of resultant hypovolemia and electrolyte imbalances (e.g., hyperkalemia, hypocalcemia) is necessary. Measures to help stabilize temperature are often necessary. Patients’ input and output should be monitored and documented. Pain management is often necessary, and patients should be assessed for severity and quality of ongoing pain.

SPECIFIC DISORDERS OF THE JOINTS

Because of their location and constant use, joints are particularly susceptible to stress, injury, and inflammation. In addition, many autoimmune disorders manifest in the joints.

DISORDERS OF MULTIFACTORIAL ORIGIN

A wide variety of joint conditions are multifactorial in origin, including ankylosing spondylitis, reactive arthritis, psoriatic arthritis, gout and pseudogout, low back pain, scoliosis, Charcot arthropathy, and carpal tunnel syndrome [37; 75].

Joint disorders of multifactorial origin can disrupt normal life activities, and families and job security can be negatively affected unless patients seek proper medical attention and counseling. Among the disorders discussed in this section, only gout can be cured, but the other disorders can be controlled to varying degrees so that in most instances the patient can maintain a fairly normal lifestyle [37; 75].

Psoriatic Arthritis

Psoriasis is often associated with inflammatory arthritis and a negative rheumatoid factor. Psoriatic skin lesions usually precede the development of arthritis, and in most cases, there is correlation between joint flares and skin flares. However, some patients with psoriatic arthritis have very mild or no psoriatic skin lesions. Heredity is the most specific risk factor, but environmental factors also play a role; the exact etiology is unknown [76].

Clinical Manifestations

The manifestations of psoriatic arthritis vary from patient to patient. Some have distal joint involvement, while others have widespread deformity, ankyloses, and joint destruction. The disease can be symmetrical or asymmetrical, and some patients have spondylitis, sacroiliitis, eye problems, or a combination. Nodules are not present with psoriatic arthritis [76].

In patients with psoriasis, silver-white scaly patches develop on the elbows, legs, scalp, and back. Nails are often pitted (20 pits or more per nail), and arthritis is more common with nail changes than with skin lesions. Onycholysis is common [76].

Joint symptoms usually begin with the acute onset of pain and swelling of distal interphalangeal (DIP) joints. A gout-like symptom in the great toe often gives a “sausage” appearance to the joint, but the disability is usually not as great as with rheumatoid arthritis. Spondylitis is often found in families with a strong background of psoriatic arthritis. Upon x-ray examination, some people show marked articular destruction with resorption of bone. A shortening of the middle phalanx of the DIP joints of fingers and toes has a characteristic cuplike appearance, and in some cases, an entire phalanx is destroyed. Extra-articular symptoms include conjunctivitis, episcleritis, or uveitis [76].

Laboratory studies of patients with psoriatic arthritis reveal mild anemia, an elevated ESR, negative rheumatoid factor, positive ANA, and an elevated uric acid level. Clinical diagnosis is made by considering nails, peripheral arthritis, and spinal involvement. Nail and skin changes in psoriatic arthritis may be hard to differentiate from those in reactive arthritis [76].

Therapeutic Measures

Therapeutic measures for psoriatic arthritis are aimed at both the arthritis and the psoriasis [76]. Nonpharmacologic approaches include physical and occupational therapy, exercise, smoking cessation, weight loss, and massage therapy. Symptoms may be controlled with NSAIDs and/or glucocorticoids

(oral or injection). In treatment-naive patients with active psoriatic arthritis, a tumor necrosis factor (TNF) inhibitor is recommended over oral, small-molecule drugs (OSMs) as a first-line option. However, OSMs may be used instead of a TNF inhibitor in patients without severe disease, particularly if they prefer an oral treatment option [77].

Gout

Gout is a metabolic disorder associated with elevated urate levels in the body and is the most common cause of inflammatory arthritis in the United States. Gouty arthritis is characterized by recurring episodes of acute, usually monoarticular, arthritis that tend to remit over several days to weeks; however, undiagnosed, untreated patients are at risk for developing a chronic deforming arthritis. An estimated 9.2 million adults in the United States are affected [78]. Gout is rarely encountered in persons younger than 30 years of age, with the predominant age range being 30 to 60 years. However, onset may occur in men in their early 20s who have a genetic predisposition and lifestyle risk factors. The peak age of onset in women is the sixth to eighth decade of life [78]. The estimated prevalence of gout is 5.9% in men and 2.0% in women [78]. The prevalence and incidence of gout has increased over the past several decades [78; 79].

Gout develops in persons with hereditary or acquired chronic hyperuricemia or in those with marked perturbations in serum urate associated with such factors as alcohol consumption, drug use, eating foods high in purines, overweight/obesity, and myeloproliferative disorders [78; 80]. The normal serum urate is generally considered to be ≤ 6.8 mg/dL. The majority of patients at the time of an acute flare have demonstrable hyperuricemia (in excess of 7 mg/dL); however, about 20% do not. The presence of hyperuricemia in the absence of symptoms is not diagnostic of gout [78]. In all cases, the hyperuricemia is caused by some dysregulation in the balance between production and excretion of urate. An estimated 80% to 90% of gout cases are due to urate underexcretion and not overproduction [78]. Hyperuricemia can occur without precipitating gout, and in the absence of symptoms, it may not warrant intervention [78; 81].

Uric acid is a final metabolic product of purine nucleotides found in many foods and in human tissue. Intermediary processes of purine metabolism include the initial breakdown of purines to inosine and then to hypoxanthine. Hypoxanthine is metabolized to xanthine, and xanthine to uric acid, with both stages catalyzed by the enzyme xanthine oxidase (the primary site for pharmacologic intervention by allopurinol) [82].

The human body is limited in its capacity to excrete a heavy urate load. In the setting of persistent hyperuricemia, often combined with stress to weight-bearing joints such as the great toe, monosodium urate crystals precipitate within joint synovial fluid, producing an intense inflammatory reaction. With chronicity, adjacent tissues may become saturated with urate, leading to deposits within articular, periarticular, bursal, bone, auricular, and cutaneous sites. These deposits, termed tophi, are detectable on physical exam or by radiographs and are a cardinal pathognomonic feature of gout. The presence of crystals, within joint fluid or in tissue, activates monocytes and macrophages to clear the crystals through phagocytosis. The release of proinflammatory cytokines and chemokines into the immediate area triggers an acute inflammatory reaction and influx of neutrophils into the joint space [83; 84; 85].

Clinical Manifestations

The clinical presentation of gout is typically one of arthritis and intense pain, and patients may exhibit inflammation and edema in the afflicted joint. Although the great toe is the most common site, other joints and their surrounding tissue can be affected, including the insteps, ankles, heels, knees, wrists, fingers, and elbows [78]. Gout may be confused with other causes of arthritis as all forms share the cardinal signs of inflammation: pain, redness, warmth, tenderness, and swelling [80; 86]. While gout initially manifests in severe, discrete episodes of pain, the condition may progress to more frequent attacks with shorter asymptomatic periods between attacks [78; 86]. Synovial fluid analysis is the gold standard for diagnosing gout, confirmed by the presence of monosodium urate.

Therapeutic and Nursing Measures

Gout is perhaps the most easily treated, and preventable, form of arthritis. This is due to widespread understanding of its underlying mechanisms and the availability of effective treatment [80]. It is managed by controlling the current acute attack and preventing future attacks. Medications addressing the underlying pathophysiology include the xanthine oxidase inhibitors (XOIs) allopurinol and febuxostat and the uricosuric agents probenecid, fenofibrate, and losartan [80; 87]. (Note: The use of fenofibrate and losartan for the treatment of gout is off label.)

The initial steps include patient education, testing to rule out other causes of hyperuricemia, and evaluation of the disease burden to determine appropriate treatment. All patients with hyperuricemia and established gout should be advised to begin dietary modification. This involves avoiding organ meat high in purine content, high-fructose corn syrup, and excessive



The American College of Rheumatology conditionally recommends that patients with gout limit their consumption of purine-rich foods (e.g., meat and seafood), alcohol, and high-fructose corn syrup (particularly in sweetened soft drinks and energy drinks).

(<https://www.rheumatology.org/Portals/0/Files/Gout-Guideline-Final-2020.pdf>. Last accessed September 26, 2022.)

Certainty of Evidence: Low or very low

alcohol use. Portions of high purine-content seafood, sugar, and salt should be limited. The ideal diet will include low- or non-fat dairy products and vegetables. Other lifestyle modifications can also assist in managing gout, including weight loss in overweight patients, regular exercise, smoking cessation, and adequate hydration [83; 87; 88].

The acute pain of gout may be treated with NSAIDs, a cyclooxygenase-2 (COX-2) inhibitor, systemic corticosteroids, or oral colchicine monotherapy in mild-to-moderate disease (≤ 6 on a 10-point pain scale). Combination therapy (i.e., colchicine and NSAIDs, oral corticosteroids and colchicine, or intra-articular steroids with each of the other options) may be used in cases of severe disease with intense pain and polyarticular presentation. Intramuscular triamcinolone acetonide is recommended in patients unable to take oral medication or likely to be poorly adherent to the multidose oral regimen [87].

An inadequate response to therapy after escalation ($< 20\%$ pain reduction within 24 hours or $< 50\%$ pain reduction after ≥ 24 hours) should prompt reconsideration of the diagnosis. If gout is confirmed, switching to another form of monotherapy or adding a second agent may prove effective [83; 87].

Urate-lowering therapy should be initiated in all patients with tophaceous gout, radiographic damage due to gout, or frequent gout flares [88]. Therapy should be started within 24 to 36 hours of the onset of an acute gout attack unless otherwise contraindicated. Urate-lowering therapy is not recommended for patients experiencing their first flare, or for patients with asymptomatic hyperuricemia (serum urate > 6.8 mg/dL) with no prior gout flares or subcutaneous tophi [88]. Allopurinol (≤ 100 mg/day) is the preferred first-line agent. Febuxostat (≤ 40 mg/day) is an acceptable alternative [88]. Probenecid may be used as an alternative to allopurinol or febuxostat if there is contraindication or intolerance to these preferred agents. However, probenecid should be avoided in patients with a history of urolithiasis [83; 87; 88].

Clinicians may also consider screening for the HLA-B*5801 allele, which is associated with high risk of severe allopurinol hypersensitivity reaction. High-risk persons include Koreans with an estimated glomerular filtration rate < 60 mL/min/1.73 m² or those with Han Chinese or Thai ancestry [89].

Anti-inflammatory prophylaxis (against precipitating an acute flare) is recommended when initiating urate-lowering therapy in asymptomatic patients [88]. Colchicine was once the treatment of choice but is now less commonly used than NSAIDs because of its narrow therapeutic window and risk of toxicity [90]. To be effective, colchicine therapy is ideally initiated within 36 hours of onset of the acute attack [78]. In the case of colchicine intolerance or contraindication, prednisolone may be used [88]. Prophylaxis should continue after achieving target serum urate level for three months in patients without tophi, for six months in patients with resolved tophi, and with any remaining signs of gout activity in all patients [88].

Patients with intermittent symptoms or chronic synovitis with tophi (chronic tophaceous gouty arthritis) should be treated with a single-agent XOI, such as allopurinol, at a dose to achieve and maintain the serum urate level within normal range [83; 87]. If the serum urate target is not achieved or disease activity persists, a uricosuric agent may be added to the XOI. Pegloticase therapy should be considered if the serum uric target is not achieved, disease activity persists, more than seven attacks occur per year and no tophi, two or more attacks per year and tophi, or chronic tophaceous gouty arthritis is present [88; 89].

Pseudogout

Calcium pyrophosphate deposition disease, or “pseudogout,” is a similar crystalline arthritis that occurs in patients with underlying osteoarthritis and is identified by the presence in synovial fluid of calcium pyrophosphate dehydrate crystals [78; 80; 85]. X-ray findings of articular cartilage calcification usually accompany it. Many patients with pseudogout have other disorders, such as diabetes, hypothyroidism, and gout [37].

Clinical Manifestations

In pseudogout, arthritis occurs in a large joint, which is erythematous, swollen, warm, and painful. Like gout, pseudogout is usually monoarticular, but involvement of other joints can follow in succession. Attacks are often precipitated by trauma, surgery, or medical illness. Onset of symptoms is rapid, with a peak in 12 to 36 hours. Episodes are intermittent, usually involve the same joint, and typically last about one to two weeks. Joints are normal between attacks [37].

Therapeutic and Nursing Measures

The deposition of calcium pyrophosphate dihydrate crystals cannot be reversed. Acute attacks of pseudogout are treated with NSAIDs, colchicine, and/or oral corticosteroids; in more severe cases, drainage of the affected joint may be helpful [37].

Nursing measures for pseudogout include careful joint assessment, thermo- or cryotherapy, and monitoring for symptoms and signs of systemic illness and side effects of medications [18]. Patient education for home care includes instruction in the safe application of heat or ice, range of motion exercises, and the nature and side effects of prescribed medications. A weight-reduction diet can be helpful promote the long-term health of weight-bearing joints [18].

Low Back Pain

When it occurs, back pain is most often localized to the lower back, and chronic back pain is almost always chronic low back pain. Although acute-onset low back pain is a common problem that usually resolves within four to six weeks, many patients develop a persistent, disabling pain syndrome with a diminishing prognosis for return to normal function. When low back pain continues beyond 12 weeks, the prospect for subsequent remission is poor and progression to chronic low back pain is likely. Chronic low back pain imposes a great burden: for patients, pain and disability; for society and the healthcare system, an enormous expense in direct and indirect costs.

Risk factors for developing low back pain can be generally categorized as nonmodifiable, such as old age, female sex, poverty, and lower education level, and modifiable, including higher body mass index (BMI), smoking, lower perceived general health status, physical activity (e.g., bending, lifting, twisting), repetitive tasks, job dissatisfaction, and depression. The greatest contributors to low back pain episodes are single-event or repetitive exposures to mechanical stress and age-related degenerative spinal changes. With chronic low back pain, mechanical and biophysiologic factors play a minimal secondary role to the primary contribution from psychosocial factors [91].

Clinical Manifestations

The onset of low back pain is described as discomfort in the vicinity of the low back ranging from a dull ache to a sudden, sharp, shooting or stabbing pain and may include limited flexibility and/or range of motion or inability to stand straight [92]. Although the symptoms of back pain can originate anywhere from the thoracic spine to the sacrum and coccyx, most cases originate in the lumbar spine, as this is the site of support for upper body weight [92].

With low back pain, the clinical presentation varies according to etiology. In general, radicular pain suggests nerve root involvement, while axial pain suggests disk degeneration, facet arthropathy, sacroiliac (SI) joint arthropathy, or myofascial pathology of the spine.

Nonspecific Low Back Pain. Up to 85% of low back pain in patients presenting to the primary care setting is nonspecific, meaning that it lacks a clear origin and is not caused by specific local or systemic disease or spinal abnormality [93]. Nonspecific low back pain is a diagnosis of exclusion made after ruling out serious causes of the back pain. Although pain can originate from ligaments, facet joints, muscle, fascia, nerve roots, the vertebral periosteum, or outer portions of the disk, the effective management of nonspecific low back pain does not require a precise anatomic diagnosis [94]. The pain is usually unilateral and may radiate to the buttocks or posterior thigh but not past the knee. This can lead to incorrect diagnosis of radiculopathy or disk herniation. However, true radicular symptoms radiate below the knee in a dermatomal distribution and can involve sensory loss, weakness, or reflex changes. Painful spasm may be present, and pain may be worsened by movement, while lying flat decreases the pain. Complaints of numbness, weakness, or bowel or bladder dysfunction are absent [95]. Degenerative changes revealed by lumbar imaging should usually be considered nonspecific, because they poorly correlate with symptom severity [96].

Lumbosacral Radiculopathy. Lumbosacral radiculopathy is a clinical diagnosis of nerve root irritation and compression, resulting in a symptom distribution of the affected lumbar or sacral nerve root such as numbness, weakness, or paresthesia.

Sciatica is the most common symptom of lumbar radiculopathy and refers to pain that radiates down the leg below the knee in the distribution of the sciatic nerve to indicate nerve root compromise from mechanical pressure or inflammation [96].

Causes of lumbar radiculopathy include disk herniation, arthritic degeneration, cord compression, spinal stenosis, tumor, and infection. With herniated disk, the pain is described as a deep, aching, axial midline pain concurrent with radicular pain. Discogenic pain results from a tear in the outer disk layer (annulus fibrosus) that causes the inner gelatinous material (nucleus pulposus) to prolapse, inflame, and compress a nerve root [95]. The resulting pain from pressure and nerve irritation improves with the resolution of local inflammation, and the disk protrusion may spontaneously remit with time. Although disk herniation and radiculopathy are often viewed as causally linked, herniation is often asymptomatic and only occasionally the cause of sciatica [95].

Lumbar Spinal Stenosis. Lumbar spinal stenosis refers to the frequently age-related narrowing of the spinal canal that may result in bony constriction of the cauda equina and the emerging nerve roots [96]. Spinal stenosis can produce pain in the low back that radiates down the back of both legs, often worsened with standing or walking. To make the pain more bearable, patients often walk a short distance with a hunched back, and then sit down for relief. The pain will then dissipate after several minutes. Congenital lumbar canal stenosis is a predisposing factor. Patients show less tenderness over the lumbar spine than those with acute lumbar disk herniation, and the straight leg-raising test may be normal [97].

Most persons 60 years of age and older exhibit varying degrees of spinal stenosis from disk herniation, osteophytes, or degenerative spondylolisthesis. Fortunately, clinical pain manifests in less than 30% and, just as with degenerative disk disease, there is poor correlation between symptom severity and extent of spinal canal stenosis revealed by MRI [95].

Myofascial Pain. Myofascial pain of the low back or neck is common, especially following trauma or repetitive motion injury. This is thought to result from strain or sprain to the muscles and ligaments. Myofascial pain is described as a deep, aching, poorly localized discomfort made worse by activity. It can be limited to discomfort in the paraspinal muscles or may extend to the buttocks and upper thigh areas [98].

Epidural Compression Syndrome. Epidural compression syndrome is an umbrella term that encompasses spinal cord compression, cauda equina syndrome, and conus medullaris syndrome. While these conditions differ in the level of neurologic deficit at presentation, they are otherwise similar in symptoms, evaluation, and management. Massive herniation of a midline disk, typically at the L4 to L5 disk level, is the most common cause of epidural compression syndrome. Tumor, epidural abscess, spinal canal hematoma, or lumbar spine spondylosis represent other causes [95].

In these patients, neurologic status at diagnosis is the greatest predictor of ultimate neurologic outcome and underscores the importance of early accurate diagnosis. The dominant symptom is back pain with accelerating pain severity. Pain from epidural spinal cord compression is made worse with recumbent positioning, and unilateral or bilateral radiculopathy may develop over time. For many patients, leg pain or neurologic symptoms are more dominant than back pain. Also common at diagnosis is symmetrical lower extremity weakness that may have progressed to gait disturbance or paralysis. Decreased lower extremity reflexes are associated with cauda equina syndrome [95].

Lumbar Facet Joint Syndrome. Lumbar facet joint syndrome is seen in as many as 35% of patients with low back pain and is frequently associated with arthritis or lumbar facet joint injury [97]. Dominant symptoms include unilateral low back pain that may radiate down the back or front of the thigh and morning stiffness with isolated facet arthropathy [99]. Tenderness is usually found over the lumbar paraspinal muscles and facet joints. Back pain is worsened with back extension and lateral rotation to the side of the pain, and the leg-raising test is negative. MRI and CT findings of facet joint arthropathy do not correlate with clinical findings [97].

Sacroiliac Joint Syndrome. SI joint syndrome typically manifests as localized pain in the lower back or upper buttock area that overlies the SI joint. Pain is intensified by attempts to walk up stairs, and while pain may be referred to the posterior thigh, extension below the knee is unusual [100]. Tenderness over the SI joint is often found in physical examination, and pain is aggravated by the Patrick test or single-leg standing [97]. The onset of SI joint pain is usually gradual (over months to years), and although etiology is often elusive, trauma, infection, and tumor represent infrequent yet known causes of SI joint pain [100].

Assessment and Diagnosis

Most patients with acute low back pain have ligamentous or muscle strain syndrome, follow a benign course, and show significant improvement within two to three weeks. The challenge for clinicians is to recognize early the possibility of serious disease, such as spinal metastatic cancer or vertebral and epidural space infection, and then to identify those with herniated disk, radiculopathy, or spinal stenosis.

The proper assessment of the patient with back pain requires vigilance and careful attention for factors and warning signs suggestive of serious or life-threatening disorders. A thorough history and physical examination should be performed on all patients, during which the patient is assessed for the presence of warning signs or “red flags.” Red flags represent alarm symptoms or signs that warrant prompt, specific diagnostic testing, urgent treatment, or referral to a specialist. Among these are weight loss, prior history of cancer, nocturnal or rest pain, age older than 50 years, recent trauma, fever and

chills, history of injection drug use, chronic corticosteroid therapy, difficulty urinating, bowel or bladder incontinence, and neurologic deficits such as saddle anesthesia, perianal or perineal sensory loss, or motor weakness in the extremities [93; 95; 101]. As an example, there is a common association between spontaneous vertebral fracture and any combination of age older than 70 years, female gender, recent trauma, and prolonged corticosteroid use. There is also a moderate to highly significant predictive value for age older than 50 years, history of prior cancer, unexplained weight loss, and failure of conservative therapy in identifying spinal malignancy [101].

Patients should also be assessed for “yellow flags,” or risk factors for poor prognosis and chronicity [94; 101]. Areas to explore include maladaptive beliefs, attitudes, and behaviors regarding the back pain and recovery, such as passivity or reluctance to self-manage, dependency on the provider to “cure,” fear avoidance beliefs, and beliefs that harm will come from activity and discomfort. Other areas include depression, anxiety, maladaptive coping response to stress, social withdrawal or isolation, and lack of social support. Adverse economic and work environment circumstances, such as job dissatisfaction, excessive and inflexible physical workplace demands, high levels of work-related stress, poor workplace social support, and adversarial or dysfunctional workplace relationships should also be noted [94]. Early detection and intervention (if indicated) for problematic motivational, emotional, or social dysfunction are important because these factors influence the selection and effectiveness of therapeutic interventions.

In the absence of red flags, use of imaging and diagnostic tests for acute low back pain is discouraged, as imaging findings rarely change clinical management. Overuse of lumbar imaging in low back pain correlates with, and likely contributes to, the two- to threefold increase in surgical rates for low back pain over the last 10 years [93]. Assigning significance to imaging anomalies requires skill at the specialist level to integrate historical, clinical, and imaging findings. Imaging abnormalities are essentially normative by 40 years of age; for instance, 80% of persons 60 years of age and older exhibit a protruding disk, which is symptomatic for only a fraction of patients. Incorrect communication of imaging findings to the patient may lead to patient fixation, contribute to fear-avoidance behaviors, and increase the risk of iatrogenic aggravation of chronic low back pain. Guidelines suggest that physicians without advanced training defer imaging tests to qualified specialists [93; 102].

However, imaging and other testing should be performed in patients with new-onset or progressive neurologic deficits and those with suspicion of serious underlying conditions. In patients with persistent pain and symptoms consistent with radiculopathy or spinal stenosis, MRI should be performed only when such patients are candidates for surgery or epidural steroid injection. CT scanning is an alternative option to first-line MRI [93].

Therapeutic Measures

Although acute low back pain improves in most patients within three to six weeks using conservative therapy, up to 33% of patients with low back pain report pain of moderate or greater severity at one-year follow-up and 20% report ongoing pain severe enough to limit activity [96]. With chronicity, low back pain may become disabling and impose a severe emotional and functional burden. The management goals for chronic low back pain are to minimize pain and disability, improve functional status, and facilitate restoration of normal activity, while limiting the use of marginally effective or inappropriate medication [93].

Many pharmacologic therapies and minimally invasive or invasive procedures have been utilized in a strategy designed simply to relieve pain—with variable results. However, there is little evidence these focused pain approaches are comparable or superior to interventions that focus primarily on restoration of function instead of pain relief. This contradicts the biomedical model in medicine that emphasizes escalation of costly and invasive therapies to achieve “pain cure” in patients lacking response to lower-intensity approaches [102; 103]. It is now recognized that treatment for conditions such as chronic low back pain persisting in the absence of a unique underlying pathologic lesion must address potential contributory factors such as affective disorders, maladaptive beliefs and coping skills, and interpersonal and occupational dysfunction. Dysregulated cortical, pre-frontal, and higher neural level mechanisms associated with chronic low back pain are being identified and may represent therapeutic targets in functional restoration-based approaches. As with other chronic pain syndromes, greater understanding of pain pathway alterations will better inform therapy selection.

Virtually universal among practice guidelines for chronic low back pain is the emphasis on a multidisciplinary, multi-modal approach that includes exercise and activity, cognitive restructuring of maladaptive attitudes and coping skills, a behavioral component addressing fear avoidance, physiotherapy and manual therapy, and analgesics as indicated [93; 96; 102; 104; 105; 106; 107]. This is often best accomplished by consultation or referral to an established pain treatment center. Multidisciplinary functional restoration programs, which are intensive (more than 100 hours) biopsychosocial interventions whereby physical rehabilitation is combined with cognitive-behavioral therapy and delivered by an interdisciplinary team, embody this recommendation. Moderate-to-strong evidence supports their efficacy in chronic low back pain. They have been found effective in reducing pain and improving physical function, work readiness, and return to work. Weaker outcomes are found in programs that are less intensive or lacking a behavioral component. Patients who do not improve with less intensive therapy options and have high levels of pain, distress, and disability should be considered for multidisciplinary functional restoration programs [94].



According to the Institute for Clinical Systems Improvement, clinicians should advise patients with acute and subacute low back pain to stay active and continue activities of daily living within the limits permitted by their symptoms.

(<https://www.icsi.org/guideline/low-back-pain>. Last accessed September 26, 2022.)

Strength of Recommendation/Level of Evidence:
Strong Recommendation/Moderate Quality Evidence

Scoliosis

Scoliosis is a lateral curvature of the spine, most commonly in the thoracic area with convexity to the right and compensatory convex curve to the left in the cervical and lumbar areas. Scoliosis can be functional (a result of poor posture or leg-length discrepancy) or structural (a result of deformity of the vertebral bodies, paralysis, congenital malformations, or idiopathic causes). Idiopathic causes are the most common and appear with increased growth during adolescence. It disproportionately affects girls, who are 10 times more likely to be diagnosed than boys at 10 years of age or older [108].

Clinical Manifestations

Symptoms of backache, fatigue, and dyspnea occur only after scoliosis is well established. Untreated scoliosis can result in pulmonary insufficiency from decreased lung capacity, back pain, degenerative arthritis of the spine, intervertebral disease, and sciatica [108]. While screening for idiopathic scoliosis has typically occurred between 10 and 18 years of age, the current evidence is insufficient to assess the balance of benefits and harms [109].

Older patients may exhibit kyphosis, a postural curvature of the spine that is due to aging, disc degeneration, atrophy of spinal muscles, osteoporosis, or vertebral collapse. Adults with kyphosis have a rounded back and possible weakness and generalized fatigue. Kyphosis rarely produces local tenderness except in severe osteoporosis with compression fractures [108].

Therapeutic Measures

Early treatment of scoliosis consists of a combination of physical therapy, bracing, and/or surgery. If the condition is untreated in adolescence, problems that develop can only be treated symptomatically. Any upper respiratory tract infections are treated aggressively to prevent pneumonia and atelectasis [108].

Specific Nursing Measures

Patients with scoliosis often have body image issues and difficulty finding clothes that fit properly. When patients with scoliosis are hospitalized for any problem, careful attention to positioning is essential; improper positioning is not only extremely uncomfortable for the patient, but it can precipitate a vertebral fracture, especially in those with osteoporosis [108].

Carpal Tunnel Syndrome

Carpal tunnel syndrome is generally associated with such umbrella terms as repetitive stress injuries, work-related upper extremity disorders, musculoskeletal disorders, entrapment neuropathies, and cumulative trauma disorders [110; 111]. Specifically, carpal tunnel syndrome is a painful disorder of the wrist and hand that occurs when the median nerve (which runs from the hand to the forearm) becomes compressed [112; 113].

The carpal tunnel is a narrow passageway on the palm side of the wrist. Surrounded by bones and ligaments, the carpal tunnel houses and protects the tendons of the hand and the median nerve, which controls sensations to the thumb and fingers. When the median nerve becomes pinched or compressed (due to swelling or irritation in adjacent tissues or tendons), the result can be pain, numbness, hand weakness, and in extreme cases, loss of hand function. Cases of bilateral carpal tunnel syndrome have been reported, but typically only one hand is affected [112; 114; 115]. Carpal tunnel syndrome is rare in children; it usually occurs only in adults [116].

Clinical Manifestations

The symptoms of carpal tunnel syndrome typically appear gradually and may include [114; 116]:

- Numbness, burning, or tingling in the fingers and palm of the hand
- Pain in the wrist, palm, or forearm, especially during use
- Decreased grip strength
- Weakness in the thumb
- Sensation of swollen fingers, whether or not swelling is apparent
- Difficulty distinguishing between hot and cold

Symptoms may cause waking during the night with the urge to “shake out” the hand or wrist. Symptoms may occur with activities that require prolonged grasping and/or flexing of the wrist (e.g., driving, holding a book). Left untreated, carpal tunnel syndrome can progress to persistent numbness and permanent loss of hand function. In severe and chronic cases, irreversible muscle damage or atrophy may occur [112; 116; 117]. Complete sensory loss in the hand has also been reported.

Assessment and Diagnosis

Early diagnosis of carpal tunnel syndrome is important to prevent muscle atrophy or damage to the median nerve that cannot be reversed by treatment [112; 116]. Early diagnosis, including a physical examination, medical history, routine laboratory tests, and imaging, can also help to identify or rule out other health conditions that may present with similar signs and symptoms and require specialized treatment [118; 119]. The physical examination should include specific testing, such as Phalen’s maneuver or Tinel’s sign, that can produce the symptoms of carpal tunnel syndrome [114; 116]. In elderly patients, particular attention should be given to the objective evidence of carpal tunnel syndrome rather than subjective complaints [120].

Therapeutic and Nursing Measures

Surgery, corticosteroids, NSAIDs, diuretics, wrist splints, exercise, ultrasound therapy, laser therapy, and yoga are among the methods that have been recommended for the treatment of carpal tunnel syndrome [121; 122; 123; 124; 125]. Although no single treatment method has been universally accepted, there is agreement that the treatment of carpal tunnel syndrome should begin as early as possible and should include attention to underlying causes, such as diabetes or rheumatoid arthritis. There is also agreement that successful treatment depends on patient compliance with the treatment program [116; 126].

Corticosteroid injection has been found to improve patient satisfaction, symptoms, and function when measured at intervals of 2, 4, 8, and 12 weeks. As noted, it demonstrates a more significant overall improvement in the symptoms of carpal tunnel syndrome than oral corticosteroids but does not appear to provide a better long-term outcome (greater than six months) than splinting or NSAIDs. Two treatment injections do not appear to provide any added benefit when compared to one treatment injection [124; 127].



According to the American Academy of Orthopaedic Surgeons, strong evidence supports that the use of steroid (methylprednisolone) injection should improve patient-reported outcomes in those with carpal tunnel syndrome.

(https://www.aaos.org/globalassets/quality-and-practice-resources/carpal-tunnel/cts_cpg_4-25-19.pdf.)

Last accessed September 26, 2022.)

Strength of Recommendation: Strong (Evidence from two or more “high” strength studies with consistent findings for recommending for or against the intervention)

Splinting has been found to improve patient satisfaction, symptoms, and function when measured at intervals of 2, 4, and 12 weeks. The American Academy of Orthopaedic Surgeons suggests that splinting be considered before surgery. This may be particularly helpful when weighing the risks of surgery versus the benefits. Splinting is not recommended for use after routine carpal tunnel release surgery. The benefit of splinting for postoperative rehabilitation is undetermined [126; 127; 128].

NSAIDs are used to treat a variety of acute and chronic pain conditions, including carpal tunnel syndrome, but opinion varies as to their effectiveness and safety for long-term use [129; 130; 131]. Specifically, NSAIDs have been associated with gastrointestinal and cardiovascular risks and toxicity with long-term use [132].

Diuretics and vitamin B6 (pyridoxine) may also help with temporary relief of symptomatic carpal tunnel syndrome, but their long-term benefits are unproven [127; 131; 133]. Acupuncture, yoga, exercise, laser therapy, activity modification, and ergonomic workplace modifications also have been mentioned as non-surgical treatment alternatives, but most experts agree that further research is needed to determine the viability and efficacy of these methods [116; 124; 126; 127; 131; 134; 135].

Carpal tunnel release is the preferred treatment for patients with chronic or severe carpal tunnel syndrome. It is achieved by either an open or endoscopic procedure [116; 122; 126; 128]. Both types of surgery are generally performed on an outpatient basis under local anesthesia. Open release surgery involves making an incision of up to 2 inches at the base of the palm of the hand and cutting the transverse carpal ligament, which releases pressure on the median nerve [116; 136]. Endoscopic surgery involves making a small, one-half inch incision at the wrist and introducing an arthroscope beneath the transverse carpal ligament. Using the scope as a guide, the ligament is cut, relieving pressure on the median nerve [116; 134; 136].

DEGENERATIVE DISORDERS

Osteoarthritis

Osteoarthritis is the most common form of arthritis and is characterized by degeneration of cartilage and its underlying bone within a joint, with resultant bony overgrowth. This process of tissue breakdown eventually leads to pain and joint stiffness [137]. Osteoarthritis develops most frequently in the knee, hip, and hand. Although pain in the lower back and the neck are the most frequently occurring musculoskeletal conditions and are the leading cause of functional limitation and work absences, the etiology of back and neck pain is often unclear, with many cases involving muscles and ligaments rather than osteoarthritic changes [138; 139; 140].

Osteoarthritis is classified as primary or secondary. The cause of primary osteoarthritis is idiopathic; no abnormality is the cause of changes in the joint [141]. Secondary osteoarthritis is the result of a known cause, most often trauma/injury

or systemic diseases. Secondary osteoarthritis is most often found in the shoulder, elbow, and ankle and is more likely to become clinically apparent at a younger age than primary osteoarthritis [141; 142; 143; 144]. A population-based study showed that secondary osteoarthritis related to trauma accounts for approximately 12% of the overall prevalence of symptomatic osteoarthritis of the knee, hip, or ankle [145]. Injuries sustained in sports activities comprise a large portion of post-traumatic osteoarthritis [146]. A wide variety of systemic diseases have been identified as frequent causes of secondary osteoarthritis; these conditions include metabolic diseases, endocrine disorders, bone dysplasias, and crystal deposition diseases [141; 147].

Clinical Manifestations

The diagnosis of osteoarthritis at most joints is made primarily on the basis of clinical findings, with imaging studies and laboratory tests more useful for ruling out other diagnoses rather than for confirming the diagnosis of osteoarthritis [148; 149; 150]. Although radiographic findings are considered to be diagnostic criteria for osteoarthritis, radiographs are not usually part of the initial diagnostic evaluation for several reasons. The primary reasons are the lack of evidence of early osteoarthritic changes on radiographs and the poor correlation between symptoms and radiographic evidence of osteoarthritis [148; 151; 152; 153]. Thus, the absence of radiographic evidence of osteoarthritis in the presence of joint-related symptoms should not exclude the diagnosis of osteoarthritis.

When obtaining a history, questions should focus on the nature of joint-related symptoms, patients' self-reports of limitations in function or activities, and information related to established risk factors for osteoarthritis. The following questions can help elicit important information needed for a diagnosis:

- Do you have any joints that hurt? If so, how long have they been bothering you?
- When does the pain occur? After certain physical activities? At rest?
- Do you have relief of pain if you rest?
- Does the pain bother you at night? Does pain wake you up at night?
- Are your joints stiff when you wake up in the morning? If so, how long does the stiffness last?
- Do the joints that hurt ever lock up or give out on you?
- Do you have a family history of osteoarthritis or rheumatoid arthritis?
- What types of recreational activities or sports do you participate in? If you play sports, do you do so for leisure or competitively?
- What is your occupation? Are there tasks or activities that are part of your job that bother any joints?

- Have you ever had an injury to a joint?
- Are there daily activities or other tasks that you cannot do because of pain or other symptoms in any joint?

The primary symptom of osteoarthritis of the knee is pain, especially with weight-bearing exercise or activity, that improves with rest. Stiffness in the joint occurs in the morning, lasting 30 minutes or less, and may occur after periods of inactivity [154]. The clinical presentation of hip osteoarthritis is similar to that of knee osteoarthritis, with pain being the most common symptom driving individuals to seek medical care [155]. Pain related to hip osteoarthritis is an ache—most often diffuse—that is usually felt during use of the joint and relieved by rest. Pain is typically gradual, variable, or intermittent; the joint may feel stiff after a period of inactivity [155]. The loss of function or mobility is usually related to the degree of pain.

Osteoarthritis of the hand is characterized by pain with use, which affects one or a few joints at any one time, and mild stiffness in the morning and/or after a period of inactivity [158]. The severity of osteoarthritis-related pain varies, and the pain may be intermittent. The joints most often affected are the distal and proximal interphalangeal joints and the base of the thumb [156; 157; 158]. Individuals who have evidence of osteoarthritis at several joints in the hand are at increased risk for generalized osteoarthritis, and clinicians should evaluate such patients as appropriate [158].

Pain related to osteoarthritis of the shoulder is typically progressive, related to activity, deep in the joint, and often localized posteriorly [142]. Pain is usually present at rest and interferes with sleep, with nocturnal pain becoming more common as the disease progresses. More advanced disease is also associated with stiffness that limits function.

Individuals with osteoarthritis of the elbow typically have pain, stiffness, and weakness in the joint [143]. Later stage disease is associated with pain when carrying a heavy object at the side of the body with the elbow in extension. The history is important when evaluating symptoms related to the elbow because of the strong relationship between trauma or occupation with osteoarthritis, especially in individuals who are younger than 40 years of age [159].

A history of ankle fracture or ligamentous injury is a hallmark feature of osteoarthritis of the ankle [144]. Diagnostic evaluation includes radiographs of the ankles made with the patient standing. MRI is also recommended, as it can provide evidence of osteonecrosis as well as indicate the amount of involvement, the extent of bone loss, and the size of subchondral cysts [144].

Therapeutic Measures

There is currently no curative therapy for osteoarthritis, and treatments to alter or arrest the disease process are few and mostly ineffective [151]. As clinicians on the frontline of care, primary care providers and nurses are typically the first to see individuals with symptoms indicative of osteoarthritis. Primary

care providers can coordinate the management of osteoarthritis, and a multidisciplinary approach is best. The ACR and the Association of Rheumatology Health Professionals (a division of the ACR) support such an approach, noting that the healthcare team may include a rheumatologist, primary physician, nurse, nurse practitioner, physician assistant, physical therapist, occupational therapist, physiatrist, psychiatrist, psychologist, orthopedic surgeon, social worker, registered dietician, vocational counselor, and others [160].

The optimal management of osteoarthritis encompasses both nonpharmacologic and pharmacologic measures, beginning with basic modalities and following a so-called pyramid approach as the disease progresses or symptoms do not respond [161]. Several factors should be considered when selecting treatment modalities, including risk factors (e.g., age, comorbidity, overweight/obesity), the level of pain and functional limitations, signs of inflammation, and degree of structural damage [162].



According to the American Academy of Orthopaedic Surgeons, oral acetaminophen is recommended to improve pain and function in the treatment of knee osteoarthritis when not contraindicated.

(<https://www.aaos.org/globalassets/quality-and-practice-resources/osteoarthritis-of-the-knee/oak3cpg.pdf>. Last accessed September 26, 2022.)

Strength of Recommendation: Strong (Evidence from two or more “high” quality studies with consistent findings for recommending for or against the intervention.)

Many treatment options are associated with benefits and risks, and the clinician should discuss the benefits and risks with patients and support their participation in the decision-making process [163; 164]. Patient preferences are an important consideration when choosing treatment options and establishing treatment goals, and the ACR advocates care that addresses treatment goals that are meaningful to the individual patient [160]. Decision aids can help enhance patients’ knowledge of treatment options, improve patients’ participation in their care, and produce realistic expectations of outcomes [164]. Decision aids for osteoarthritis have been developed in a variety of media (e.g., print, online, video) and are available online (<https://decisionaid.ohri.ca>) [164].

The pain and disability associated with osteoarthritis often has a substantial psychologic and social effect. It is important to discuss these aspects with patients and to address psychologic issues, especially depression, in order for treatment measures to be effective [165].

Specific Nursing Measures

Education and self-management, through lifestyle modifications are universally recognized as the core of treatment in clinical guidelines [166]. This recommendation is based on research showing that education helps patients become more involved in their care, leading to improved outcomes [163]. The Agency for Healthcare Research and Quality notes that an effective partnership is the key to the effective management of osteoarthritis; the healthcare professional's role in this partnership is to [163]:

- Encourage patients to change their behavior to improve symptoms or slow disease progression
- Promote the proper use of medications
- Instruct patients on how to interpret and report symptoms accurately
- Support patients' efforts to maintain normal activities
- Help patients adjust to new social and economic circumstances and cope with emotional consequences

Nurses should emphasize to patients that adhering to the management program will alleviate their symptoms, improve their function, and enhance their quality of life. Education should be tailored to address individual needs. For example, patients who participate in sports should be advised to avoid sports with direct contact and high impact and to wear protective equipment to prevent injury [167]. Similarly, for patients in occupations with high risk for osteoarthritis, clinicians should discuss the importance of avoiding high-risk tasks. It is also essential to encourage patients with osteoarthritis of the glenohumeral joint or the elbow to modify activities that led to the development of the disease [143; 159]. Periodic contact during follow-up can help promote self-management [166].

IMMUNOLOGIC DISORDERS

Rheumatoid Arthritis

Rheumatoid arthritis is defined as a chronic inflammatory disease characterized by uncontrolled proliferation of synovial tissue and a wide array of multisystem comorbidities [24]. In its most common presentation, rheumatoid arthritis affects the joints, causing inflammation of the synovium and cartilage and bone loss. The precise etiology of rheumatoid arthritis is presently unknown [26]. Most likely it has an autoimmune origin (whereby an individual's immune system confuses healthy synovial tissue for foreign substances, thereby attacking the synovial joint surfaces) given that autoantibodies (e.g., rheumatoid factor, ACPA) are present and often precede the clinical manifestation of rheumatoid arthritis by many years [22; 25; 168].

The course and severity of the illness can vary considerably, and infection, genetic factors, and hormones may contribute to the disease. Rheumatoid arthritis appears to require the complex interaction of genetic and environmental factors with the immune system and ultimately in the synovial tissues throughout the body. Triggers for rheumatoid arthritis have long been the target of active research. Purported triggers have included bacteria (*Mycobacterium*, *Streptococcus*, *Mycoplasma*, *Escherichia coli*, *Helicobacter pylori*), viruses (rubella, Epstein-Barr virus, parvovirus), and superantigens [25; 26; 27].

Although rheumatoid arthritis has a clear genetic component, only about 1 in 25 White individuals with the so-called shared epitope develop rheumatoid arthritis [27]. Even if one monozygotic twin has rheumatoid arthritis, there is only approximately a one in six chance that the other twin will develop the same disease. Thus, other factors in addition to genetics are active as precipitators or triggers of rheumatoid arthritis [27].

Clinical Manifestations

Findings on general physical examination are normal except for an occasional low-grade fever (38°C) and a slightly elevated pulse rate. The characteristic patient with rheumatoid arthritis initially presents with complaints of pain and stiffness in multiple joints. There is prominent and prolonged morning stiffness (lasting more than one hour) that usually begins gradually with fatigue, loss of appetite, widespread muscle aches, and weakness [23; 25; 27].

After this initial presentation, joint pain appears. When the joint is not used for some time, it can become warm, tender, and stiff. After inflammation of the joint, increased synovial fluid is produced and the joint becomes swollen. There is accompanying soft tissue swelling, and joint pain is often felt bilaterally, affecting the fingers, wrists, elbows, shoulders, hips, knees, ankles, toes, and neck [25]. Though the joints are tender, the small joints of the hands and feet are not usually painful when the patient is at rest. Palmar erythema and prominent veins on the dorsum of the hand and wrist indicate increased blood flow. Distal interphalangeal joints are rarely involved. The temperature over the involved joints (except the hip) can be elevated, but there is usually no accompanying erythema. There are limitations in the range of motion, muscle strength, and function around inflamed joints.

In addition, soft, poorly delineated subcutaneous nodules (rheumatoid nodules) are often found in the extensor surface of the forearm. Soft, small lymph nodes are found occasionally in epitrochlear, axillary, and cervical areas [24]. Other symptoms that may present include anemia due to deficits in bone marrow production; eye burning, itching, and discharge; or lung inflammation (pleurisy) [23; 24; 25; 27]. Joint destruction may occur within one to two years after the appearance of the disease.

Rheumatoid arthritis is not solely a disease of joint destruction; it can involve almost all organs. Approximately 18% to 41% of patients with rheumatoid arthritis develop extra-articular manifestations [169; 170]. Rheumatoid arthritis may cause inflammation of the outer cardiac lining (pericarditis) and cardiac muscle (myocarditis), leading to congestive heart failure. In a population-based cohort study, patients with rheumatoid arthritis had a significantly higher risk of cardiovascular disease than those without rheumatoid arthritis [171]. More than half of the patients 50 to 59 years of age and all of those older than 60 years of age with a new diagnosis of rheumatoid arthritis had a more than 10% increased risk of cardiovascular disease within 10 years of rheumatoid arthritis onset.

Pulmonary manifestations are also seen in patients with rheumatoid arthritis, occurring in approximately 30% to 40% of patients. In approximately 10% to 20% of these patients, involvement of the respiratory system is the first manifestation of rheumatoid arthritis [170]. There are several types of potential pulmonary manifestations of rheumatoid arthritis: pleural disease, interstitial pneumonitis, and fibrosis. Pleural effusions and pulmonary rheumatoid nodules are the most common manifestations, along with high rheumatoid factor titers [172; 173; 174]. Pleuritis is more often found in autopsies of patients with rheumatoid arthritis than in living patients. In about 20% of patients, pleuritis develops concurrently with rheumatoid arthritis onset [174]. Although pleuritic pain is not usually a major complaint, the effusions may be large enough to cause dyspnea. Pulmonary fibrosis can either be slowly progressive or result from pulmonary inflammatory disease; on physical exam of the lungs, they present with fine, diffuse, dry rales.

Ocular involvement is another major manifestation of rheumatoid arthritis, usually manifesting as scleritis, development of anterior uveitis, and peripheral ulcerative keratitis (corneal melt) [175; 176]. These disorders are associated with inflammatory cytokines produced by ocular mononuclear cell infiltrates [176; 177].

Osteopenia and osteoporosis are very common extra-articular complications in patients with rheumatoid arthritis [178]. The development of osteopenia in patients with rheumatoid arthritis appears to occur independent of corticosteroid use and is directly linked to elevated levels of the RANK ligand expressed by T cells, which promotes osteoclastic bone resorption [178; 179; 180].

Diagnosis

Rheumatoid arthritis is a clinical diagnosis [181]. As discussed, several laboratory tests are recommended for the diagnosis of rheumatoid arthritis, including rheumatoid factor, ESR, CRP, and anti-CCP antibody [22]. While the results of these tests are relatively sensitive and specific, false positives are possible. In 2010, a multi-biomarker disease activity test, Vectra DA, was introduced. This test uses a unique algorithm to derive a composite score (1 to 100) based on the results of 12 blood pro-

tein biomarkers, including vascular cell adhesion molecule-1, epidermal growth factor, vascular endothelial growth factor A, interleukin-6 (IL-6), TNF receptor type 1, matrix metalloproteinase-1 or collagenase-1, matrix metalloproteinase-3 or stromelysin-1, YKL-40, leptin, resistin, serum amyloid, and CRP [182; 183]. Vectra DA has been independently verified and found to correlate well to disease activity measured with rheumatoid arthritis assessment tools (e.g., Disease Activity Score in 28 joints using the CRP level). The test is validated for use in adults already diagnosed with rheumatoid arthritis but is not intended to diagnose rheumatoid arthritis [184].

There are several other laboratory tests used in the differential diagnosis of rheumatoid arthritis. Complete blood count may reveal mild normochromic and either normocytic or microcytic anemia (hemoglobin 10 g/dL); white blood cell count and differential may reveal thrombocytosis [24; 29]. Although baseline evaluation of renal and hepatic function is not sensitive or specific for rheumatoid arthritis, it is recommended because the findings will guide medication choices.

Popular imaging tests for rheumatoid arthritis include joint ultrasound, MRI, and joint x-rays. Imaging studies may show normal findings or osteopenia and erosions near joint spaces in early disease; wrist and ankle films are useful as baselines for comparison with future studies [24; 185]. Implementing the modern treatment strategy in rheumatoid arthritis (i.e., early initiation and optimal adjustments of aggressive therapies) requires methods for early diagnosis and sensitive monitoring of the disease process.

A number of different medical conditions may be considered in the differential diagnosis of rheumatoid arthritis [181; 186; 187; 188]. These include:

- Connective tissue diseases (e.g., lupus, scleroderma, polymyositis)
- Fibromyalgia
- Hemochromatosis
- Infectious endocarditis
- Lyme arthritis
- Osteoarthritis
- Polyarticular sepsis
- Sarcoidosis
- Thyroid disease
- Viral arthritis

Therapeutic Measures

Rheumatoid arthritis has no known prevention or cure. Lifelong treatment is usually required, including medication, physical therapy, exercise, and possibly surgery. In order to provide the best outcomes, patients should be educated regarding the most appropriate treatment regimens for their disease manifestations, as earlier rheumatoid arthritis diagnosis can

assist in aggressive early treatment for rheumatoid arthritis (when indicated), thereby delaying joint destruction. The 2010 ACR/EULAR Classification Criteria for Rheumatoid Arthritis is now a well-established diagnostic and prognostic tool; as such, guidelines (e.g., the 2016 update of the EULAR Recommendations for the Management of Rheumatoid Arthritis with Synthetic and Biological Disease-Modifying Antirheumatic Drugs) recommend that patients start treatment with a disease-modifying antirheumatic drug (DMARD) immediately following a rheumatoid arthritis diagnosis [189]. Therapeutic goals include preservation of function and quality of life, minimization of pain and inflammation, joint protection, and control of systemic complications, with the ultimate aim being low disease activity or remission [23; 24; 27; 189; 190].

Today, the recommended standard of treatment is a tightly controlled, aggressive strategy tailored to each patient, with modifications to the individual medication regimen to achieve a particular target (remission, or alternatively, low disease activity) in a specific period of time (usually six months) [189; 191]. The “treat-to-target” approach for a patient with early high disease activity and poor prognostic features typically involves initiation of methotrexate and/or another DMARD(s) immediately upon diagnosis [189; 190; 191]. Initial combination therapies with DMARDs, particularly those including a biologic anti-TNF agent, appear to provide earlier clinical improvement and less joint damage progression in patients with early moderate or highly active disease; they can be withdrawn successfully, and fewer treatment adjustments are needed than with initial monotherapies [189; 191; 192; 193; 194]. Patients with active disease are monitored closely (every one to three months), and it is recommended that treatment adjustments be made if there is no improvement at three months (or if the six-month target has not been reached) [189; 191]. Patients with low-to-moderate disease activity or high disease activity without poor prognostic features are typically started on DMARD monotherapy. NSAIDs, glucocorticoids, or COX-2 inhibitors are often used concurrently to treat rheumatoid arthritis-associated joint pain and inflammation. However, they do not alter the disease course and should not be used as single therapy.

Occasionally, surgery is needed to correct severely affected joints. Surgeries serve to relieve joint pain, correct deformities, and modestly improve joint function [23; 24; 27]. The most successful locations of surgery are those performed on the knees and hips [23; 24; 27]. The first surgical treatment performed is a synovectomy, which removes part or all of the joint lining (synovium). This procedure may only provide temporary relief, but it can be effective for patients for whom pharmacologic treatment has not resulted in improvements. Surgeries performed in later-onset disease include total joint replacement with a joint prosthesis. In extreme cases, total knee or hip replacement can have enhanced importance, making the difference between a dependent or independent lifestyle for a patient.

Range-of-motion exercises and individualized exercise programs prescribed by a physical therapist can also delay the loss of joint function. Joint protection techniques, heat and cold treatments, and splints or orthotic devices to support and align joints may be of assistance [23; 24; 27]. Some therapists will use specialized devices to apply deep heat or electrical stimulation to reduce pain and improve joint mobility [23; 24; 27]. Occupational therapists can construct splints for the hand and wrist and teach patients with rheumatoid arthritis how to protect and use their joints most effectively. In addition to physiotherapy, occupational therapists can also show patients with rheumatoid arthritis how to better cope with limitations that can affect their daily tasks at work and at home. For example, many clinicians have recommended frequent rest periods between activities and proper sleeping habits (e.g., 8 to 10 hours of sleep per night) [195].

In addition to the medical management of rheumatoid arthritis, several lifestyle changes may improve symptom severity and decrease the number of flare-ups. The National Institute of Arthritis and Musculoskeletal and Skin Disorders recommends advising patients regarding rest and exercise, use of orthotic devices, stress reduction, and healthful diet [23].

INFECTIOUS DISORDERS

Infectious Arthritis

Infectious arthritis (also known as septic arthritis) is the inflammation of a joint resulting from an invading organism that attacks the synovium and synovial fluid. Viral, bacterial, and fungal infections all predispose susceptible people to arthritis involvement. Pathogens present in the host circulate freely in the bloodstream and become trapped in the richly perfused synovial membrane, leading to inflammation and subsequent degenerative changes. Infectious arthritis is an opportunistic disease that primarily occurs in patients with immunocompromise or who already have joint destruction from another disorder (e.g., rheumatoid arthritis). Early diagnosis and treatment can prevent serious degenerative changes [76; 196].

Patients with infectious arthritis undergo repeated arthrocentesis, which can be stressful. Additional treatment will depend on the underlying pathogen, with antibiotics, antiviral, or antifungals prescribed as appropriate [76; 196].

The nurse should be available to both the patient and family for psychological support, physical care, health education, and monitoring the patient’s response to therapy. The control of pain and protection of the involved joint or joints are priorities of nursing management [76; 196].

Therapeutic and Nursing Measures

The patient history is key to diagnosis, and nurses should be careful to obtain a complete and accurate history. This should include any recent viral (e.g., parvovirus, alphavirus, hepatitis, Epstein-Barr virus) and bacterial (e.g., *Streptococcus pneumoniae*) infection. In young, sexually active patients, the most com-

mon causative pathogen is *Neisseria gonorrhoea*. For patients who develop infectious arthritis following trauma, puncture wounds, or injection drug use, *Pseudomonas aeruginosa* is the most likely cause [18; 197].

It is important that the pathogen responsible for the infectious arthritis be identified and treatment begun as quickly as possible to prevent joint destruction. Isolating the organism will guide in the selection of intravenous antimicrobial therapy and the level of aggressiveness needed to control the infection. Pathogens are identified through the aspiration of synovial fluid, synovial fluid cultures, and synovial biopsy. Empiric therapy is started after joint aspiration is complete and cultures are obtained [76]. Other therapeutic measures for infectious arthritis include surgical excision of the affected synovium in instances where destruction of the joint cartilage, tendons, or both appears imminent [76].

To protect the intra- and extra-articular structure from future damage and reduce the patient's discomfort, the involved joint should be immobilized during the acute stage. However, after two to three days, aggressive physical therapy is recommended to prevent long-term damage and disability [18].

The involved joint should be assessed frequently for drainage and any change in condition. Sterile technique should be maintained with any dressing changes [18]. For patients receiving parenteral fluids, intake and output should be measured and documented accurately. Laboratory test results will be monitored daily, especially the results of culture and sensitivity tests [18].

Patient education should include instruction about range-of-motion exercises to maintain joint mobility; dressing change techniques and wound care, if appropriate; and adherence to prescribed medications. The patient should be advised of symptoms and signs of repeated infection (e.g., increased pain, fever, swelling, redness, drainage) and to avoid any trauma to the joint [18].

NEOPLASTIC DISORDERS

In this section, the discussion of masses and tumors of the joints and surrounding structures will be limited to the benign and malignant lesions generally included in a differential diagnosis for arthritis.

Masses and Benign Tumors of the Joint

Patients with a benign lesion of the joint may experience years of intermittent minor problems with the involved joint, with a history of discomfort and joint instability. Because the symptoms of a benign tumor can remain innocuous for long periods, joint damage may result prior to diagnosis. No matter the extent of damage, joint surgery is required to resolve the issue and prevent further deterioration [37].

Lipoma

A lipoma of a joint is a lobulated fatty mass. Lipomas develop frequently in the elbow or knee joint of patients with osteoarthritis [37]. In many cases, patients seek medical attention when the involved joint begins locking or when pain, decreased motion, or an effusion occurs. Effusion aspirate is clear, and x-rays are non-diagnostic [37].

Hemangioma

Hemangiomas are rare vascular tumors often associated with arteriovenous malformations of skin vascular disease. They tend to affect younger individuals, often teenage girls who have been symptomatic since childhood. The knee is the most commonly involved joint [37].

In patients with joint hemangioma, there is a history of episodic, unilateral "doughy" joint swelling; pain; limitation of motion; and locking, buckling, or both. Aspiration of the joint repeatedly produces serosanguineous fluid in the absence of trauma. X-rays early in the course of the hemangioma may appear normal; ultrasound is often more helpful. With more advanced disease, enlarged epiphyses, joint narrowing, and enlargement of the intercondylar notch layer may be visualized [37].

Surgical removal yields good results in the treatment of a localized hemangioma. If accessible, therapeutic embolization of a major feeder vessel may be effective as well. Because of the vascular nature of this tumor, a diffuse form may involve the entire joint capsule and make resection impossible. For these patients, radiation therapy is the usual treatment [37].

Synovial Chondromatosis

Synovial chondromatosis is a condition of unknown etiology in which numerous cartilaginous nodules form. These nodules involve the joint, bursae, and in some cases the tendon sheaths of a knee, hip, elbow, shoulder, or ankle. It is a self-limiting disease that most frequently affects young and middle-aged men [37].

Synovial chondromatosis has an insidious onset, and many years often pass before the patient seeks evaluation for the problem. Presenting symptoms usually consist of swelling, pain, stiffness, limitation of motion, and joint locking. Particularly with synovial chondromatosis, the patient experiences joint crepitation or a grating sensation from the multiple intra-synovial nodules. X-ray findings may demonstrate calcified, free-floating bodies in the synovium [37].

Excision of the involved synovium and removal of all loose bodies is the treatment of choice. Surgical therapy has a good prognosis, although the condition may recur if removal is incomplete [37].

Pigmented Villonodular Synovitis

Pigmented villonodular synovitis is a condition in which the synovial lining cells have a marked proliferation that results in the appearance of numerous villi and folds. The formation of the finger-like projections can affect not only the synovial lining but also the tendon sheath, bursa, and bone. Unilateral involvement of the knee, hip, ankle, or elbow joint of young adults is most common [37].

There are two forms of pigmented villonodular synovitis: localized and diffuse. The diffuse type causes pain and mild, episodic joint swelling over a period of months to years. Occasionally, the patient may note an acutely painful, warm, swollen joint with limited motion. Repeated aspirations of joint fluid yield dark serosanguineous fluid in the absence of trauma [37].

The localized type of pigmented villonodular synovitis occurs in either the medial or lateral knee compartment as a solitary nodule. It, too, may begin with episodic pain and mild swelling and can be misdiagnosed as torn meniscus. Serosanguineous fluid is rarely aspirated with this type of lesion [37]. Imaging studies may show soft tissue density, but angiographies are more useful because the vascularity of these enable the condition to be diagnosed [37].

Surgical resection is the treatment for both types of pigmented villonodular synovitis. The localized form is usually cured with simple excision. With the diffuse type, lesions may recur if synovectomy is incomplete [37].

Specific Nursing Measures

During the diagnostic phase, the health history is important. The nurse should concentrate on any significant joint trauma and inquire regarding any past history of arthritis. The patient should be asked for a detailed description of swelling, pain, limitation of motion, and/or joint instability [7].

In cases of significant joint instability, a cane or crutches may be necessary. Patients should also receive education on appropriate pain management measures, including thermotherapy, over-the-counter medications, and biofeedback. In most cases, the patient will also require education to prepare for surgery and postoperative recovery [7].

Postoperative care includes monitoring for wound drainage or other signs of infection, dressing changes (as necessary), and splinting. Discharge planning includes instruction in the care of the surgical site, activity restrictions, and follow-up appointments [7].

Malignant Tumors of the Joint

Malignant tumors of the joint are rare. However, if a patient has a slow-growing monoarticular mass, malignancy should be suspected [198].

Synovial Sarcoma

Though it is the most common primary tumor of the joint, synovial sarcoma is rare. This malignancy can appear at any age, although it seems to predominate in young adults. The growth generally appears on a lower extremity, but synovial sarcomas can also develop in an upper extremity, the neck, or the chest [198].

Patients with synovial sarcoma often present with a slow-growing mass that may have been present for months to years, depending on how deeply seated it is in tissue. Pain may be present, or the patient may have a vague sensation of discomfort over the involved area. There may also be localized swelling. In cases involving the neck, tumor invasion may produce hoarseness, dysphagia, or dyspnea [198].

As with most cancer, survival time is dependent on the size of the tumor, site in the body, and age at diagnosis. The five-year survival rate is approximately 60%; this increases to 75% in patients 30 years of age or younger. If the tumor is in an extremity, five-year survival is about 65%; if the tumor is present in the trunk, the rate decreases to 40% [199].

The goals of treatment for synovial sarcoma are to eliminate the tumor, preserve a functional limb, and minimize mortality and morbidity. Preoperative chemotherapy and radiation therapy may be undertaken to decrease the size of the tumor [198]. Wide surgical resection is typically undertaken, followed by continued radiation therapy.

Clear Cell Sarcoma

Clear cell sarcoma is a rare tumor that involves tendons rather than joint spaces. It can occur in any age group and usually is found in an extremity, particularly the foot. However, it can develop in the trunk, head, genitals, stomach, and intestines. Because of this lesion's location and its predisposition to metastasis, it can be difficult to remove entirely, making treatment complicated and prognosis poor [198]. The average age at diagnosis is 25 years [200].

The primary treatment of clear cell sarcoma is radical resection of the tumor. In some cases, an extremity may be amputated. Preoperative and postoperative radiation therapy are employed. In some cases, chemotherapy may be used, but it is not particularly effective [200].

TRAUMATIC DISORDERS OF THE JOINT

Permanent structural changes may occur in a joint as a result of cartilage and capsular tears, detachment of menisci, hemorrhagic effusions, articular fractures, or repetitive trauma. Because of the realignment of involved bone, bursa, and tendons, a mechanical deterioration of articular cartilage results in osteoarthritis [201; 202].

Traumatic arthritis may result from unexpected force (e.g., sports, motor vehicle accidents) or from repetitive trauma—a chronic injury resulting from repeated smaller stresses to a joint

through vibrations, blows, abnormal strain, or position. Injury resulting from repetitive stress is often related to occupation and lifestyle, for example, the stress placed on the metatarsophalangeal joints of a ballet dancer or the knees of a jogger. Over time, repetitive trauma may realign the joint and lead to the same result as an acute injury [201; 202].

The patient with traumatic arthritis must make lifestyle changes. In order to continue participating in chosen sports or occupations, patients may require braces, splints, or special equipment; in some cases, they may need to halt participation or seek accommodations in their workplace. Although these options are effective in halting the progression of traumatic arthritis, they are often undesired or impossible for patients. The responsibility of the nurse is to provide accurate information about the alternatives available [201; 202].

Nurses also play an important role in the therapeutic and preventive care of traumatic disorders. If they are the first person on the scene of an accident, nurses may be able to prevent any residual damage by splinting, elevating the area, and not allowing weight on the area to protect the joint. It may be possible to identify tasks that expose employees to repetitive trauma; employees then can rotate through the jobs rather than be assigned permanently to potentially harmful tasks [201; 202].

Joint Effusions

Joint effusions can occur as a result of simple trauma or secondary to fractures, internal derangements, or severe sprains. Within 24 hours after a blow to the joint, synovial fluid accumulates. If blood vessels in the synovium are broken, hemarthrosis also occurs. The knee is most commonly affected by this injury, although it can occur in other joints as well [75].

Clinical Manifestations

In simple cases of traumatic synovitis, joint swelling with mild pain occurs. Aspiration of the joint produces clear fluid with elevated protein content and decreased viscosity. Hemarthrosis, which usually develops within 15 minutes to 2 hours after the trauma, is usually more painful than clear effusion and is accompanied by low-grade fever. Diagnosis of traumatic synovitis is primarily by physical examination, but x-ray examination is done to rule out fracture [75].

Therapeutic Measures

Immediately following injury, patients should be advised to apply cryotherapy (e.g., ice) for 30 minutes; this can be repeated up to four times per day to reduce swelling and relieve pain. After the first 24 hours, patients should switch to moist heat to relax surrounding muscles and reduce pain. If fluid accumulates in the joint, repeated joint aspirations may be necessary. Compression dressings applied to the joint, along with limited weight bearing, may be used, depending on the severity of the injury [75].

Dislocation and Subluxation

Dislocation is the complete displacement of a joint's articulating surfaces following trauma. Partial displacement of the articulating surfaces results in subluxation. Both subluxations and dislocations can damage soft tissues, nerves, or blood vessels if not attended to promptly. The joints most often affected are shoulders, wrists, elbows, fingers, hips, knees, and toes [75].

Clinical Manifestations

After injury, the joint appears deformed; it is tender, and motion is limited. The involved extremity may be visibly shortened. Joint pain may be intense, especially if articular surface fractures are present. With immediate treatment, there is a good prognosis. However, bone necrosis can result if reduction of the subluxation or dislocation is delayed [75]. Diagnosis is made through physical examination and patient history, with x-rays taken to evaluate joint displacement and to determine whether fractures are present [75].

Therapeutic Measures

The longer the delay in correcting a joint displacement, the more difficult the procedure becomes because of edema and muscle spasms. Two types of procedures can correct this injury. Closed reduction is manual traction done under local or general anesthesia. The pain associated with this procedure can be intense, and pain management techniques (including strong analgesics) are necessary. If muscle spasms are an issue, tranquilizers and/or muscle relaxants may be administered. Open reduction is done when wire fixations of the joint or repair of torn ligaments is also necessary [75].

If present at the site of injury, the joint should be splinted—even if crooked—to prevent further damage. Cold compresses can be applied to decrease pain and swelling [35]. The area distal to the injury should be observed for evidence of vascular damage (e.g., pallor, absent pulse, abnormal coolness) and nerve damage (e.g., paresthesia, paralysis) [35].

If analgesics, muscle relaxants, or tranquilizers are administered, it is important to monitor the patient's respiratory status. Any dressings or casts should be checked for pressure that may impair blood flow. Patients should receive education on gradual mobilization and return to activities [7].

CASE STUDIES

SYSTEMIC LUPUS ERYTHEMATOSUS

Patient A is a woman, 29 years of age, with two small children. She presents to her primary care provider with complaints of rashes developing on her arms and legs whenever she spends time in the sun. She also reports several small patches of hair loss on her head that she attributes to the stress of new motherhood and to a recent trip and her fear of flying. She reports a lack of energy, being easily fatigued, and always needing to

PATIENT A'S FIRST PHYSICAL EXAM RESULTS	
Parameter	Findings
General appearance	Significantly underweight, with a decrease in weight of 23 pounds since last exam one year prior Height: 5 feet 5 inches (165.5 cm) Weight: 108 pounds (49 kg)
Skin and nails	Multiple rash-like lesions on sun-exposed areas of the body, primarily on the arms and legs Slightly jaundiced
Head and nose	Nares clear Oropharynx benign and without obvious lesions Mucous membranes moist
Eyes	Some yellowing within the sclera Pupils equal, round, reactive to light and accommodation Conjunctiva normal No retinal exudates
Ears	Tympanic membranes intact
Neck	Supple No signs of lymphadenopathy, jugular vein distension, or thyromegaly
Chest	Clear to auscultation throughout Equal air entry bilaterally No wheezing or crackles Chest resonant on percussion
Abdomen	Soft and nontender Active bowel sounds No masses or organ enlargement
Extremities	No cyanosis, clubbing, or edema Rash-like lesions present
Genitourinary system	Normal female
Neurologic status	Alert and oriented Deep tendon reflexes 2+ with symmetrical flexor plantar responses No focal deficits noted
Cardiovascular system	Regular rate and rhythm Prominent S ₁ and S ₂
Vital Signs	
Blood pressure	110/70 mm Hg
Temperature	99.8° F
Heart rate	70 beats per minute with regular rhythm
Respiratory rate	15 breaths per minute
Source: Author	

Table 2

nap during the day. Patient A also reports mild pain in her fingers and elbows but attributes the joint discomfort to carrying for the children. She states that these problems have been ongoing for approximately four months.

Medical History

Patient A has no known allergies and takes no prescription or over-the-counter medication aside from occasional naproxen for joint pain and antacid for heartburn. She neither smokes nor drinks alcohol. Her youngest child is 2 years of age, and she reports unremarkable childbirths and postpartum periods. Aside from the current complaints, the patient's medical history is unremarkable.

She has four brothers and three sisters. The family history indicates an older sister with rheumatoid arthritis, an aunt with pernicious anemia, and mother with hyperthyroidism.

Assessment and Diagnosis

The primary care provider conducts a full physical assessment (**Table 2**). Several laboratory tests are ordered, with the following results:

- Hematocrit (HCT): 23%
- Red blood cell (RBC) count: 3.5 million cells/mcL
- White blood cell (WBC) count: 5,500 cells/mcL
- Platelets: 350,000 cells/mcL
- ESR: 25 mm/hour
- Urinalysis: Normal
- ANA: 1:640
- Anti-DNA antibody test: Elevated
- Complement assay: Decreased C3 level at 43 mg/dL and decreased C4 level at 14 mg/dL

Further, a tissue biopsy of one of the lesions is taken and reveals vasculitis (i.e., white blood cells within the walls of blood vessels).

Based on the results of the assessment and laboratory studies, Patient A is diagnosed with SLE.

Management

A one-month course of prednisone with tapered doses is prescribed. Nabumetone, an anti-inflammatory, is added to the regimen prior to the prednisone being weaned off. After one month of treatment, all signs and symptoms of lupus have resolved.

However, 13 years later, Patient A again presents to her primary care provider, this time with complaints of a productive cough and transient stiffness and pain in her hands and feet (migratory polyarthritis). She is afraid that she is developing rheumatoid arthritis like her sister. The provider conducts a physical examination (**Table 3**) and is concerned that the patient may be showing signs of pneumonia. A chest x-ray revealed mild pulmonary edema but no white blood cell infiltrates in the terminal airways. Laboratory tests reveal:

- HCT: 43%
- Platelet: 330,000 cells/mcL
- WBC count: 1,200 cells/mcL
- Urinalysis: Within normal limits

Patient A is diagnosed as experiencing a lupus flare and is prescribed a one-month course of prednisone along with a 10-day course of antibiotics to prevent pneumonia. Within three months, all signs and symptoms have resolved.

Five years later, Patient A returns to her primary care provider complaining of fatigue, anorexia, weight loss (25 pounds in the last four months), and significant swelling in her abdomen,

face, and ankles. The nurse practitioner notes a “butterfly-shaped” rash present across the bridge of the patient’s nose and cheeks. Blood tests reveal an HCT of 24% and a WBC count of 2,400 cells/mcL. A dipstick examination of the urine reveals an abnormal protein concentration, and microscopy indicates the presence of significant numbers of red and white blood cells. A 24-hour urine protein collection reveals excretion of 2.5 g protein in 24 hours.

Study Questions

1. What is the significance of the patient’s family history?
2. Is this patient underweight, normal weight, overweight, or obese?
3. What underlying pathologic process is responsible for Patient A’s hair loss? What is the relevance of the abnormal ESR?
4. Vasculitis in lupus results from the trapping of antigen antibody complexes in blood vessel walls followed by an intense inflammatory response to the immune complexes. Why is prednisone effective in relieving vasculitis?
5. What is the most likely cause of jaundice in this patient?
6. What pathophysiology underlies lymph node enlargement in this patient?
7. The patient’s WBC differential was: 75% neutrophils, 15% lymphocytes, 5% monocytes/macrophages, 4% eosinophils, and 1% basophils. Which one of these white blood cell types has been specifically targeted by the patient’s immune system?
8. Why was Patient A experiencing swelling throughout her body?

LOW BACK PAIN

Patient B is a woman, 35 years of age, who has worked as a housekeeper for the past 10 years. She is 5 foot 3 inches in height with a weight of 178 pounds. She presents to her primary care provider with complaints of low back pain. She reports having had this pain intermittently for several years; however, for the past two days, it has been worse than ever. The recent exacerbation started after vacuuming a rug (i.e., pulling and twisting at the waist). Patient B reports that the pain is primarily on the right lower side and radiates down her posterior right thigh to her knee; it is not associated with any numbness or tingling. The pain can be relieved by lying flat on her back with her legs slightly elevated and is lessened somewhat when she takes ibuprofen 400 mg. Except for moderate obesity and difficulty maneuvering onto the examination table because of pain, the patient’s examination is fairly normal. The only abnormalities noted are a positive straight leg raise test, with raising the right leg eliciting more pain than the left. Her strength, sensation, and deep tendon reflexes in all extremities are normal.

PATIENT A'S SECOND PHYSICAL EXAM RESULTS	
Parameter	Findings
General appearance	Healthy and calm White woman Height: 5 feet 5 inches (165.5 cm) Weight: 131 pounds (59.5 kg)
Skin and nails	No lesions or abnormalities noted
Head and nose	Nares clear Oropharynx irritated but without obvious lesions Mucous membranes moist
Eyes	Pupils equal, round, reactive to light and accommodation Conjunctiva normal
Ears	Tympanic membranes intact
Neck	Supple Lymph nodes slightly enlarged
Chest	Auscultation reveals abnormal lung sounds (bronchitis) No wheezing, but some crackles
Abdomen	Soft and nontender Active bowel sounds No masses or organ enlargement
Extremities	No cyanosis, clubbing, or edema Axillary lymph nodes swollen
Genitourinary system	Normal female Inguinal lymph nodes slightly enlarged
Neurologic status	Alert and oriented Deep tendon reflexes 2+ with symmetrical flexor plantar responses No focal deficits noted
Cardiovascular system	Regular rate and rhythm Prominent S ₁ and S ₂
Vital Signs	
Blood pressure	140/90 mm Hg
Temperature	100.0° F
Heart rate	105 beats per minute with regular rhythm
Respiratory rate	15 breaths per minute
Source: Author	

Table 3

Study Questions

1. What is the patient's likely diagnosis?
2. How will the patient be treated?

RHEUMATOID ARTHRITIS

Patient C is a woman, 50 years of age, who presents to her primary care provider for her annual exam. She reports having been very tired for the past month and also experiencing stiffness, pain, and swelling in multiple joints. She states, "I ache all over, and I have pain in different places all the time. One day it is in my right shoulder, the next day in my right wrist, and the following day my left wrist. I'm stiff everywhere when I get up in the morning or if I sit for any length of time. And I feel so tired, like I have a case of the flu that won't go away."

PATIENT C'S PHYSICAL EXAM RESULTS	
Parameter	Findings
General appearance	Pleasant and alert, but appears very tired and is in moderate acute distress from joint pain Height: 5 feet 4 inches (162.5 cm) Weight: 140 pounds (63.5 kg)
Skin and nails	Intact, warm, pink, and dry No rashes Normal turgor
Head and nose	Head atraumatic
Eyes	Pupils equal, round, reactive to light and accommodation Normal funduscopic examination
Ears	Tympanic membranes intact
Neck	Supple with no jugular vein distention or thyromegaly No bruits Mild lymphadenopathy bilaterally
Chest	Clear to auscultation and percussion No lumps, dimpling, discharge, or discoloration noted in breast exam
Abdomen	Soft, non-tender, and non-distended Positive bowel sounds throughout No superficial veins or organomegaly
Extremities	No clubbing or ankle edema Hands: Swelling of the 3rd, 4th, and 5th proximal interphalangeal joints bilaterally. Pain in the 4th and 5th metacarpophalangeal joints bilaterally. Poor grip strength bilaterally. Wrists: Good range of motion. Fixed nodule at pressure point on left side. Elbows: Good range of motion. Fixed nodule at pressure point in right side. Shoulders: Pain and decreased range of motion bilaterally. Hips: Good range of motion. Knees: Pain, significant edema, and decreased range of motion bilaterally. Feet: No edema. Full plantar flexion and dorsiflexion and full pedal pulse bilaterally.
Genitourinary system	Last menstrual period 16 months ago Normal pelvic exam
Neurologic status	Alert and oriented Cranial nerves II–XII intact Muscle strength 5/5 in upper extremities and 4/5 lower extremities Deep tendon reflexes 2+ in biceps, triceps, and patella
Cardiovascular system	Regular rate and rhythm Normal S ₁ , S ₂ , no S ₃ or S ₄ No murmurs, rubs, or gallops
Vital Signs	
Blood pressure	125/80 mm Hg
Temperature	100.0° F
Heart rate	80 beats per minute with regular rhythm
Respiratory rate	15 breaths per minute
<i>Source: Author</i>	

Table 4

The patient has been diagnosed with hypothyroidism in the past, for which she is taking levothyroxine. She is also prescribed venlafaxine to treat major depressive disorder, and she indicates that her mood has been good, despite the fatigue. She is also taking an over-the-counter multivitamin and calcium supplement. Patient C reports rarely using alcohol and never smoking. There is no family history of autoimmune disorders.

Assessment and Diagnosis

The primary care provider does a complete physical exam (**Table 4**) and orders laboratory tests. The laboratory blood test results are:

- Sodium: 140 meq/L
- ANA: Negative
- HCT: 43%
- Uric acid: 2.9 mg/dL
- Potassium: 3.7 meq/L
- ESR: 38 mm/hour
- WBC count: 15,100 cells/mcL
- Cholesterol: 189 mg/dL
- Chloride: 104 meq/L
- Creatinine: 1.0 mg/dL
- Platelets: 270,000 cells/mcL
- Albumin: 4.0 g/dL
- Bicarbonate: 23 meq/L
- Blood glucose: 94 mg/dL
- RBC count: 4.7 million cells/mcL
- Thyroid stimulating hormone (TSH): 1.7 mcU/mL
- Blood urea nitrogen: 18 meq/L
- Hemoglobin: 14.9 g/dL
- Calcium: 8.8 mg/dL
- Rheumatoid factor: Positive

A urinalysis is performed and is normal, with no RBCs, WBCs, or protein. A chest x-ray finds no fluid, masses, infection, or cardiomegaly. An x-ray of the hand shows soft tissue swelling and bone demineralization but no erosions. Synovial fluid removed from the left knee (7.4 mL) is cloudy and pale yellow in appearance; analysis indicates 14,000 white blood cells/mcL (primarily neutrophils) and a glucose level of 60 mg/dL.

Based on these findings, Patient C is diagnosed with rheumatoid arthritis and referred to a rheumatologist for follow-up.

Study Questions

1. Which of Patient C's vital signs is consistent with a diagnosis of rheumatoid arthritis and why?
2. Are there any other abnormal findings from the patient's physical exam that are consistent with a diagnosis of rheumatoid arthritis?
3. What is the association between the fixed nodules at pressure points on the left wrist/right elbow and a diagnosis of rheumatoid arthritis?
4. Why is it reasonable that this patient has no stiffness, pain, or swelling in the DIP joints of the fingers?
5. Which of these patient's laboratory test results are consistent with a diagnosis of rheumatoid arthritis?
6. In terms of the progression of the disease, what do the results of the hand x-ray suggest?
7. Which findings in the examination of the synovial fluid are consistent with a diagnosis of rheumatoid arthritis?
8. What causes limitation of joint motion that occurs early in the clinical course of rheumatoid arthritis? What causes limitation of joint motion that occurs late in the clinical course of rheumatoid arthritis?

CONCLUSION

With knowledge of the structures and function of the muscles, joints, and connective tissue and the dynamic pathology that intrudes and impedes normal function, nurses can readily provide quality and often life-saving actions. An awareness of why symptoms appear leads to quicker reporting to physicians of changes in the patient's condition. Nurses can also perform immediate interventions based on standing orders and recognition of what needs to be done in order to provide safe quality care. This knowledge changes what could be only technical care to professional care through the use of decision-making skills built upon the knowledge of pathophysiology.

SELF-ASSESSMENT QUESTIONS

#38950 PATHOPHYSIOLOGY: MUSCLES, JOINTS, AND CONNECTIVE TISSUES

After reviewing the course, complete the following Self-Assessment Questions. Receive immediate feedback by reviewing the Study Guide provided on pages 115–117. Please record the number of questions you correctly answered on the Evaluation. The Evaluation must be completed in order to receive credit for this course.

This 15 contact hour activity must be completed by September 30, 2025.

- The prime mover is the muscle that**
 - contracts to produce movement.
 - works to assist in performing the movement.
 - remains inert during movement to provide stability.
 - works opposite to synergists by relaxing during their contraction.
- A bursa is a**
 - connective tissue covering.
 - fibrous connective tissue that connects bones to one another.
 - cord of connective tissue that attaches muscles to the periosteum of the bones.
 - fluid-filled sac that facilitates motion of structures that move against each other.
- All of the following statements regarding muscle fibers are true, EXCEPT:**
 - Type 1 fibers have a high excitability.
 - Type 1 fibers have a large cell diameter.
 - Type 2 fibers are numerous in quantity.
 - Type 2 fibers have a fast conduction velocity.
- What musculoskeletal structure is most frequently influenced by degenerative disease?**
 - Joints
 - Fasciae
 - Muscles
 - Tendons
- Heberden nodes may be noted on the distal interphalangeal joint of patients with**
 - osteoporosis.
 - osteoarthritis.
 - rhabdomyolysis.
 - systemic lupus erythematosus.
- The elbow's normal flexion is**
 - 0°.
 - 90°.
 - 160°.
 - 240°.
- Which of the following statements regarding the antinuclear antibody (ANA) test is FALSE?**
 - The ANA is the most commonly used autoantibody test.
 - The ANA is the most specific and sensitive test for lupus.
 - A positive ANA test alone indicates a conclusive diagnosis of lupus.
 - Nearly all patients with lupus (97%) have a positive ANA blood test.
- According to the American Academy of Physical Medicine, ultrasound is an essential component in the diagnosis of all of the following, EXCEPT:**
 - Nerve entrapments
 - Rheumatoid arthritis
 - Tendinopathies/tendon tears
 - Acute or chronic muscle injury
- The most common ocular symptom of Marfan syndrome is**
 - uveitis.
 - myopia.
 - glaucoma.
 - retinal detachment.
- Patients with Ehlers-Danlos syndrome who become pregnant are at risk for**
 - pre-eclampsia.
 - uterine rupture.
 - postpartum depression.
 - low birth weight infants.

Self-Assessment questions continue on next page →

11. Which of the following should be included in the differential diagnosis of bursitis and tendonitis?
 - A) Gout
 - B) Infection
 - C) Rheumatoid arthritis
 - D) All of the above
12. Which of the following is recommended for the management of bursitis and tendinitis?
 - A) Opioid analgesics
 - B) Long-term immobilization
 - C) Warm compresses to the affected area
 - D) Physical therapy and structured exercise after the initial period of rest
13. The average age at diagnosis of polymyalgia rheumatica is
 - A) 5 years of age.
 - B) 25 years of age.
 - C) 50 years of age.
 - D) 70 years of age.
14. Which type of lupus mainly affects the skin?
 - A) Drug-induced lupus
 - B) Systemic lupus erythematosus (SLE)
 - C) Peripheral lupus erythematosus (PLE)
 - D) Cutaneous lupus erythematosus (CLE)
15. Which of the following is thought to be the prime cause of SLE?
 - A) Hormones
 - B) Environment
 - C) Genetic factors
 - D) Immune dysregulation
16. The most universal symptom of lupus is
 - A) rash.
 - B) fever.
 - C) fatigue.
 - D) joint pain.
17. Which of the following drug classes is involved in the treatment of lupus?
 - A) NSAIDs
 - B) Opioids
 - C) Antivirals
 - D) Antibiotics
18. The most frequent presentation in systemic scleroderma is the clinical triad of
 - A) anemia, kidney failure, and hypothermia.
 - B) fatigue, skin changes, and esophageal hypermotility.
 - C) skin changes, Raynaud phenomenon, and esophageal hypomotility.
 - D) mucosal membrane inflammation, hyperthermia, and joint swelling.
19. For patients with systemic scleroderma who experience digital ulcers, the recommended treatment is
 - A) NSAIDs.
 - B) methotrexate.
 - C) intravenous iloprost.
 - D) atovaquone-proguanil.
20. A sprain is an injury to a ligament caused by
 - A) muscle overexertion.
 - B) repetitive motion over time.
 - C) Fracture of the attached bone.
 - D) forcing a joint beyond its normal range of motion.
21. The most common sprains affect the
 - A) knee.
 - B) wrist.
 - C) ankle.
 - D) shoulder.
22. The standard of care for patients with rhabdomyolysis is
 - A) mannitol.
 - B) fluid therapy.
 - C) temperature stabilization.
 - D) extracorporeal blood purification.
23. Which of the following statements regarding psoriatic arthritis is TRUE?
 - A) The disease is always symmetrical.
 - B) All patients eventually develop sacroiliitis.
 - C) Nodules are present with psoriatic arthritis.
 - D) The manifestations of psoriatic arthritis vary from patient to patient.
24. Urate-lowering therapy should be initiated in patients with gout and
 - A) one or more tophi.
 - B) no history of urolithiasis.
 - C) stage 1 chronic kidney disease.
 - D) fewer than two attacks per year.

25. Which of the following is a nonmodifiable risk factor for developing low back pain?
- A) Smoking
 - B) Female sex
 - C) Job dissatisfaction
 - D) Higher body mass index (BMI)
26. The majority of low back pain cases are
- A) nonspecific.
 - B) myofascial pain.
 - C) lumbar spinal stenosis.
 - D) lumbosacral radiculopathy.
27. Which of the following statements regarding sacroiliac (SI) joint syndrome is TRUE?
- A) Pain usually extends below the knee.
 - B) The onset of SI joint pain is usually abrupt.
 - C) Pain is intensified by attempts to walk downstairs.
 - D) It typically manifests as localized pain in the lower back or upper buttock area that overlies the SI joint.
28. Which of the following is considered a “red flag” when assessing the patient with chronic low back pain?
- A) Prior history of cancer
 - B) Fear avoidance beliefs
 - C) Depression and anxiety
 - D) Reluctance to self-manage
29. Which of the following is a management goal for chronic low back pain?
- A) Improve functional status
 - B) Minimize pain and disability
 - C) Facilitate restoration of normal activity
 - D) All of the above
30. Scoliosis
- A) is a postural curvature of the spine.
 - B) is more common in boys than girls.
 - C) most commonly occurs in the cervical area of the spine.
 - D) is most commonly the result of idiopathic causes that appear with increased growth during adolescence.
31. Which of the following statements most accurately defines carpal tunnel syndrome?
- A) Pain and abnormal nerve sensations in the neck, shoulder, arm, and/or hand
 - B) Painful disorder of the wrist and hand that occurs when the median nerve becomes compressed
 - C) Injury to the nerves, muscles, and tendons of the wrist and hand due to the repetitive use of vibrating hand tools
 - D) Medial elbow pain, hand fatigue, and sensations in the fourth and fifth fingers resulting from ulnar nerve damage
32. As an intervention for carpal tunnel syndrome, splinting
- A) is of clear benefit as postoperative rehabilitation.
 - B) has only been shown to improve patient function after 16 weeks.
 - C) should be begun immediately after routine carpal tunnel release surgery.
 - D) may be particularly helpful when weighing the risks of surgery versus the benefits.
33. The primary symptom of osteoarthritis of the knee is pain, especially
- A) with extension of the extremity.
 - B) accompanied by stiffness after use.
 - C) present at rest and interfering with sleep.
 - D) with weight-bearing exercise or activity, that improves with rest.
34. Which of the following is TRUE about the etiology of rheumatoid arthritis?
- A) It is likely an autoimmune disease.
 - B) It is believed that infectious, genetic, and hormonal factors may be contributing factors.
 - C) If one monozygotic twin has rheumatoid arthritis, there is a 1 in 2 chance that the other twin will develop the same disease.
 - D) Both A and B
35. Which of the following conditions should be included in the differential diagnosis of rheumatoid arthritis?
- A) Bursitis
 - B) Influenza
 - C) Sarcoidosis
 - D) Rhabdomyolysis

Self-Assessment questions continue on next page →

36. Which of the following best describes the treatment of rheumatoid arthritis?

- A) Rheumatoid arthritis does not require lifelong treatment.
- B) Rheumatoid arthritis is a curable disease that can be treated with a set pharmaceutical regimen.
- C) Patients with suspected rheumatoid arthritis should be referred within one month of presentation.
- D) The earlier a patient's rheumatoid arthritis disease is diagnosed, the earlier aggressive treatment for rheumatoid arthritis can be started, thereby delaying joint destruction.

37. In young, sexually active patients with infectious arthritis, the most common causative pathogen is

- A) hepatitis.
- B) *Neisseria gonorrhoea*.
- C) *Chlamydia trachomatis*.
- D) *Pseudomonas aeruginosa*.

38. Hemangiomas are

- A) common vascular tumors.
- B) most likely to affect older men.
- C) often associated with arteriovenous malformations of skin vascular disease.
- D) typically identified in the joints of the upper extremities (e.g., wrist, elbow).

39. Which of the following statements regarding synovial sarcoma is TRUE?

- A) The growth generally appears on a lower extremity.
- B) Though it is the most common primary tumor of the joint, synovial sarcoma is rare.
- C) In cases involving the neck, tumor invasion may produce hoarseness, dysphagia, or dyspnea.
- D) All of the above

40. Joint effusions

- A) are primarily diagnosed by MRI.
- B) most commonly occur in the elbow.
- C) occur only secondary to fractures, internal derangements, or severe sprains.
- D) result in synovial fluid accumulation within 24 hours after a blow to the joint.

STUDY GUIDE

COURSE #38950 PATHOPHYSIOLOGY: MUSCLES, JOINTS, AND CONNECTIVE TISSUES

This Study Guide has been included to provide immediate feedback about your achievement for the course learning objectives. The correct answers for the Self-Assessment Questions and their locations within the text are indicated below.

- The prime mover is the muscle that**
 - contracts to produce movement. (*Muscles, Joints, and Connective Tissues: Structural and Functional Inter-Relationships; Structure and Function of Skeletal Muscles; Producing Skeletal Movement*)
- A bursa is a**
 - fluid-filled sac that facilitates motion of structures that move against each other. (*Muscles, Joints, and Connective Tissues: Structural and Functional Inter-Relationships; Other Connective Tissue Structures*)
- All of the following statements regarding muscle fibers are true, EXCEPT:**
 - Type 1 fibers have a large cell diameter. (*The Process of Skeletal Muscle Contraction*)
- What musculoskeletal structure is most frequently influenced by degenerative disease?**
 - Joints (*Pathophysiologic Influences and Effects; Degenerative Changes*)
- Heberden nodes may be noted on the distal interphalangeal joint of patients with**
 - osteoarthritis. (*Nursing Assessment: Establishing the Data Base; Objective Data; Structural Changes*)
- The elbow's normal flexion is**
 - 160°. (*Nursing Assessment: Establishing the Data Base; Objective Data; Range of Motion*)
- Which of the following statements regarding the antinuclear antibody (ANA) test is FALSE?**
 - A positive ANA test alone indicates a conclusive diagnosis of lupus. (*Nursing Assessment: Establishing the Data Base; Diagnostic Studies; Serum Tests for Antibodies and Antigens*)
- According to the American Academy of Physical Medicine, ultrasound is an essential component in the diagnosis of all of the following, EXCEPT:**
 - Rheumatoid arthritis (*Nursing Assessment: Establishing the Data Base; Diagnostic Studies; Musculoskeletal Ultrasound*)
- The most common ocular symptom of Marfan syndrome is**
 - myopia. (*Specific Disorders of Connective Tissue and Muscles; Genetic disorders; Marfan Syndrome; Clinical Manifestations*)
- Patients with Ehlers-Danlos syndrome who become pregnant are at risk for**
 - uterine rupture. (*Specific Disorders of Connective Tissue and Muscles; Genetic disorders; Ehlers-Danlos Syndromes; Clinical Manifestations*)
- Which of the following should be included in the differential diagnosis of bursitis and tendonitis?**
 - All of the above (*Specific Disorders of Connective Tissue and Muscles; Inflammatory Disorders; Bursitis, Tendinitis, and Tendinosis; Clinical Manifestations*)
- Which of the following is recommended for the management of bursitis and tendinitis?**
 - Physical therapy and structured exercise after the initial period of rest (*Specific Disorders of Connective Tissue and Muscles; Inflammatory Disorders; Bursitis, Tendinitis, and Tendinosis; Therapeutic Measures*)

13. **The average age at diagnosis of polymyalgia rheumatica is**
D) 70 years of age. (Specific Disorders of Connective Tissue and Muscles; Inflammatory Disorders; Polymyalgia Rheumatica)
14. **Which type of lupus mainly affects the skin?**
D) Cutaneous lupus erythematosus (CLE) (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Lupus Erythematosus)
15. **Which of the following is thought to be the prime cause of SLE?**
D) Immune dysregulation (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Lupus Erythematosus)
16. **The most universal symptom of lupus is**
C) fatigue. (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Lupus Erythematosus; Clinical Manifestations)
17. **Which of the following drug classes is involved in the treatment of lupus?**
A) NSAIDs (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Lupus Erythematosus; Therapeutic Measures)
18. **The most frequent presentation in systemic scleroderma is the clinical triad of**
C) skin changes, Raynaud phenomenon, and esophageal hypomotility. (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Scleroderma; Clinical Manifestations)
19. **For patients with systemic scleroderma who experience digital ulcers, the recommended treatment is**
C) intravenous iloprost. (Specific Disorders of Connective Tissue and Muscles; Immunologic Disorders; Systemic Scleroderma; Therapeutic Measures)
20. **A sprain is an injury to a ligament caused by**
D) forcing a joint beyond its normal range of motion. (Specific Disorders of Connective Tissue and Muscles; Traumatic Disorders; Sprains and Strains)
21. **The most common sprains affect the**
C) ankle. (Specific Disorders of Connective Tissue and Muscles; Traumatic Disorders; Sprains and Strains; Clinical Manifestations)
22. **The standard of care for patients with rhabdomyolysis is**
B) fluid therapy. (Specific Disorders of Connective Tissue and Muscles; Traumatic Disorders; Rhabdomyolysis; Therapeutic Measures)
23. **Which of the following statements regarding psoriatic arthritis is TRUE?**
D) The manifestations of psoriatic arthritis vary from patient to patient. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Psoriatic Arthritis; Clinical Manifestations)
24. **Urate-lowering therapy should be initiated in patients with gout and**
A) one or more tophi. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Gout; Therapeutic and Nursing Measures)
25. **Which of the following is a nonmodifiable risk factor for developing low back pain?**
B) Female sex (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Low Back Pain)
26. **The majority of low back pain cases are**
A) nonspecific. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Low Back Pain; Clinical Manifestations; Nonspecific Low Back Pain)

27. Which of the following statements regarding sacroiliac (SI) joint syndrome is TRUE?
D) It typically manifests as localized pain in the lower back or upper buttock area that overlies the SI joint. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Low Back Pain; Clinical Manifestations; Sacroiliac Joint Syndrome)
28. Which of the following is considered a “red flag” when assessing the patient with chronic low back pain?
A) Prior history of cancer (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Low Back Pain; Assessment and Diagnosis)
29. Which of the following is a management goal for chronic low back pain?
D) All of the above (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Low Back Pain; Therapeutic Measures)
30. Scoliosis
D) is most commonly the result of idiopathic causes that appear with increased growth during adolescence. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Scoliosis)
31. Which of the following statements most accurately defines carpal tunnel syndrome?
B) Painful disorder of the wrist and hand that occurs when the median nerve becomes compressed (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Carpal Tunnel Syndrome)
32. As an intervention for carpal tunnel syndrome, splinting
D) may be particularly helpful when weighing the risks of surgery versus the benefits. (Specific Disorders of the Joints; Disorders of Multifactorial Origin; Carpal Tunnel Syndrome; Therapeutic and Nursing Measures)
33. The primary symptom of osteoarthritis of the knee is pain, especially
D) with weight-bearing exercise or activity, that improves with rest. (Specific Disorders of the Joints; Degenerative Disorders; Osteoarthritis; Clinical Manifestations)
34. Which of the following is TRUE about the etiology of rheumatoid arthritis?
D) Both A and B (Specific Disorders of the Joints; Immunologic Disorders; Rheumatoid Arthritis)
35. Which of the following conditions should be included in the differential diagnosis of rheumatoid arthritis?
C) Sarcoidosis (Specific Disorders of the Joints; Immunologic Disorders; Rheumatoid Arthritis; Diagnosis)
36. Which of the following best describes the treatment of rheumatoid arthritis?
D) The earlier a patient’s rheumatoid arthritis disease is diagnosed, the earlier aggressive treatment for rheumatoid arthritis can be started, thereby delaying joint destruction. (Specific Disorders of the Joints; Immunologic Disorders; Rheumatoid Arthritis; Therapeutic Measures)
37. In young, sexually active patients with infectious arthritis, the most common causative pathogen is
B) *Neisseria gonorrhoea*. (Specific Disorders of the Joints; Infectious Disorders; Infectious Arthritis; Therapeutic and Nursing Measures)
38. Hemangiomas are
C) often associated with arteriovenous malformations of skin vascular disease. (Specific Disorders of the Joints; Neoplastic Disorders; Masses and Benign Tumors of the Joint; Hemangioma)
39. Which of the following statements regarding synovial sarcoma is TRUE?
D) All of the above (Specific Disorders of the Joints; Neoplastic Disorders; Malignant Tumors of the Joint; Synovial Sarcoma)
40. Joint effusions
D) result in synovial fluid accumulation within 24 hours after a blow to the joint. (Specific Disorders of the Joints; Traumatic Disorders of the Joint; Joint Effusions)

Course Availability List

These courses may be ordered by mail on the Customer Information form located between pages 88–89. We encourage you to **GO GREEN**. Access your courses **online** or download as an **eBook** to save paper and **receive a discount** or sign up for **One Year of Unlimited Online CE starting at only \$85!** Additional titles are also available.

www.NetCE.com



Like us at NetCE4Nurses

POSTOPERATIVE COMPLICATIONS

#30763 • 15 ANCC /

1 **ADVANCED PHARM HOUR**

BOOK BY MAIL – \$98 • ONLINE – \$90

Purpose: The purpose of this course is to provide nurses and all allied health professionals who care for postsurgical patients the knowledge necessary to recognize and manage common postoperative complications, improving patient care and outcomes.

Faculty: Susan Engman Lazear, RN, MN

Audience: This course is designed for all nurses and allied professionals involved in the care of patients who undergo surgical procedures, especially those who work in the preoperative area, the operating room, or the postanesthesia unit in hospitals or free-standing surgical centers.

Additional Approval: AACN Synergy CERP Category A, CCMC



COLORECTAL CANCER

#90782 • 15 ANCC /

5 **ADVANCED PHARM HOURS**

BOOK BY MAIL – \$98 • ONLINE – \$90

Purpose: The purpose of this course is to provide healthcare professionals with information regarding the screening, diagnosis, and treatment of colorectal cancer in order to improve adherence to established guidelines and, by extension, patient outcomes.

Faculty: Mark Rose, BS, MA, LP

Audience: This course is designed for physicians, physician assistants, nurses, and other healthcare providers who may improve the identification and care of patients with colorectal cancer.

Additional Approval: AACN Synergy CERP Category A, CCMC



ACUTE CORONARY SYNDROME: AN OVERVIEW FOR NURSES

#30993 • 15 ANCC / 10 **ADVANCED PHARM HOURS**

BOOK BY MAIL – \$98 • ONLINE – \$90

Purpose: The purpose of this course is to reduce the widening gap between care according to guidelines and actual care delivered by providing nurses with knowledge necessary to implement the most appropriate approach to diagnosis and treatment.

Faculty: Karen Majorowicz, RN; Lori L. Alexander, MTPW, ELS, MWC

Audience: This course is designed for nurses practicing in primary care, inpatient, outpatient, and home care settings to enhance their knowledge of the evidence-based guidelines related to the assessment, management, and secondary prevention of acute coronary syndrome.

Additional Approval: AACN Synergy CERP Category A, CCMC

MULTIMODAL PHARMACOTHERAPY FOR PAIN MANAGEMENT

#35270 • 5 ANCC / 5 **ADVANCED PHARM HOURS**

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to provide healthcare providers with a clear understanding of the concept of multimodal pharmacotherapy for pain relief, including available classes of analgesics.

Faculty: Richard E. Haas, BSN, MSN, EdM, PhD, CRNA, PHRN

Audience: This course is designed for nurses involved in the care of patients with pain.

Additional Approval: AACN Synergy CERP Category A



HIPAA PRIVACY AND SECURITY

#91140 • 5 ANCC HOURS

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to provide information that will allow health and mental health professionals to more easily comply with the Privacy and Security Rules defined by HIPAA.

Faculty: Carol Shenold, RN, ICP

Audience: This course is designed for all members of the interprofessional healthcare team.

Additional Approval: AACN Synergy CERP Category B



SMOKING AND SECONDHAND SMOKE

#91784 • 10 ANCC /

5 **ADVANCED PHARM HOURS**

BOOK BY MAIL – \$68 • ONLINE – \$60

Purpose: The purpose of this course is to provide physicians, nurses, behavioral health professionals, and other members of the interdisciplinary team with a formal educational opportunity that will address the impact of tobacco smoking and secondhand exposure in public health and disease as well as interventions to promote smoking cessation among their patients.

Faculty: Mark S. Gold, MD, DFASAM, DLFAPA

Audience: This course is designed for physicians, nurses, and other healthcare professionals who may intervene to stop patients from smoking.

Additional Approval: AACN Synergy CERP Category A, CCMC



Prices are subject to change. Visit www.NetCE.com for a list of current prices.

Course Availability List (Cont'd)

MONKEYPOX: THE 2022 GLOBAL OUTBREAK

#94040 • 3 ANCC /

0.5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$26 • ONLINE – \$18

Purpose: The purpose of this course is to address these knowledge gaps to enable timely diagnosis, treatment, and prevention of monkeypox, thereby promoting public health strategies to limit spread of the outbreak.

Faculty: John M. Leonard, MD

Audience: This course is designed for physicians, physician assistants, nurses, pharmacy professionals, and other healthcare professionals who may identify and care for patients with suspected or confirmed human monkeypox infection.

Additional Approval: AACN Synergy CERP Category A

NEW!

HYPERTENSION: STRATEGIES TO IMPROVE OUTCOMES

#94223 • 5 ANCC / 5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to provide healthcare professionals with the information necessary to develop treatment regimens associated with optimal adherence and provide adequate patient education, counseling, and support to patients with hypertension.

Faculty: John J. Whyte, MD, MPH

Audience: This course is designed for all physicians, physician assistants, nurses, and pharmacy professionals involved in the care of patients with hypertension.

Additional Approval: AACN Synergy CERP Category A, CCMC

UPDATE

OSTEOARTHRITIS

#94954 • 10 ANCC /

5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$68 • ONLINE – \$60

Purpose: The high prevalence of osteoarthritis and its substantial burden at both the individual and healthcare system levels demands sound knowledge and clinical skills in diagnosing and managing the disease. The purpose of this course is to provide healthcare professionals with the information necessary to adequately assess osteoarthritis symptoms, treat osteoarthritis patients based on evidence-based guidelines, and appropriately refer to specialists.

Faculty: Lori L. Alexander, MTPW, ELS, MWC

Audience: This course is designed for physicians, physician assistants, nurses, and other healthcare professionals involved in the care of patients with osteoarthritis.

Additional Approval: AACN Synergy CERP Category A, CCMC

New Guidelines

OPIOID SAFETY: BALANCING BENEFITS AND RISKS

#95500 • 5 ANCC / 5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to provide clinicians who prescribe or distribute opioids with an appreciation for the complexities of opioid prescribing and the dual risks of litigation due to inadequate pain control and drug diversion or misuse in order to provide the best possible patient care and to prevent a growing social problem.

Faculty: Mark Rose, BS, MA, LP

Audience: This course is designed for all physicians, osteopaths, physician assistants, pharmacy professionals, and nurses who may alter prescribing and/or dispensing practices to ensure safe opioid use.

Additional Approval: AACN Synergy CERP Category A

Special Approval: This course fulfills the 2-hour requirement for pain management, identification of addiction, or the practices of prescribing or dispensing opioids for Pennsylvania CRNPs with prescriptive authority.

NEW!
Pain Mgmt/
Opioids

PSYCHEDELIC MEDICINE AND INTERVENTIONAL PSYCHIATRY

#96790 • 10 ANCC /

8 ADVANCED PHARM HOURS

BOOK BY MAIL – \$68 • ONLINE – \$60

Purpose: The purpose of this course is to provide medical and mental health professionals with the knowledge and skills necessary to effectively treat mental disorders using emerging psychedelic and interventional techniques.

Faculty: Mark S. Gold, MD, DFASAM, DLFAPA

Audience: The course is designed for all members of the interprofessional team, including physicians, physician assistants, nurses, and mental health professionals, involved in caring for patients with mental disorders resistant to traditional treatment approaches.

Additional Approval: AACN Synergy CERP Category A

NEW!

CANNABIS AND CANNABIS USE DISORDERS

#96973 • 5 ANCC / 5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to allow healthcare professionals to effectively identify, diagnose, treat, and provide appropriate referrals for patients with cannabis use disorders.

Faculty: Mark Rose, BS, MA, LP

Audience: This course is designed for health and mental health professionals who are involved in the evaluation or treatment of persons who use cannabis, either illicitly or as an adjunct to medical treatment.

Additional Approval: AACN Synergy CERP Category A

Prices are subject to change. Visit www.NetCE.com for a list of current prices.

Course Availability List (Cont'd)

PALLIATIVE CARE AND PAIN MANAGEMENT AT THE END OF LIFE

#97383 • 15 ANCC / 10 ADVANCED PHARM HOURS

BOOK BY MAIL – \$98 • ONLINE – \$90

Purpose: The purpose of this course is to bridge the gap in knowledge of palliative care by providing an overview of the concept of palliative care and a discussion of the challenges, benefits, and strategies of optimum palliative care at the end of life.

Faculty: Lori L. Alexander, MTPW, ELS, MWC

Audience: This course is designed for all members of the interprofessional team, including physicians, physician assistants, nurse practitioners, nurses, pharmacists, pharmacy technicians, social workers, marriage and family therapists, and other members seeking to enhance their knowledge of palliative care.

Additional Approval: AACN Synergy CERP Category A, CCMC

COMMONLY ABUSED SUPPLEMENTS

#98020 • 2 ANCC HOURS

BOOK BY MAIL – \$23 • ONLINE – \$15

Purpose: The purpose of this course is to provide healthcare professionals in all practice settings the knowledge necessary to increase their understanding of the commonly abused supplements and their adverse effects.

Faculty: Chelsey McIntyre, PharmD

Audience: This course is designed for healthcare professionals whose patients are taking or are interested in taking dietary supplements.

Additional Approval: AACN Synergy CERP Category A

GETTING TO THE POINT: ACUPUNCTURE AND ACUPOINT THERAPIES

#98030 • 4 ANCC HOURS

BOOK BY MAIL – \$32 • ONLINE – \$24

Purpose: The purpose of this course is to provide healthcare professionals in all practice settings the knowledge necessary to increase their understanding of acupoint and acupressure therapies.

Faculty: Chelsey McIntyre, PharmD

Audience: This course is designed for healthcare professionals whose patients are using or are interested in using acupoint and acupressure therapies.

Additional Approval: AACN Synergy CERP Category A

DIZZINESS AND VERTIGO

#98401 • 10 ANCC / 5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$68 • ONLINE – \$60

Purpose: The purpose of this course is to provide clinicians with the information necessary to appropriately diagnose and treat causes of dizziness and vertigo and improve patients' quality of life.

Faculty: Mark Rose, BS, MA, LP

Audience: This course is designed for physicians and nurses involved in the diagnosis, treatment, and care of patients with dizziness and/or vertigo.

Additional Approval: AACN Synergy CERP Category A, CCMC

PARKINSON DISEASE

#98772 • 10 ANCC /

5 ADVANCED PHARM HOURS

BOOK BY MAIL – \$68 • ONLINE – \$60

Purpose: The purpose of this course is to provide physicians, nurses, and other members of the interprofessional healthcare team a review of pathogenesis, disease progression, diagnosis, and management of Parkinson disease, in order to improve patient care and quality of life.

Faculty: Mark Rose, BS, MA, LP

Audience: This course is designed for all healthcare providers in the primary care setting who may encounter patients with Parkinson disease.

Additional Approval: AACN Synergy CERP Category A, CCMC

UPDATE

ANEMIA IN THE ELDERLY

#99083 • 5 ANCC / 2 ADVANCED PHARM HOURS

BOOK BY MAIL – \$38 • ONLINE – \$30

Purpose: The purpose of this course is to provide healthcare providers with the knowledge and tools necessary to identify anemia early and respond appropriately. Better health outcomes for the geriatric population will result from an increase in evidence-based clinical practices.

Faculty: Susan Waterbury, MSN, FNP-BC, ACHPN

Audience: This course is designed for physicians, physician assistants, nurses, and other healthcare professionals involved in the care of elderly patients.

Additional Approval: AACN Synergy CERP Category A, CCMC

Are you licensed in other states?
Our Unlimited Online Special Offer includes all state mandates!
www.NetCE.com/NurseUnlimited

Prices are subject to change. Visit www.NetCE.com for a list of current prices.



Customer Information

(Incomplete information may delay processing.)

For office use only:
PA23

Please print your Customer ID # located on the back of this catalog. (Optional)

--	--	--	--	--	--	--	--	--	--	--	--

Order/complete by mail
P.O. Box 997571
Sacramento, CA 95899-7571

Order/complete by fax
(916) 783-6067

Contact us
(800) 232-4238

Email us
help@NetCE.com

Order/complete online
www.NetCE.com/PA23

Last Name _____ First Name _____ MI _____

State _____ License # _____ Exp. _____

State _____ Additional License # _____ Exp. _____

License Type (circle one): RN / CNS / CRNP / CRNA / CNM / Other: _____

Address _____

City _____ State _____ Zip _____

Phone () _____

Fax () _____

Email _____

Sign me up for NetCE Alerts, Promotions, License Reminders, and More!

Receive certificate(s) by:

Online Access - FREE! Email required

Email - FREE!

Fax - FREE!

Mail - Add \$6 for shipping and handling

<p>Complete before December 31, 2023, pay</p> <p>\$52</p> <p>Complete after December 31, 2023, pay</p> <p>\$62</p>	<p>ENCLOSED SPECIAL OFFER: 31 HOURS</p> <p>You may complete ALL four of these courses for a maximum payment of \$52 (or pay the individual course price).</p>																					
	<table border="1"> <thead> <tr> <th>✓</th> <th>Course #</th> <th>Course Title / Contact Hours</th> <th>Price</th> </tr> </thead> <tbody> <tr> <td> </td> <td>97542</td> <td>Child Abuse Identification & Reporting: The Pennsylvania Req. / 3 Contact Hours</td> <td>\$18</td> </tr> <tr> <td> </td> <td>96690</td> <td>Anxiety Disorders in Older Adults / 3 Contact Hours</td> <td>\$18</td> </tr> <tr> <td> </td> <td>90240</td> <td>Pancreatic Cancer / 10 Contact Hours</td> <td>\$60</td> </tr> <tr> <td> </td> <td>38950</td> <td>Pathophysiology: Muscles, Joints, and Connective Tissues / 15 Contact Hours</td> <td>\$90</td> </tr> </tbody> </table>	✓	Course #	Course Title / Contact Hours	Price		97542	Child Abuse Identification & Reporting: The Pennsylvania Req. / 3 Contact Hours	\$18		96690	Anxiety Disorders in Older Adults / 3 Contact Hours	\$18		90240	Pancreatic Cancer / 10 Contact Hours	\$60		38950	Pathophysiology: Muscles, Joints, and Connective Tissues / 15 Contact Hours	\$90	
✓	Course #	Course Title / Contact Hours	Price																			
	97542	Child Abuse Identification & Reporting: The Pennsylvania Req. / 3 Contact Hours	\$18																			
	96690	Anxiety Disorders in Older Adults / 3 Contact Hours	\$18																			
	90240	Pancreatic Cancer / 10 Contact Hours	\$60																			
	38950	Pathophysiology: Muscles, Joints, and Connective Tissues / 15 Contact Hours	\$90																			

Additional Courses Available by Mail (ACCESS ONLINE FOR A DISCOUNT!)
 Payment must accompany this form. To order by phone, please have your credit card ready.

✓	Course #	Course Title / Contact Hours	Price	✓	Course #	Course Title / Contact Hours	Price
<input type="checkbox"/>	30763	Postoperative Complications / 15.....	\$98	<input type="checkbox"/>	95500	Opioid Safety: Balancing Benefits and Risks / 5	\$38
<input type="checkbox"/>	30993	Acute Coronary Syndrome / 15	\$98	<input type="checkbox"/>	96790	Psychedelic Medicine & Interventional Psychiatry / 10 .	\$68
<input type="checkbox"/>	35270	Multimodal Pharmacotherapy for Pain Mgmt / 5	\$38	<input type="checkbox"/>	96973	Cannabis and Cannabis Use Disorders / 5	\$38
<input type="checkbox"/>	90782	Colorectal Cancer / 15	\$98	<input type="checkbox"/>	97383	Palliative Care and Pain Management at the EOL / 15..	\$98
<input type="checkbox"/>	91140	HIPAA Privacy and Security / 5.....	\$38	<input type="checkbox"/>	98020	Commonly Abused Supplements / 2	\$23
<input type="checkbox"/>	91784	Smoking and Secondhand Smoke / 10.....	\$68	<input type="checkbox"/>	98030	Getting to the Point: Acupuncture & Acupoint Therapies / 4..	\$32
<input type="checkbox"/>	94040	Monkeypox: The 2022 Global Outbreak / 3	\$26	<input type="checkbox"/>	98401	Dizziness and Vertigo / 10	\$68
<input type="checkbox"/>	94223	Hypertension: Strategies to Improve Outcomes / 5.....	\$38	<input type="checkbox"/>	98772	Parkinson Disease / 10	\$68
<input type="checkbox"/>	94954	Osteoarthritis / 10	\$68	<input type="checkbox"/>	99083	Anemia in the Elderly / 5	\$38

Check or Money Order (payable to NetCE)

VISA / MasterCard / AmEx / Discover

Please print name (as shown on credit card) _____

Credit card # _____

Expiration date _____ Security code _____

Signature _____

Security code is last three numbers in the signature area on back of credit card or four numbers above the account number on front of AmEx cards.

Special Offer (before December 31, 2023) **\$52**

\$62 (after December 31, 2023) _____

I would like my certificates mailed for an additional \$6 _____

Additional Courses _____

Subtotal _____

Expedited Delivery _____

Grand Total _____

Expedited mail delivery within 2 to 3 days is available in most areas at an additional charge of \$35. Call for information on international delivery.

Prices are subject to change. Visit www.NetCE.com for a list of current prices.

Last Name _____ First Name _____ MI _____
 State _____ License # _____ Expiration Date _____

To receive continuing education credit, completion of this Evaluation is mandatory.
Please answer all of the following questions and provide your signature at the bottom of this page.
Your postmark or facsimile date will be used as your completion date.

Please read the following questions and choose the most appropriate answer for each course completed.

1. Was the course content new or review?
2. How many self-assessment questions did you answer correctly? _____ Questions
3. How much time did you spend on this activity, including the questions? _____ Hours
4. Would you recommend this course to your peers?
5. Did the course content support the stated course objective?
6. Did the course content demonstrate the author's knowledge of the subject?
7. Was the course content free of bias?
8. Before completing the course, did you identify the necessity for education on the topic to improve your professional practice?
9. Have you achieved all of the stated learning objectives of this course?
10. Has what you think or feel about this topic changed?
11. Were self-assessment questions with evaluation an effective teaching strategy?
12. Did evidence-based practice recommendations assist in determining the validity or relevance of the information?
13. Are you more confident in your ability to provide patient care after completing this course?
14. Do you plan to make changes in your practice as a result of this course content?

In accordance with the reporting requirements of Act 31, please provide the following information for course #97542:

15. Please provide the last four digits of your social security number.
16. Please provide your date of birth.

#97542
 PA Child Abuse
 Identification and Reporting
 3 Contact Hours

1. New Review
2. _____ Questions
3. _____ Hours
4. Yes No
5. Yes No
6. Yes No
7. Yes No
8. Yes No
9. Yes No
10. Yes No
11. Yes No
12. Yes No
13. Yes No
14. Yes No
15. _____
16. ____/____/____
(mm/dd/yyyy)

#96690
 Anxiety Disorders
 in Older Adults
 3 Contact Hours

1. New Review
2. _____ Questions
3. _____ Hours
4. Yes No
5. Yes No
6. Yes No
7. Yes No
8. Yes No
9. Yes No
10. Yes No
11. Yes No
12. Yes No
13. Yes No
14. Yes No

#90240
 Pancreatic
 Cancer
 10 Contact Hours

1. New Review
2. _____ Questions
3. _____ Hours
4. Yes No
5. Yes No
6. Yes No
7. Yes No
8. Yes No
9. Yes No
10. Yes No
11. Yes No
12. Yes No
13. Yes No
14. Yes No

#38950
 Pathophysiology: Muscles,
 Joints, & Connective Tissues
 15 Contact Hours

1. New Review
2. _____ Questions
3. _____ Hours
4. Yes No
5. Yes No
6. Yes No
7. Yes No
8. Yes No
9. Yes No
10. Yes No
11. Yes No
12. Yes No
13. Yes No
14. Yes No

Additional Comments / Criticisms / Concerns: _____

I have read the course(s) and completed the Evaluation(s) in full. I understand my postmark or facsimile date will be used as my completion date.

Signature _____
Signature required to receive continuing education credit.

NOW AVAILABLE UNLIMITED ONLINE CE FOR ONE YEAR STARTING AT ONLY \$85

Purchase our Unlimited Special Offer and receive access to more than 1,500 hours of CE, including special offers and state-required courses.

Anxiety Disorders	Dizziness and Vertigo
Ischemic Stroke	Sepsis
Responsible Opioid Prescribing	Autism Disorder
Herbal Medications	Alcohol Use Disorder
Diabetes Pharmacology	Fibromyalgia
Influenza	...and many more!
Parkinson Disease	

To access your Special Offer, please visit:
NetCE.com/NurseUnlimited

 **NetCE**
a trchealthcare brand



 **NatMed**
a trchealthcare brand

Access the Most Comprehensive Database for Alternative Medicines

Answer patient questions on natural medicines and integrative therapies with confidence.

Scan the QR code
to learn more.



Want More CE Choices?

Get One Year of Unlimited Online CE starting at only \$85*!

Includes access to our entire course library of more than 1,500 hours, including over 500 pharmacology hours, special offers, and state-required courses!

The following Online Specials are included with your Online Unlimited Subscription or may be purchased individually.

NURSES:

NATURAL MEDICINE SPECIAL OFFER

Cannabinoids • Top-Selling Supplements • Commonly Abused Supplements • Acupuncture • Microbiome Medley • Glucosamine and Chondroitin • The Scoop on Collagen
17.5 Hours \$49

WOUND MANAGEMENT SPECIAL OFFER

Pressure Injuries and Skin Care • Hyperglycemia and Wound Management • Treating Pressure Injuries and Chronic Wounds
15 Hours \$41

TRAUMA SPECIAL OFFER

Postoperative Complications • Transport Methods for the Critically Ill Patient
30 Hours \$62

ADVANCED PRACTICE NURSES:

DESIGN A DEAL SPECIAL OFFER

Choose from our entire library of courses to create your own unique Special Offer.

\$52 to \$73 (Based on number of hours selected)
[NetCE.com/Design](https://www.netce.com/Design)

ADVANCED PRACTICE SPECIAL OFFER

Bacterial Sexually Transmitted Infections • Diagnosing and Managing Headaches • Autoimmune Diseases
30 Hours (includes 22 Pharm Hours) \$73

PHARMACOLOGY SPECIAL OFFER

Epidural Analgesia Update • Diabetes Pharmacology • Moderate Sedation/Analgesia
30 Hours (includes 28 Pharm Hours) \$73

***APRNs, get One Year of Unlimited Online CE for \$125. Includes over 500 Pharmacology hours.**

For more details, go to [NetCE.com/NurseUnlimited](https://www.netce.com/NurseUnlimited)

BUSINESS HOURS: Monday through Friday, 8am-5pm Pacific Time. We are closed on weekends and holidays.

CUSTOMER SERVICE: 800-232-4238 or help@netce.com. Call or email us for customer assistance, course catalogs, additional certificates, or transcripts. If you require special assistance, please contact the Sr. Director of Development and Academic Affairs to inform her of your needs by calling 800-232-4238.

RETURN POLICY: Satisfaction guaranteed or your money back within 30 days of purchase, unless certificates have been issued. Please return the materials and include a brief note of explanation. For more information, please contact help@netce.com.


TURNAROUND TIME: If sent by mail, your order is processed within 2 to 3 weeks from the day it was received. **For the fastest processing time, visit www.netce.com to purchase, complete for credit, and receive your certificates instantly.**

MAILING PREFERENCES: To modify your mailing preferences or to view our privacy policy, please go to www.netce.com.

PRICING: Prices are subject to change. Visit www.netce.com for a list of current prices.

RETURNED CHECKS: If, for any reason, your check is returned, you will be contacted requesting a cashier's check or money order for the full amount of the order plus a \$35 reinstatement fee. In addition, we are unable to accept temporary checks.

If you have questions about your license or certification renewal or state requirements, please contact your board. A list of approvals and accreditations is available on our website at www.netce.com.



NatMed
a trchealthcare brand

Access the Most Comprehensive Database for Alternative Medicines

Answer patient questions on natural medicines and integrative therapies with confidence.

Scan the QR code to learn more.





P.O. Box 997571
Sacramento, CA 95899

PRESORTED
STANDARD
U.S. POSTAGE PAID
PERMIT NO. 182
SACRAMENTO, CA

Vol. 148
No. 20
PA23

Complete online at NetCE.com/PA23

Quick Code#

Customer ID#

Scan this QR code to get started.
If you do not have a smartphone or
QR code reader, please visit
NetCE.com/PA23.



Like us at **NetCEContinuingEducation**



We Recycle

